



2025 Herbicide x Variety Guide

The purpose of this guide is to assist you in managing herbicide programs with Pioneer® brand soybeans. Pioneer uses molecular markers, lab, and/or field testing to evaluate soybean variety tolerance to several herbicides.

- Research has shown good correlation between molecular markers and varietal response to **preplant or pre-emergence applications** of the PPO herbicides sulfentrazone and saflufenacil, but low correlation with response to the PPO herbicide flumioxazin (e.g. Afforia®, Enlite®, Envive®, Trivence®, Surveil® and Valor®) when the herbicides are used at normal field rates.
- Research has also shown good correlation between lab assays and field tolerance to preplant and preemergence applications of metribuzin.
- Research has also shown good correlation between molecular markers and tolerance to preplant or preemergence applications of rimsulfuron.

Please note that these ratings are not correlated with tolerance to exposure or application of these herbicides after soybean emergence. Metribuzin, rimsulfuron, and all PPO herbicides can injure soybeans when applied after emergence. Crop injury can also occur when metribuzin or PPO-treated soils are splashed onto soybean stems, cotyledons, or foliage.

Challenging environments such as heavy rainfall during seed germination or seedling emergence; sandy soils, soils low in organic matter or high pH soils; or during periods of excessively cold, hot, dry or wet weather can result in higher herbicide activity or reduced crop tolerance. In such cases, crop injury may occur on varieties rated as having acceptable tolerance to the herbicide. University research indicates herbicides within an herbicide class may vary in their degree of crop selectivity. The potential for herbicide injury may also be impacted by the labeled herbicide rate used and the method or timing of application.

Herbicides that contain **Sulfentrazone and Saflufenacil** include Spartan® brands, Authority® brands, Sonic®, Optill®, Optill® PRO, Sharpen® and Verdict®. **Always read and follow herbicide label directions.**

Herbicides that contain **Metribuzin** include Canopy® Blend herbicide, Trivence® herbicide, Sencor®, Axiom®, Boundary®, Domain® and Authority® MTZ. **Always read and follow herbicide label directions.**

The following herbicide sensitivity ratings are for sulfentrazone, saflufenacil, and metribuzin:

- **Adequate Tolerance.** Available research and/or field observations suggest this herbicide is unlikely to result in material crop injury to this particular variety under normal circumstances.
- ▼ **Requires Careful Management.** Available research and/or field observations suggest this herbicide may exhibit crop injury to this particular variety in challenging environments.
- **Response Warning.** Available research and/or field observations suggest this herbicide has a high potential for crop injury to this variety.
- **Insufficient Data.** Additional testing is needed to evaluate this variety.

Herbicides that contain **Rimsulfuron** include LeadOff® and Basis® Blend. **Always read and follow herbicide label directions.**

The following herbicide sensitivity ratings are for rimsulfuron:

- ++++ **Varieties with BOLT® technology.** Growers may apply LeadOff® or Basis® Blend herbicides 0 days or more prior to planting this particular variety.
- +++ **Varieties with the STS® gene.** This particular variety has a shorter plant-back interval for LeadOff® and Basis® Blend herbicides. See product labels for details on plant-back intervals.
- ++ **High degree of rimsulfuron tolerance.** Available research and/or field observations suggest these herbicides are unlikely to result in material crop injury to this particular variety under normal circumstances. See product labels for details on plant-back intervals.
- + **Low degree of rimsulfuron tolerance.** Available research and/or field observations suggest these herbicides have a high potential for crop injury to this particular variety. Do not plant this particular variety into rimsulfuron-treated fields within 10 months of application if soil is excessively cold or wet or if soil pH exceeds 6.5. Soil temperature should be >50° F and trending warmer. See product labels for details on plant-back intervals.
- **Insufficient Data.** Additional testing is needed to evaluate this variety.



2025 Herbicide x Variety Guide

Pioneer makes no warranty regarding the herbicide crop response information in this guide. Other varieties may be added when data become available.

			Herbicide Sensitivity Ratings		
Brand**	Relative Maturity ¹	Technology Segment ²	Sulfentrazone and Saflufenacil	Metribuzin	Rimsulfuron
P0009A28E™	000.9	E3	●	●	++
P002A42E™	00.2	E3	●	●	++
P003Z08E™	00.3	E3	●	●	++
P004Z87E™	00.4	E3	●	●	++
P006Z63E™	00.6	E3	●	●	++
P007A68E™	00.7	E3	●	●	++
P007Z45E™	00.7	E3	●	●	++
P007Z82E™*	00.7	E3	●	●	++
P008Z25E™	00.8	E3	●	●	++
P009Z94E™	00.9	E3	●	●	++
P00Z65E™*	0.0	E3	●	●	++
P01Z13E™	0.1	E3	●	●	++
P02A78E™	0.2	E3	●	●	++
P02Z34E™	0.2	E3	●	●	++
P04A98E™	0.4	E3	●	●	++
P04Z49E™	0.4	E3	●	●	++
P05Z60E™	0.5	E3	●	●	++
P06A38E™	0.6	E3	●	●	++
P06A85E™	0.6	E3	●	●	++
P06Z90E™	0.6	E3	●	●	++
P07A10™	0.7	-	●	●	++
P08A44E™	0.8	E3	●	●	++
P09A20™	0.9	-	■	●	+
P09Z31E™*	0.9	E3	●	●	++
P09Z79E™	0.9	E3	●	●	++
P11A10™	1.1	-	●	▼	++
P11A50™	1.1	-	●	●	++
P11A67	1.1	-	□	●	+
P11T36E™	1.1	E3	●	●	++
P11Z72E™	1.1	E3	●	●	++
P11Z91E™*	1.1	E3	●	●	++
P13A03™	1.3	-	●	●	++
P13Z28E™	1.3	E3	●	●	++
P14Z08E™*	1.4	E3	●	●	++
P14Z16E™	1.4	E3	●	●	++
P14Z67E™	1.4	E3	●	●	++

Adequate Tolerance
 Requires Careful Management
 Crop Response Warning
 Insufficient Data



2025 Herbicide x Variety Guide

			Herbicide Sensitivity Ratings		
Brand**	Relative Maturity ¹	Technology Segment ²	Sulfentrazone and Saflufenacil	Metribuzin	Rimsulfuron
P15A20™	1.5	-	●	●	++
P15Z80E™	1.5	E3	●	●	++
P16Z25E™*	1.6	E3	●	●	++
P16Z42E™	1.6	E3	●	●	++
P16Z96E™	1.6	E3	●	●	++
P17A87E™	1.7	E3	●	●	++
P17Z39E™	1.7	E3	●	●	++
P18A73E™	1.8	E3	●	●	++
P18A82™	1.8	-	●	●	++
P18Z01E™	1.8	E3	●	●	++
P19A27PR™	1.9	Plenish,R	●	■	++
P19A37E™	1.9	E3	●	●	++
P19A66E™	1.9	E3	●	●	++
P19Z52E™	1.9	E3	●	●	++
P20A10™	2.0	-	●	●	++
P20Z14E	2.0	E3	●	●	++
P21A20™	2.1	-	●	▼	++
P21A31PR™	2.1	Plenish,R	●	●	++
P21Z05PE™*	2.1	Plenish,E3	●	●	++
P21Z71E™	2.1	E3	●	●	++
P21Z88E™	2.1	E3	●	●	++
P22A36PR™	2.2	Plenish,R	●	●	++
P22A67E™	2.2	E3	●	●	++
P22Z02E™*	2.2	E3	●	●	++
P22Z50™*	2.2	-	●	●	++
P23Z58E™	2.3	E3	●	●	++
P23Z82E™	2.3	E3	●	●	++
P24A07E™	2.4	E3	●	●	++
P24A46PR™	2.4	Plenish,R	●	●	++
P25A16E™	2.5	E3	●	●	++
P25Z11E™*	2.5	E3	●	●	++
P26A20™	2.6	-	●	●	++
P26Z78E™	2.6	E3	●	●	++
P26Z86E™*	2.6	E3	●	●	++
P26Z94PE™*	2.6	Plenish,E3	●	●	++
P27A26PR™	2.7	Plenish,R	●	●	++
P27Z41E™	2.7	E3	●	●	++
P28A39E™	2.8	E3	●	●	++

Adequate Tolerance
 Requires Careful Management
 Crop Response Warning
 Insufficient Data



2025 Herbicide x Variety Guide

			Herbicide Sensitivity Ratings		
Brand**	Relative Maturity ¹	Technology Segment ²	Sulfentrazone and Saflufenacil	Metribuzin	Rimsulfuron
P28A83PR™	2.8	Plenish,R	●	●	++
P28Z30E™	2.8	E3	●	●	++
P28Z44PE™*	2.8	Plenish,E3	●	●	++
P28Z89E™	2.8	E3	●	●	++
P29A57PR™	2.9	Plenish,R	●	●	++
P29T50	2.9	-	●	▼	++
P29Z61E™*	2.9	E3	●	●	++
P29Z96PE™*	2.9	Plenish,E3	●	●	++
P30A75E™	3.0	E3	●	●	++
P30A93PR™	3.0	Plenish,R	●	●	++
P30Z35PE™*	3.0	Plenish,E3	●	●	++
P31A48PR™	3.1	Plenish,R	●	●	++
P31Z03E™	3.1	E3	●	●	++
P31Z32E™	3.1	E3	●	●	++
P32A10™	3.2	-	●	●	++
P32Z91E™	3.2	E3	●	●	++
P33A62E™	3.3	E3	●	●	++
P33A85E™	3.3	E3	●	●	++
P33T60	3.3	-	●	●	++
P33Z17E™	3.3	E3	●	●	++
P33Z27E™*	3.3	E3	●	●	++
P34A50™	3.4	-	●	●	++
P34A59PR™	3.4	Plenish,R	●	●	++
P34A65PR™	3.4	Plenish,R	●	●	++
P34A98E™	3.4	E3	●	●	++
P35A20™	3.5	-	●	●	++
P35Z76E™	3.5	E3	●	●	++
P36A94PR™	3.6	Plenish,R	●	●	++
P36Z04E™*	3.6	E3	●	●	++
P36Z47BE™	3.6	Bolt,E3	●	□	++++
P37A18E™	3.7	E3	●	●	++
P37T51PR	3.7	Plenish,R	●	●	++
P37Z06E™	3.7	E3	●	●	++
P37Z90™*	3.7	-	●	●	++
P38A28E™	3.8	E3	●	●	++
P38Z41E™*	3.8	E3	●	●	++
P38Z63E™	3.8	E3	●	●	++
P39A78™	3.9	-	●	●	++

Adequate Tolerance
 Requires Careful Management
 Crop Response Warning
 Insufficient Data



2025 Herbicide x Variety Guide

			Herbicide Sensitivity Ratings		
Brand**	Relative Maturity ¹	Technology Segment ²	Sulfentrazone and Saflufenacil	Metribuzin	Rimsulfuron
P39A82S™	3.9	STS	■	▼	+++
P40A40™	4.0	-	●	●	++
P40Z57E™	4.0	E3	●	●	++
P41A61BE™	4.1	Bolt,E3	●	●	++++
P41T65PR	4.1	Plenish,R	■	●	++++
P41Z11PE™*	4.1	Plenish,E3	●	●	++
P41Z16E™*	4.1	E3	●	●	++
P41Z80BLX™	4.1	Bolt,LL,RR2X	●	●	++++
P42A13PR™	4.2	Plenish,R	●	●	++
P42A84E™	4.2	E3	●	●	++
P42Z50BLX™*	4.2	Bolt,LL,RR2X	●	●	++++
P43Z44SE™	4.3	STS,E3	■	●	++++
P44A91E™	4.4	E3	●	●	++
P44Z67BE™*	4.4	Bolt,E3	●	●	++++
P45A40LX™	4.5	LL,RR2X	●	●	++
P45A81E™	4.5	E3	●	●	++
P45Z75E™	4.5	E3	●	▼	++
P46A09E™	4.6	E3	■	●	+
P46A35X™	4.6	RR2X	■	●	+
P46A45PR™	4.6	Plenish,R	●	●	++
P46A90LX™	4.6	LL,RR2X	●	●	++
P46T97SE™	4.6	STS,E3	■	●	+++
P46Z26E™*	4.6	E3	●	●	++
P46Z53E™	4.6	E3	■	●	+
P47A10™	4.7	-	●	▼	++
P47Z15BE™	4.7	Bolt,E3	●	●	++++
P48A04LX™	4.8	LL,RR2X	●	●	++
P48A14E™	4.8	E3	●	●	++
P48A94PR™	4.8	Plenish,R	●	●	++
P48Z58PE™*	4.8	Plenish,E3	●	●	++
P48Z70BLX™	4.8	Bolt,LL,RR2X	●	●	++++
P48Z73E™*	4.8	E3	●	●	++
P49Z02E™	4.9	E3	■	●	+
P50A08LX™	5.0	LL,RR2X	●	●	++
P50Z95E™	5.0	E3	■	●	+
P51A85BE™	5.1	Bolt,E3	●	●	++++
P51Z31SE™*	5.1	STS,E3	●	●	++
P52A14SE™	5.2	STS,E3	■	●	+++

Adequate Tolerance
 Requires Careful Management
 Crop Response Warning
 Insufficient Data



2025 Herbicide x Variety Guide

			Herbicide Sensitivity Ratings		
Brand**	Relative Maturity ¹	Technology Segment ²	Sulfentrazone and Saflufenacil	Metribuzin	Rimsulfuron
P53Z60LX™	5.3	LL,RR2X	●	●	++
P56A71E™	5.6	E3	■	●	+
P56Z22E™*	5.6	E3	●	●	++
95Y70	5.7	STS,R	■	▼	+++
P59Z15SE™*	5.9	STS,E3	■	●	+++
P60Z06E™*	6.0	E3	●	●	++
P64Z52E™	6.4	E3	●	●	++
P68A41BE™	6.8	Bolt,E3	●	▼	++++
P69Z71E™*	6.9	E3	●	●	++
P70A62E™	7.0	E3	●	▼	++

* New Product

** All Pioneer products denoted with ™ are brand names.

1 RELATIVE MATURITY: Shows the relative maturity group rating, with the digits preceding the decimal representing the general maturity group, and the digit following the decimal showing relative maturity within the group on a scale of 0 to 9, with 0 early and 9 late. For example, a soybean product with a relative maturity rating of 1.8 would be a late product in Group 1 maturity.

2 TECHNOLOGY SEGMENT:



Always follow stewardship practices in accordance with the Product Use Guide (PUG) or other product-specific stewardship requirements including grain marketing and pesticide label directions. **Varieties with BOLT® technology** provide excellent plant-back flexibility for soybeans following application of sulfonylurea (SU) herbicides such as LeadOff® or Basis® Blend as a component of a burndown program or for double-crop soybeans following SU herbicides such as Finesse® applied to wheat the previous fall.

Glyphosate Tolerant

Always follow grain marketing, stewardship practices and pesticide label directions. **Varieties with the Glyphosate Tolerant trait** (including those designated by the letter "R" in the product number) contain genes that confer tolerance to glyphosate herbicides. Glyphosate herbicides will kill crops that are not tolerant to glyphosate.

STS®

herbicide tolerant trait

Varieties with the STS® trait are tolerant to certain sulfonylurea (SU) herbicides. This technology allows post-emergent applications of Synchrony® XP and Classic® herbicides without crop injury or stress (see herbicide product labels). NOTE: A soybean variety with a herbicide tolerant trait does not confer tolerance to all herbicides. Spraying herbicides not labeled for a specific soybean variety will result in severe plant injury or plant death. Always read and follow herbicide label directions and precautions for use.



Varieties with the LibertyLink® gene (LL) are resistant to Liberty® herbicide.

LibertyLink® and the Water Droplet Design are registered trademarks of Bayer.

● Adequate Tolerance ▼ Requires Careful Management ■ Crop Response Warning □ Insufficient Data



2025 Herbicide x Variety Guide



DO NOT APPLY DICAMBA HERBICIDE IN-CROP TO SOYBEANS WITH Roundup Ready 2 Xtend® (RR2X) technology unless you use a dicamba herbicide product that is specifically labeled for that use in the location where you intend to make the application. IT IS A VIOLATION OF FEDERAL AND STATE LAW TO MAKE AN IN-CROP APPLICATION OF ANY DICAMBA HERBICIDE PRODUCT ON SOYBEANS WITH Roundup Ready 2 Xtend® technology, OR ANY OTHER PESTICIDE APPLICATION, UNLESS THE PRODUCT LABELING SPECIFICALLY AUTHORIZES THE USE. Contact the U.S. EPA and your state pesticide regulatory agency with any questions about the approval status of dicamba herbicide products for in-crop use with soybeans with Roundup Ready 2 Xtend® technology. ALWAYS READ AND FOLLOW PESTICIDE LABEL DIRECTIONS. Soybeans with Roundup Ready 2 Xtend® technology contain genes that confer tolerance to glyphosate and dicamba. Glyphosate herbicides will kill crops that are not tolerant to glyphosate. Dicamba will kill crops that are not tolerant to dicamba. Roundup Ready 2 Xtend® is a registered trademark of Monsanto Technology LLC used under license.



Varieties with Enlist E3® technology (E3): The transgenic soybean event in Enlist E3® soybeans is jointly developed and owned by Corteva Agriscience and M.S. Technologies L.L.C.

Following burndown, Enlist Duo® and Enlist One® herbicides with Colex-D® technology are the only herbicides containing 2,4-D that are authorized for preemergence and postemergence use with Enlist® crops. Consult Enlist® herbicide labels for weed species controlled. Enlist Duo and Enlist One herbicides are not registered for use or sale in all states and counties; are not registered in AK, CA, CT, HI, ID, MA, ME, MT, NH, NV, OR, RI, UT, VT, WA and WY; and have additional subcounty restrictions in AL, GA, TN and TX, while existing county restrictions still remain in FL. All users must check "Bulletins Live! Two" no earlier than six months before using Enlist One or Enlist Duo. To obtain "Bulletins," consult epa.gov/espp/, call 1-844-447-3813, or email ESPP@epa.gov. You must use the "Bulletin" valid for the month and state and county in which Enlist One or Enlist Duo are being applied. Contact your state pesticide regulatory agency if you have questions about the registration status of Enlist® herbicides in your area. ALWAYS READ AND FOLLOW PESTICIDE LABEL DIRECTIONS. IT IS A VIOLATION OF FEDERAL AND STATE LAW TO USE ANY PESTICIDE PRODUCT OTHER THAN IN ACCORDANCE WITH ITS LABELING. ONLY USE FORMULATIONS THAT ARE SPECIFICALLY LABELED FOR SUCH USE IN THE STATE OF APPLICATION. USE OF PESTICIDE PRODUCTS, INCLUDING, WITHOUT LIMITATION, 2,4-D-CONTAINING PRODUCTS NOT AUTHORIZED FOR USE WITH ENLIST CROPS, MAY RESULT IN OFF-TARGET DAMAGE TO SENSITIVE CROPS/AREAS AND/OR SUSCEPTIBLE PLANTS, IN ADDITION TO CIVIL AND/OR CRIMINAL PENALTIES. Additional product-specific stewardship requirements for Enlist crops, including the Enlist Product Use Guide, can be found at www.traitstewardship.com.



Plenish® (P) high oleic soybeans have an enhanced oil profile and are produced and channeled under contract to specific grain markets. Growers should refer to the Pioneer Product Use Guide on www.pioneer.com/us/stewardship for more information.

(-) = Variety does not contain a herbicide resistant gene.

PIONEER® brand products are provided subject to the terms and conditions of purchase which are part of the labeling and purchase documents.™® Trademarks of Corteva Agriscience and its affiliated companies. © 2025 Corteva.

REV. 1/16/25

