

Pellicar-F

- Used to compensate for deficiencies in exogenous amino acids (histidine, tryptophan, threonine)
- It is a nutritionally incomplete food and cannot be used as the sole source of nourishment
- Must be used under medical supervision
- Intended for adults
- Should not be used by pregnant women, breastfeeding women or in case of hypersensitivity to any of the product ingredients
- Comorbidities (e.g., diabetes, kidney failure) are not a contraindication to use
- Not intended for parenteral use
- Does not pose a health risk for people without pressure sores or hard-healing wounds

Storage:

Store in a dry place, at a temperature below 25°C. Keep out of the reach of small children.

PelliCarnos

Food for special medical purposes

Pellicar-F

For dietary management
in the treatment of pressure sores
or hard-healing wounds

Carnosine, threonine, tryptophan

30 tablets

Pellicar-F

Dosage and method of use:

1 tablet once a day or 1 tablet 2 times a day during or after a meal.

The tablets can be chewed, swallowed whole or allowed to dissolve in the mouth.

Do not exceed the recommended daily dose.

Do not use in case of hypersensitivity to the ingredients of the preparation.

The product is not intended for children, pregnant or breastfeeding women.

Contains a sweetener (izomalt E953). The product does not contain gluten, lactose, and preservatives.

Batch No.:

Best before the end:

Net weight: 20.7g (30 tablets of 0.69g each)

Manufactured for: PelliCarnos S.A.
ul. Zajęcza 15, 00-351 Warsaw
www.pellicarnos.com