



Midodine



POVIDONE IODINE 10% SOLUTION (1% ACTIVE IODINE)

Preoperative and antiseptic skin
treatment of burns to reduce
the risk of infecti

Ready to use

Lower irritant effect

Wide application

Forms a protective, germicidal film

Effective in the treatment of burns

Wide spectrum of exposure
(including HIV, HBV, Herpes)



120 ml

500 ml

1000 ml

5000 ml

MIDO PHARM®

www.midopharm.com

Midodine



Midodine is a professional produkt for pre-operative skin and burn antiseptic treatment to reduce the risk of the most popular infections.

Can be used for hygienic and surgical hand disinfection, urological, ophthalmological and oral treatment. It forms a protective, germicidal film over the treated area, stimulates wound healing, is harmless to mucous membranes and soft tissues. Destroys microorganisms: viruses, bacteria, fungi, spores, protozoa. The development of antibacterial resistance to iodinated povidone has not been observed during more than 60 years of intensive use in hospitals, dentistry and doctors' practices. Furthermore, antibiotic resistance does not affect sensitivity to povidone-iodine.

In the povidone complex, iodine has a significantly lower irritating effect compared to alcohol solutions of iodine.

APPLICATION:

Midodine is intended for use before operations, gynecological and obstetric manipulations, bladder catheterization, biopsies, injections, punctures, blood sampling, treatment of minor burns and insect stings, prevention of oral inflammation, treatment of surgical instruments, emergency disinfection after contact with infected material and surfaces.

For use in hospitals, polyclinics, health centers, nursing homes, social centers and other health care institutions.

Clinically tested. Microbiologically tested. For non-professional and professional use. Keep out of the sight and reach of children!

USAGE

For external use only. Do not put in hot water and heat only to about body temperature. Used on the skin in undiluted and diluted form. The diluted solution should be prepared shortly before use. The presence of large amounts of blood may reduce the effectiveness of the product.

A diluted solution can be used before oral and ophthalmological manipulations, as well as for disinfection of drinking water.

USE OF UNDILUTED SOLUTION

For hygienic hand disinfection.

2x3ml of undiluted solution with an exposure time of 30 sec each time.

For surgical hand disinfection.

2x5ml of undiluted solution with an exposure time of 5 minutes.

Disinfection of the skin and skin bordering with genital area.

Before applying, make sure that the surface of the skin is clean. Apply the undiluted solution to the skin, completely covering the area to be treated, let it work for at least 60 seconds, let it dry.

For skin and scar care.

Use on the treated surface 2-3 times a day. After application, tie up.



MIDO PHARM®

MANUFACTURER:

SPODRA SIA
VAT: LV40003941352
Lubanas street 17-1,
Riga, Latvija, LV-1019,
Tel.: +371 29624524
E-mail: rikra@sveiks.lv

www.spodra.com



www.midopharm.com



USE OF DILUTED SOLUTION

INDICATION	Dilution	Examples
Wet compresses	1:5 to 1:10	200 to 100 ml per 1 liter
Immersion or sitz bath	1:25	40 ml to 1 liter
Preoperative bath, stomatitis, pharyngitis, oral moniliasis and trichomoniasis	1:100	10 ml per 1 liter
Hygienic bath, drinking water disinfection	1:1000	10 ml per 10 liters
Vaginal douching, ITS insertion, perineal douching, urological douching	1:25	4 ml per 100 ml
Irrigation of chronic and postoperative wounds	1:2, 1:20	50 ml to 5 ml per 100 ml
Rinsing in traumatology and orthopedics, rinsing in oral cavity surgery	1:10	10 ml per 100 ml
Use in ophthalmology (pre-surgery)	1:8	12,5 ml per 100 ml (physiological solution)

USAGE

Do not use to disinfect open wounds with a large area. May cause skin irritation or allergic skin reactions if there is an allergy to povidone-iodine, iodine or if there are thyroid diseases. If irritation or allergic reaction is felt, use should be stopped.

Special care is required for pregnant women and patients who are breastfeeding. In such cases, the benefit/risk ratio should be evaluated and povidone iodine should be used only in case of absolute indications.

Attention! It is not recommended for use in children under 2 years of age and is contraindicated in newborns and children under 1 year of age. For newborns or infants up to 6 months of age, use only in case of absolute indications.

Do not use together with chlorhexidine and other active substances to prevent mutual inactivation.

BENEFITS OF USING

- Fast effect
- Long lasting action
- Broad-spectrum antiseptic effect
- Alleviation of pain syndrome in burns
- Excellent skin compatibility, pH level similar to physiological

1 ml of solution contains 100 mg of iodized povidone, which corresponds to 10 mg of active iodine, glycerin, sodium hydroxide, sodium benzoate.

AVAILABLE PACKAGING TYPES

UNITS	PACKAGING	ARTICLE
120 ml bottle with cap	24 pcs	008714
500 ml bottle with cap	12 pcs	008912
1000 ml bottle with cap	12 pcs	008721
5000 ml can	1 pcs	008738

ACTIVE INGREDIENT

Polyvinylpyrrolidone iodine (CAS No.: 25655-41-8): 10 g / 100 g (10 wt %)



MANUFACTURER:

SPODRA SIA
VAT: LV40003941352
Lubanas street 17-1,
Riga, Latvija, LV-1019,
Tel.: +371 29624524
E-mail: rikra@sveiks.lv
www.spodra.com

