

DATE: 6/18/14 6:20/14 6/23/14

ORDER NUMBER:

Please examine carefully for style, arrangement, completeness of copy, spelling and color. It is expressly understood that all artwork, special dies, plates, etc., manufactured by or for us in connection with this order, shall remain permanently in our possession, and any payment made by buyer for or on account of such material shall be deemed part of the consideration for our services in completing this order.

We attempt to ensure this label layout conforms to any applicable labeling laws or regulations. However, it is the client's ultimate responsibility to obtain proper final approval.

218.963.2575 • 800.247.6573 • 218.963.4849 FAX

NISSWA, MN 56468

P.O. BOX 10 • 5652 COUNTY ROAD 18

LABEL & PACKAGING INC

VIKING

VIKINGLABEL.COM

Label Size: 5.375 X 8.625

Die: 1076

4 Color Process

BLACK LINE

I state this proof is:

☐ OK as is.

☐ In need of changes and reproof.

☐ OK for production **AFTER** changes.

Authorized Customer Signature.

We cannot proceed with your order until a returned, SIGNED proof is received.

If you have any questions regarding this proof, please contact your Viking Label representative.

HAR00043

PANTONE® Check standard Pantone Color Specifier for exact color match.

COLORS ON THIS PROOF MAY NOT REPRESENT FINAL PRINTED LABEL

Packaged/Shipped:

per Roll/Box:

per Ctn:

Hard Core:

Rewind Figure:

1

THIS WAY

2

THIS WAY

3

THIS WAY

4

THIS WAY

Feeding Directions:

1. Before Feeding EquiLix for the first time, animals should be well fed and have access to plenty of clean, fresh water.

2. Have one lick block available for every five to ten head of horses.

3. Provide an adequate source of good quality forage at all times (all they can eat).

4. Place lick blocks in stalls or in pastures where horses loaf.

5. Make lick blocks available in more than one location in large pastures or where more than one water source is available.

Ingredients: Corn distillers dried grains with solubles, Corn condensed distillers solubles, Yeast culture, Salt, Monocalcium Phosphate, Diatomaceous Earth, Flax Meal, Pea Meal, Magnesium Oxide, Calcium Carbonate, Canola Oil, Calcium Sulfate, Potassium Chloride, Zinc Sulfate, Manganese Sulfate, Copper Sulfate, Zinc Polysaccharide Complex, Zinc Proteinate, Manganese Polysaccharide Complex, Manganese Proteinate, Copper Polysaccharide Complex, Copper Proteinate, Dried Aspergillus Oryzae Fermentation Extract, Liquid Aspergillus Niger Fermentation Product, Cobalt Carbonate, Ethylenediamine Dihydriodide, Vitamin A Acetate (Stability Improved), Vitamin D3 Supplement, Vitamin E Supplement, Biotin.

Control #:

Best If Used Before:

Guaranteed by:
Harvest Fuel, Inc. Walhalla, ND 58282

This product manufactured in:
Walhalla, ND 58282
Horton, KS 66437
800-327-9222 / www.equilix.com

is a registered trademark of harvest Fuel, Inc.

US Patent #5,260,089 which deals with new, advanced alternatives to molasses licks applies to this product.

Guaranteed Analysis

Crude Protein, Min.....	13.00%	Copper (Cu), Min.....	600 ppm
Crude Fat, Min.....	6.00%	Selenium (Se), Min.....	5.0 ppm
Omega 3 Fatty Acid, Min.....	0.40%	Zinc (Zn), Min.....	1200 ppm
Crude Fiber, Max.....	11.00%	Manganese (Mn), Min.....	750 ppm
Calcium (Ca), Min.....	2.50%	Iodine (I), Min.....	8.0 ppm
Calcium (Ca), Max.....	3.00%	Cobalt (Co), Min.....	4.0 ppm
Phosphorus (P), Min.....	1.50%	Fluorine, Max.....	0.8 ppm
Salt (NaCl), Min.....	12.00%	Vitamin A, Min.....	100,000 IU/lb
Salt (NaCl), Max.....	14.00%	Vitamin D, Min.....	20,000 IU/lb
Magnesium (Mg), Min.....	2.50%	Vitamin E, Min.....	350 IU/lb
Potassium (K), Min.....	1.10%	Biotin, Min.....	18 ppm
Sulfur (S), Min.....	1.00%	Omega 3 Fatty Acid, Min.....	5500 ppm

Alpha-Amylase (from Aspergillus oryzae)..... Min. 10,180 ug starch hydrolyzed/min/oz.
Amyloglucosidase (from Aspergillus niger, Aspergillus oryzae)..... Min. 1,540 ug maltose hydrolyzed/min/oz.

Net Wt. 125 lbs. (57 kg.)

6. If animals run out of lick blocks, repeat from Step #1.

7. Other salt and minerals may be provided free choice to balance specific rations.

8. A two week adaptation period will be required to establish a base consumption rate.

9. The above feeding directions are general guidelines and will vary with the overall feeding conditions.

Always use good management practices and adapt to your specific requirements.

WARNING: Contains 600 ppm copper.
DO NOT feed to sheep!

Made
in the U.S.A.