

FEEDING DIRECTIONS:

- 1. Before feeding EquiLix PI for the first time, animals should be well fed and have access to plenty of clean, fresh water.
- 2. Have one lick block available for every five to ten head of horses.
- 3. Provide an adequate source of good quality forage at all times (all they can eat).
- 4. Place lick blocks in stalls or in pastures where horses loaf.
- 5. Make lick blocks available in more than one location in large pastures or where more than one water source is available.
- 6. If animals run out of lick blocks, repeat from Step #1.
- 7. Other salt and minerals may be provided free choice to balance specific rations.
- 8. A two week adaptation period will be required to establish a base consumption rate.
- 9. The above feeding directions are general guidelines and will vary with the overall feeding conditions.

Always use good management practices and adapt to your specific requirements.

WARNING: Contains 500 ppm copper. DO NOT feed to sheep!

www.SweetPro.com











Vitamin, Mineral & Digestive Aid



Control #:

Best If Used Before:

Guaranteed by: Harvest Fuel, Inc., Walhalla, ND 58282

This product manufactured in:
Walhalla, ND 58282 ____
Horton, KS 66437 ____
800-327-9222 / www.sweetpro.com



is a registered trademark of Harvest Fuel, Inc.


US Patent #5,260,089 which deals with new, advanced alternatives to molasses licks applies to this product.

INGREDIENTS: Corn distillers dried grains with solubles, condensed corn distillers solubles, yeast culture, salt, ground flax, pea meal, diatomaceous earth, magnesium oxide, monocalcium phosphate, calcium carbonate, vegetable oil, potassium chloride, copper proteinate, zinc proteinate, manganese proteinate, copper sulfate, zinc sulfate, manganese sulfate, cobalt carbonate, ethylenediamine dihydriodide, sodium selenite, dried aspergillus oryzae fermentation extract, liquid aspergillus niger fermentation product, vitamin A acetate, vitamin D3 supplement, vitamin E supplement, biotin.

Guaranteed Analysis

Crude Protein, Min14.00%	Copp
Crude Fat, Min 6.50%	Selei
Omega 3 Fatty Acid, Min 0.40%	Zinc
Crude Fiber, Max 11.00%	Iron
ADF, Max 8.50%	Man
Calcium (Ca), Min 1.75%	lodir
Calcium (Ca), Max 2.25%	Coba
Phosphorus (P), Min 1.00%	Fluo
Salt (NaCl), Min 7.00%	Vitar
Salt (NaCl), Max 9.00%	Vitar
Magnesium (Mg), Min 2.00%	Vitar
Potassium (K), Min 1.10%	Bioti
Sulfur (S), Min	Ome

Copper (Cu), Min	500 ppm
Selenium (Se), Min	5.0 ppm
Zinc (Zn), Min	1000 ppm
Iron (Fe), Min	350 ppm
Manganese (Mn), Min	750 ppm
lodine (I), Min	8.0 ppm
Cobalt (Co), Min	4.0 ppm
Fluorine, Max	0.8 ppm
Vitamin A, Min	100,000 IU/lb
Vitamin D, Min	20,000 IU/lb
Vitamin E, Min	350 IU/lb
Biotin, Min	18 ppm
Omega 3 Fatty Acid, Min	5500 ppm



Made

in the USA

Digestive Enzyme Activity (at time of manufacture)

Alpha-amylase (from barley malt, aspergillus oryzae)......10,180 ug starch hydrolyzed/min/oz Amyloglucosidase (from aspergillus niger, aspergillus oryzae)...1,540 ug maltose hydrolyzed/min/oz

Net Wt. 125 lbs (57 kg)