1	
2	
3	
4	
5	
6	
7	
8	
9	
10	
11	
12	
13	
14	
15	

# Name:

# LINEAR ALGEBRA AND VECTOR ANALYSIS

MATH 22B

Total:

Welcome to the final exam. Please don't get started yet. We start all together at 9:00 AM after getting reminded about some formalities. You can fill out the attendance slip already. Also, you can already enter your name into the larger box above.

- You only need this booklet and something to write. Please stow away any other material and any electronic devices. Remember the honor code.
- Please write neatly and give details. Except for problems 2 and 3 we want to see details, even if the answer should be obvious to you.
- Try to answer the question on the same page. There is additional space on the back of each page. If you must, use additional scratch paper at the end.
- If you finish a problem somewhere else, please indicate on the problem page where we can find it.
- You have 180 minutes for this 3-hourly.



# Unit 39: Final Practice B

#### Problems

### Problem 39B.1) (10 points):

The graph G = (V, E) in Figure 1 represents a discrete surface in which all triangles are oriented counterclockwise. The values of a 1-form = vector field F are given.

a) (2 points) Find the line integral of F along the boundary curve oriented counter clockwise.

b) (2 points) Compute the curl H = dF and write its values into the triangles.

c) (2 points) What is the sum of all curl values? Why does it agree with the result in a)?

d) (2 points) Find also  $g = d^*F$  and enter it near the vertices.

e) (1 point) True or False:  $\sum_{x \in V} g(x) = 0.$ 

f) (1 point) True of False: we called  $L = dd^*$  the Laplacian of G.



FIGURE 1. A discrete 2-dimensional region on which a 1-form F models a vector field. You compute the curl dF and divergence  $d^*F$  of F.

Problem 39B.2) (10 points) Each question is one point:

a) Name the 3-dimensional analogue of the Mandelbrot set.

b) If A is a  $5 \times 4$  matrix, then  $A^T$  is a  $m \times n$  matrix. What is m and n?

c) Write down the general formula for the arc length of a curve

$$r(t) = [x(t), y(t), z(t)]^T$$

with  $a \leq t \leq b$ .

d) Write down one possible formula for the curvature of a curve

$$r(t) = [x(t), y(t), z(t)]^T$$
.

e) We have seen a parametrization of the 3-sphere invoking three angles  $\phi, \theta_1, \theta_2$ . Either write down the parametrization or recall the name of the mathematician after whom it this parametrization is named.

f) The general change of variable formula for  $\Phi : R \to G$  is  $\iiint_R f(u, v, w)$   $dudvdw = \iiint_G f(x, y, z) dxdydz$ . Fill in the blank part of the formula.

g) What is the numerical value of  $\log(-i)$ ?

h) We have used the Fubini theorem to prove that  $C^2$  functions f(x, y) satisfy a partial differential equation. Please write down this important partial differential equation as well as its name. (It was used much later in the course.)

i) What is the integration factor |dr| for the parametrization

$$r(u, v) = [a\cos(u)\sin(v), b\sin(u)\sin(v), c\cos(v)]^T$$
?

j) In the first lecture, we have defined  $\sqrt{\operatorname{tr}(A^T A)}$  as the length of a matrix. What is the length of the  $3 \times 3$  matrix which contains 1 everywhere?

# Problem 39B.3) (10 points) Each problem is 1 point:

a) Assume that for a Morse function f(x, y) the discriminant D at a critical point  $(x_0, y_0)$  is positive and that  $f_{yy}(x_0, y_0) < 0$ . What can you say about  $f_{xx}(x_0, y_0)$ ?

b) We have proven the identity  $|dr| = |r_u \times r_v|$ , where r was a map from  $\mathbb{R}^m$  to  $\mathbb{R}^n$ . For which m and n was this identity defined?

c) Which of the following is the correct integration factor when using spherical coordinates in 4 dimensions?

 $\begin{aligned} |d\Phi| &= r\\ |d\Phi| &= (3 + \cos(\phi))\\ |d\Phi| &= \rho^2 \sin(\phi)\\ |d\Phi| &= \rho^3 \sin(2\phi)/2 \end{aligned}$ 

d) Which of the following vector fields are gradient fields? (It could be none, one, two, three or all.)

$$\begin{split} F &= [x,0]^T \\ F &= [0,x]^T \\ F &= [x,y]^T \\ F &= [y,x]^T \end{split}$$

e) Which of the following four surfaces is a one-sheeted hyperboloid? (It could be none, one, two, three or all.)

 $\begin{array}{l} x^2+y^2=z^2-1\\ x^2-y^2=1-z^2\\ x^2+y^2=1-z^2\\ x^2-y^2=z^2+1 \end{array}$ 

f) Parametrize the surface  $x^2 + y^2 - z^2 = 1$  as

 $r(\theta, z) = [\dots, \dots, \dots, \dots]^T.$ 

g) Who was the creative person who discovered dark matter and proposed the mechanism of gravitational lensing?

h) What is the cosine of the angle between the matrices  $A, B \in M(2, 2)$ , where A is the identity matrix and B is the matrix which has 1 everywhere? You should get a concrete number.

i) We have seen the identity  $|v|^2 + |w|^2 = |v - w|^2$ , where v, w are vectors in  $\mathbb{R}^n$ . What conditions do v and w have to satisfy so that the identity holds?

j) Compute the exterior derivative dF of the differential form  $F = e^x \sin(y) dx dy + \cos(xyz) dy dz .$ 

#### Problem 39B.4) (10 points):

a) (4 points) Find the plane  $\Sigma$  which contains the three points

A = (3, 2, 1), B = (3, 3, 2), C = (4, 3, 1).

b) (3 points) What is the area of the triangle ABC?

c) (3 points) Find the distance of the origin O = (0,0,0) to the plane  $\Sigma$ .

#### Problem 39B.5) (10 points):

a) (8 points) Find all the critical points of the function

$$f(x,y) = x^5 - 5x + y^3 - 3y$$

and classify these points using the second derivative test.

b) (2 points) Is any of these points a global maximum or global minimum of f?

### Problem 39B.6) (10 points):

a) (8 points) Use the Lagrange method to find **all the maxima and all the minima** of

$$f(x,y) = x^2 + y^2$$

under the constraint

$$g(x,y) = x^4 + y^4 = 16$$
.

b) (2 points) In our formulation of Lagrange theorem, we also mentioned the case, where  $\nabla g(x, y) = [0, 0]^T$ . Why does this case not lead to a critical point here?

# Problem 39B.7) (10 points):

a) (5 points) The hyper surface

$$S = \{f(x, y, z, w) = x^2 + y^2 + z^2 - w = 5\}$$

defines a three-dimensional manifold in  $\mathbb{R}^4$ . It is poetically called a **hyper-paraboloid**. Find the tangent plane to S at the point (1, 2, 1, 1).

b) (5 points) What is the linear approximation L(x, y, z, w) of f(x, y, z, w) at this point (1, 2, 1, 1)?

Linear Algebra and Vector Analysis

# Problem 39B.8) (10 points):

Estimate the value f(0.1, -0.02) for

$$f(x,y) = 3 + x^{2} + y + \cos(x+y) + \sin(xy)$$

using quadratic approximation.

## Problem 39B.9) (10 points):

a) (8 points) We vacation in the **5-star hotel** called **MOTEL 22** in 5-dimensional space and play there ping-pong. The ball is accelerated by gravity  $r''(t) = [x(t), y(t), z(w), v(t), w(t)] = [0, 0, 0, 0, -10]^T$ . We hit the ball at  $r(0) = [4, 3, 2, 1, 2]^T$  and give it an initial velocity  $r'(0) = [5, 6, 0, 0, 3]^T$ . Find the trajectory r(t).

b) (2 points) At which positive time t > 0 does the ping-pong ball hit the **hyper ping-pong table** w = 0? (The points in this space are labeled [x, y, z, v, w].)

# Problem 39B.10) (10 points):

a) (5 points) Integrate the function  $f(x, y) = (x^2 + y^2)^{22}$  over the region  $G = \{1 < x^2 + y^2 < 4, y > 0\}.$ 

b) (5 points) Find the area of the region enclosed by the curve

$$r(t) = [\cos(t), \sin(t) + \cos(2t)]^T$$
,

with  $0 \le t \le 2\pi$ .

# Problem 39B.11) (10 points):

a) (7 points) Integrate

$$f(x, y, z) = x^2 + y^2 + z^2$$

over the solid

$$G = \{x^2 + y^2 + z^2 \le 4, z^2 < 1\}.$$

b) (3 points) What is the volume of the same solid G?

#### Problem 39B.12) (10 points):

a) (8 points) Compute the line integral of the vector field

$$F = [yzw + x^{6}, xzw + y^{9}, xyw - z^{3}, xyz + w^{4}]^{T}$$

along the path

$$r(t) = [t + \sin(t), \cos(2t), \sin(4t), \cos(7t)]^T$$

from t = 0 to  $t = 2\pi$ .

b) (2 points) What is  $\int_0^{2\pi} r'(t) dt$ ?

## Problem 39B.13) (10 points):

a) (8 points) Find the line integral of the vector field

 $F(x,y) = [3x - y, 7y + \sin(y^4)]^T$ 

along the polygon ABCDE with A = (0,0), B = (2,0), C = (2,4), D = (2,6), E = (0,4). The path is closed. It starts at A, then reaches B, C, D, E until returning to A again.

b) (2 points) What is line integral if the curve is traced in the opposite direction?

# Problem 39B.14) (10 points):

a) (8 points) What is the flux of the vector field

 $F(x, y, z) = [y + x^3, z + y^3, x + z^3]^T$ 

through the sphere  $S = \{x^2 + y^2 + z^2 = 9\}$  oriented outwards?

b) (2 points) What is the flux of the same vector field F through the same sphere S but where S is oriented inwards?

#### Problem 39B.15) (10 points):

a) (7 points) What is the flux of the curl of the vector field

$$F(x, y, z) = [-y, x + z(x^{2} + y^{5}), z]^{T}$$

through the surface

$$S = \{x^2 + y^2 + z^2 + z(x^4 + y^4 + 2\sin(x - y^2 z)) = 1, z > 0\}$$

oriented upwards?

b) (3 points) The surface in a) was not closed, it did not include the bottom part

$$D = \{ z = 0, x^2 + y^2 \le 1 \} .$$

Assume now that we close the bottom and orient the bottom disc D downwards. What is the flux of the curl of the same vector field F through this closed surface obtained by taking the union of S and D?