



Pediatric Dermatology

Epidemiology and Clinical Pattern of Atopic Dermatitis in a North Indian Pediatric Population

	<b>Study objective</b>	To study the incidences and clinical patterns of AD in Pediatric population
	<b>No. of subjects</b>	462 children with AD, 241 (52.16%) were boys and 221 (47.83%) were girls.
	<b>Duration</b>	Mean age at onset of the disease was 4.1 years and mean duration was 1.9 years
	<b>Test Method</b>	In every patient a detailed history was taken, including present age, age at onset, residence, personal and family history of atopy, degree of pruritus, history of oozing, seasonal variation, and history of allergy and associated diseases, if any.

Result:

Approximately **2% of the AD patients** had an aggravation during rainy season.<sup>1</sup>


Rise in rainfall is linked to the rise in the concentration of mold spores in the environment.<sup>2</sup>

1. Pediatric Dermatology Vol. 15 No. 5 347-35 1, 1998 | 2. Osong Public Health Res Perspect 2013 4(3), 140-145

In Chronic Dermatitis Conditions  
<sup>Rx</sup>**AVEXA**<sup>®</sup>  
Cream/Lotion



The **Emollient +plus**



**CARBOMER**  
TECHNOLOGY



QUICK ABSORPTION



MOISTURE LOCK



Non-comedogenic



Dermatologically Tested

Soothing & instant cooling

Strengthens skin barrier

Suitable for sensitive skin

💡 Think **Moisturizer...** Think **Ajanta Dermatology**