

Ministry of Health and Social Welfare

What is vaccination?


Vaccination is way people are protected against diseases by giving special medicine called vaccines

What diseases do vaccination prevent people against ?

- TB
- Diphtheria
- Poliomyelitis
- Pertusis
- Tetanus
- Measles
- Yellow Fever
- Hepatitis
- Haemophilus Influenza

VITAMIN A SUPPLEMENTATION DOSE: 100,000 UNITS 6-11 MONTHS 200,000 UNITS PAST PARTUM WOMEN (TO BE GIVEN EVERY 6 MONTHS)			
DATE OF VISIT	DOSE	DATE OF VISIT	DOSE

S/N	DATE TO RETURN TO CLINIC
1	
2	
3	
4	
5	



Sleep under you're your mosquito net every night.

When Pregnant, Go to the clinic for your Malaria Medicine

ANTI-TUBERCULOSIS VACCINATION

Date of BCG Vaccination _____
Give at birth

POLIO VACCINE Date vaccine Given

OPV0 _____

OPV1 _____

OPV2 _____

OPV3 _____

PENTAVALENT Date vaccine Given

PENTA -1 _____

PENTA -2 _____

PENTA -3 _____

MEASLES VACCINE DATE: _____
Give at 9 Months to less than 12 Months

YELLOW FEVER VACCINE DATE: _____
Give at 9 Months to less than 12 Months

Republic of Liberia

CHILD'S No.	
CHILD'S Name	M or F
FATHER'S Name	
MOTHER'S Name	
ADDRESS	
CLINIC ISSUING CARD	Date first seen

BIRTH NOTIFICATION	
PLACE OF BIRTH	WEIGHT
DATE OF BIRTH	TIME
NAME OF BIRTH ATTENDANT:	
FACILITY NAME:	

When and how should children take their vacci-

Vaccine	age	site	dose	route
BCG	At birth	Upper right arm	0.5ml	intradermal
OPV0	At birth	oral	2 drops	mouth
OPV1	6 wks	oral	2 drops	mouth
OPV2	10 wks	oral	2 drops	mouth
OPV3	14 wks	oral	2 drops	mouth
Penta 1	6 wks	Upper left thigh	0.5ml	intramuscular
Penta 2	10 wks	Upper left thigh	0.5ml	intramuscular
Penta 3	14 wks	Upper left thigh	0.5ml	intramuscular
measles	9 months	Upper left arm	0.5ml	subcutaneous
Yellow fever	9 months	Upper left arm	0.5ml	subcutaneous


Watch the direction of the line showing the child health



GOOD
Mean the child is growing well



Danger
Mean the child is not growing well



Very Dangerous
May be ill needs extra care

- Breast milk is the best milk for your baby
- Give baby only breast milk from birth to six months
- From six months add other foods according to the child's age
- Continue breast feeding up to two years

