## **Supplement Facts**

Serving Size: 3 Capsules Servings Per Container: 60

Amount Per Serving		% DV
D-Aspartic Acid	1560mg	*
Fenusterols® (Fenugreek Seed Extract) (50% Steroidal Saponins)	300mg	*
Tribulus Terrestris (40% Saponins)	300mg	*
Horny Goat Weed (10% Icariin)	100mg	*
Maca Root	100mg	
Eurycoma Longifolia 20:1	100mg	*

Boron (as Boron Citrate) \* Daily Value not established

Other Ingredients: Gelatin, Silicon Dioxide

BioPerine®

The statements made hereon have not been evaluated by the Food and Drug Administration. This product is not intended to diagnose, treat, cure, or prevent any disease

Dist. by: Advantage Pharmaceuticals Orem, UT 84057 U.S.A. ©2013 Advantage Pharmaceuticals, LLC

MADE IN USA

5ma



**NEW FORMULA** 

## ALPHA

NATURAL TESTOSTERONE COMPLEX

WARNING: KEEP OUT OF REACH OF CHILDREN. Not intended for use by persons under 18 years of age. Do not exceed recommended use. Improper use will not improve results and is not advised. Consult with your physician before using this product. Do not take this product if you have a medical condition(s) including, but not limited to diabetes, heart disease, high or low blood pressure, cardiac arrhythmia, stroke, liver or thyroid disease, psychological disorders, prostate hypertrophy, or hormonal abnormalities.

> 180 CAPSULES DIETARY SUPPLEMENT

## RECOMMENDED USE:

Take 3 capsules twice daily. For best results take first thing in the morning on an empty stomach, and before bedtime.

WARNING: Not intended for use by persons under 18 years of age. Do not exceed recommended use. Improper use will not improve results and is not advised. Consult with your physician before using this product. Do not take this product if you have a medical condition(s) including, but not limited to diabetes, heart disease, high or low blood pressure. cardiac arrhythmia, stroke, liver or thyroid disease, psychological disorders, prostate hypertrophy, or hormonal abnormalities. Consult with your physician if you are taking antidepressants, MAO inhibitors or other prescription drugs. Discontinue use two weeks prior to surgery. This product should not be used by women, those prone to dehydration, or if you are regularly exposed to excessive heat. Reduce or discontinue use if sleeplessness, tremors, dizziness, nervousness, headaches, or heart palpitations occur. Do not use if packaging has been tampered with. Store in a cool, dry place.

KEEP OUT OF REACH OF CHILDREN.