

INDEX

Note: Page numbers followed by the letter f refer to figures. Page numbers followed by the letter t refer to tables. Page numbers followed by the letter p refer to plates.

ABCs of trauma management, 2072–2073
Abdomen, injuries to, with pelvic fractures, 2256
Abdominal muscles, external oblique, for iliopsoas transfer, 1336
Abdominal reflex, 56
assessment of, in scoliosis, 223, 223f
Abduction. *See also under specific joint(s)*, e.g., Hip.
definition of, 27t
Abduction contracture(s)
metacarpal, in cerebral palsy, 1222f–1223f
of hip, Ober's test for, 40, 41f
Abductor hallucis brevis tendon, transfer to first metatarsal, for dorsal bunion, 1372–1373, 1379f, 1386f–1387f
Abductor limp
after varus femoral osteotomy, 697, 700f
in avascular necrosis, 583
Abductor pollicis brevis tendon, preservation of, for radial polydactyly, 426, 430f–431f
Above-knee amputation, 1793–1794
ischial-bearing, for osteosarcoma, 1982p–1991p
prostheses for, 1806
Above-knee orthosis, for poliomyelitis, 1326–1327
Abscess(es)
Brodie's, 155f
cold, in tuberculous arthritis, 1832–1833, 1833f
intraosseous, 1848, 1851f
of psoas muscle, 1850f
paravertebral, in tuberculous spondylitis, 1832–1833, 1833f
Abuse. *See Child abuse*.
Accessory bone(s), in foot, 895f
Accessory navicular. *See Navicular, accessory*.
Acetabular index, in developmental dysplasia of hip, 531, 532f, 533t
Acetabular teardrop figure, in developmental dysplasia of hip, 531–532, 533f
Acetabulum
anatomy of, 519, 519f–520f
augmentation of
osteotomies for, 642–646, 644f–649f
Chiari, 642, 644f, 646
shelf procedures for, 645–649, 646f
development of, in developmental dysplasia of hip, 520f, 520–522, 522f
dysplasia of
after treatment for developmental dysplasia of hip, 587–596, 619f
and developmental dysplasia of hip, 517

Acetabulum (*Continued*)
fractures of, 2251–2271. *See also Pelvis*, fractures of.
complications of, 2271
treatment of, 2268–2271, 2270f–2272f
with hip dislocation, 2276f
growth of, 2251, 2252f
ossification into labrum, after use of Pavlik harness, 537, 538f
radiographic changes in, in Legg-Calvé-Perthes disease, 679
repositioning of, pelvic osteotomies for, 596–632, 618t, 621f–633f, 622p–641p
Ganz, 633, 642, 643f
Pemberton, 596–597, 618, 620f, 622p–629p
Steel, 632, 642f
Tonnis, 632–633
Acetaminophen, postoperative, 112
Acetylcholine receptor antibody(ies), in myasthenia gravis, 1500
Achard syndrome, vs. Beals' syndrome, 1595
Achilles tendon
anterior transposition of, for equinus deformity, in cerebral palsy, 1137
contracture of, and pes planovalgus, 912
dissection of, in correction of vertical talus, 968p–969p
lengthening of
for equinus deformity, in cerebral palsy, 1136–1137, 1137f–1139f, 1140p–1141p
for pes cavus, 991, 995f
for talipes equinovarus, 931–932, 937, 938f
tenodesis of, to fibula, for valgus ankle deformity, 1268
Achondroplasia, 1510–1519, 1511f–1519f and kyphosis, 346
and postlaminectomy kyphosis, 340f
angular deformities of lower extremities in, 1517, 1518f
clinical features of, 1510–1512, 1511f–1512f
craniocervical stenosis in, 1513–1514
diagnosis of, prenatal, 1513
differential diagnosis of, 1513
genetic features of, 1510
growth hormone for, 1517, 1519f
leg lengthening for, 1517, 1519f
pathophysiology of, 1510–1512
radiographic features of, 1512–1513, 1512f–1513f
short stature in, 1517, 1519, 1519f
spinal stenosis in, 1516f, 1516–1517

Achondroplasia (*Continued*)
thoracolumbar kyphosis in, 1514f–1515f, 1514–1516
vs. osteogenesis imperfecta, 1727–1728
vs. pseudoachondroplasia, 1524t
Achterman and Kalamchi classification, of fibular deficiency, 1776, 1776f
ACL. *See Anterior cruciate ligament (ACL)*.
Acrocephalosyndactyly syndrome(s), 1615
anesthesia in, 120t
Acromioclavicular joint
dislocation of, 2118, 2120
injuries to, Rockwood classification of, 2115, 2117f
physeal fractures of, 2116f
Acromion process, partial excision of, in shoulder disarticulation, 2016p–2017p
Acro-osteolysis, in pyknodysostosis, 1564
Acrosyndactyly, 434
in congenital band syndrome, 461–462, 463f
Actinomycosis, 1870, 1870f
Active hypertrophy exercise(s), for poliomyelitis, during chronic phase, 1325–1326
Acute hematogenous osteomyelitis. *See Osteomyelitis, acute hematogenous*.
Acute infective polyneuritis, 1313t
Acute lymphocytic leukemia, back pain in, 104–105
Acute normovolemic hemodilution, 117
Adamantinoma, vs. Campanacci's disease, 1917
Adams' forward bending test
in Scheuermann's kyphosis, 324, 328f
in scoliosis, 217, 217f, 222
ADD (attention deficit disorder), and Legg-Calvé-Perthes disease, 663–664
Adduction, definition of, 27t
Adduction contracture(s), of hip, in cerebral palsy, surgical treatment of, 1173, 1178–1179, 1179f, 1180p–1183p, 1184, 1185f
Adductor hallucis brevis tendon, transfer to first metatarsal, for dorsal bunion, 1372–1373, 1379f, 1386f–1387f
Adductor longus tendon, tenotomy of, for closed reduction of hip, 542p–543p
Adductus longus muscle, dissection of, in open reduction of hip, medial approach to, 550–553
ADI. *See Atlanto-dens interval (ADI)*.
Adiposogenital syndrome, 715
Adolescence, growth spurt during, 1048, 1050f
AER (apical ectodermal ridge), and limb development, 379, 1745, 1746f

- AERMF (apical ectodermal ridge maintenance factor), and syndactyly, 433
- AFO(s). *See* Ankle-foot-orthoses (AFOs).
- AFP (alpha-fetoprotein), screening for, 1250
- Age
- and back pain, 96
 - and height, 10, 13f–18f
 - for boys, 10, 13f, 15f, 17f
 - for girls, 10, 14f, 16f, 18f
 - and interpretation of radiographs, 127
 - and joint range of motion, 28–29
 - of hip, 37–38, 38t
 - and mobility of cervical spine, 34t
 - and postlaminectomy kyphosis, 339
 - and signal intensity in magnetic resonance imaging, 154f, 154–155, 156f
 - and weight, 10, 14f–18f
 - for boys, 10, 13f, 15f, 17f
 - for girls, 10, 14f, 16f, 18f
- gestational, and incidence of cerebral palsy, 1123
- Airway, management of, in trauma, 2070
- Aitken classification, of femoral deficiency, 1756, 1758, 1758f–1759f
- Akin procedure, for hallux valgus, 1018
- Aldolase, in muscular dystrophy, 1466
- Alglucerase, for Gaucher's disease, 1645
- Aliasing, on magnetic resonance imaging, 155
- Alkaline phosphatase (ALP), serum, in osteosarcoma, 1959
- Allergic reaction(s), to latex, 117–118
 - in myelomeningocele, 1261
- Allis technique, of closed reduction in hip dislocation, 2275, 2278f, 2280f
- Allograft(s)
 - for scoliosis surgery, 240–241
 - vs.* prostheses, for limb salvage
 - after Ewing's sarcoma treatment, 2030f–2031f
 - after osteosarcoma treatment, 2028–2029, 2029f
- ALP (alkaline phosphatase), serum, in osteosarcoma, 1959
- Alpha-fetoprotein (AFP), screening for, 1250
- ALS (amyotrophic lateral sclerosis), anesthesia in, 120t
- Ambidexterity, in toddlers, 21
- AMBRI, in shoulder instability, 2131
- Ambulation
 - after bone graft epiphysiodesis, 736
 - in cerebral palsy, prognosis for, 1130
 - in Duchenne muscular dystrophy, lower limb surgery and, 1469–1470, 1470t, 1471f, 1472p–1475p
 - in hip spica cast, after femoral shaft fracture repair, 2314, 2315f
 - in myelomeningocele, 1253, 1255
 - level of lesion and, 1256–1257
 - in osteogenesis imperfecta, 1730
 - in proximal focal femoral deficiency, 1761, 1761f
 - with prosthesis, problems during, 1803
- Amelia, 1748, 1749t
 - upper, bilateral, 1757f
- American Society of Anesthesiologists, physical status classification of, 111, 111t
- American Spinal Injury Association (ASIA), evaluation form of, 2103, 2104f
- Amnion disruption sequence, and transverse upper limb deficiencies, 1784, 1785f–1786f, 1786–1787
- Amniotic band syndrome. *See* Congenital band syndrome.
- Amputation
 - above-knee, 1793–1794
 - ischial-bearing, for osteosarcoma, 1982p–1991p
 - prostheses for, 1806
 - below-knee, 1794, 1795f
 - for osteochondroma, 1996p–1999
 - prostheses for, 1806
 - bony overgrowth after, 1791–1792, 1791f–1792f, 1796
 - complications of, 1796, 1798, 1798f
 - congenital band syndrome and, 461, 463f
 - congenital tibial pseudarthrosis and, 881
 - for fibular hemimelia, 1777–1779
 - vs.* limb lengthening, 1772–1777
 - for macrodactyly, 458, 460f
 - forequarter, Littlewood technique of, 2000p–2013p
 - malignant tumors and, 1792
 - prosthesis fitting after, 1805
 - midthigh, for osteosarcoma, 1982p–1991p
 - of foot, with knee arthrodesis, for femoral deficiency, 1762f, 1762–1763, 1766p–1767p
 - of upper limb, 1793
 - psychosocial issues in, 1796
 - purpura fulminans and, 1792–1793, 1793f, 1796, 1796f
 - through arm, 2018p–2021p
 - traumatic, 1790–1792, 1790f–1793f, 1791t
 - debridement of, 1791
 - Ertl procedure in, 1792, 1793f
 - in-field care of, 1791
 - mechanism of, 1790, 1790f
 - prognosis in, 1791, 1791t
 - replantation after, 503–504
 - Amyloidosis, in rheumatoid arthritis, 1819
 - Amyoplasia. *See* Arthrogryposis.
 - Amyotonia congenita, anesthesia in, 120t
 - Amyotrophic lateral sclerosis (ALS), anesthesia in, 120t
 - Analgesia. *See also* Pain management.
 - patient-controlled. *See* Patient-controlled analgesia (PCA).
 - regional, 114–115
 - systemic, 112–114, 113t–114t
 - with sedation, 111, 111t
 - Analgia, congenital, 1448, 1449t
 - Anatomic axis
 - in femur, 1103, 1106f
 - vs.* mechanical axis, in lower extremities, 2322f
 - Anatomic snuff-box, swelling of, in scaphoid fractures, 2246, 2247f
 - Anderson-Green-Messner chart(s), for predicting remaining femoral/tibial growth, 1048–1051, 1049t, 1050f–1055f, 1052t
 - Anencephaly. *See* Myelomeningocele.
 - Anesthesia
 - and malignant hyperthermia, 118
 - asthma and, 110
 - for closed reduction of ankle fractures, 2395
 - in acrocephalosyndactyly syndromes, 120t
 - in muscular dystrophy, Emery-Dreifuss, 1479
 - in osteogenesis imperfecta, 124t, 1733
 - in surgical procedures for cerebral palsy, 1134
 - local, side effects of, 115–116
 - NPO requirements and, 110, 110t
 - orthopaedic syndromes affecting, 120t–125t
 - physical status classification for, 111, 111t
 - preoperative evaluation for, 109–111, 110t
 - Anesthesia (*Continued*)
 - regional, intravenous, 115–116
 - upper respiratory infection and, 110
 - Anesthetic(s), local, choice of, 115
 - Aneuploidy, gonosomal, anesthesia in, 122t
 - Aneurysm(s), of upper extremities, 497
 - Aneurysmal bone cyst(s). *See* Bone cyst(s), aneurysmal.
 - Angio-osteohypertrophy, anesthesia in, 122t
 - Angled thumb, 449
 - Angular deformity(ies), 25–26, 26f
 - of leg, 1103–1117, 1104f–1116f, 1105t. *See also* Leg, angular deformities of.
 - Anhidrosis, familial sensory neuropathy with, and insensitivity to pain, 1449t, 1453
 - Animal bite wound(s), 478
 - Anisomelia, and leg length inequality, 1039
 - Ankle
 - accessory ossification centers in, 2394–2395, 2399f
 - anatomy of, 2391–2392, 2392f
 - arthrodesis of
 - in poliomyelitis, 1373–1395, 1388f–1389f, 1390p–1393p, 1394f
 - distraction-compression, 1395, 1400p–1405p
 - extra-articular subtalar, 1388, 1396p–1399p
 - triple, 1383–1387, 1388–1389, 1390p–1393p
 - pantalar, with ankle fusion, 1388–1389, 1394–1395
 - disarticulation of, congenital tibial pseudarthrosis and, 881
 - dorsiflexion of, 43, 43f
 - and gait, 77, 78f
 - equinovarus deformity of
 - from unopposed activity of triceps surae and tibialis posterior muscles, in poliomyelitis, 1365, 1372f, 1374p–1377p
 - in cerebral palsy. *See* Cerebral palsy, foot and ankle in, equinovarus deformity of.
 - equinus deformity of, in Duchenne muscular dystrophy, 1467
 - eversion of, 44, 44f
 - fractures of, 2392–2414, 2393f–2417f
 - axial compression, 2394
 - classification of, 2392–2394, 2393f–2397f
 - complications of, 2403–2414, 2404f–2417f
 - delayed/nonunion as, 2405
 - premature physeal closure as, 2403–2405, 2404f, 2406f
 - valgus deformity as, 2405, 2407f
 - diagnosis of, 2394–2395, 2398f–2399f
 - pronation-eversion-lateral rotation, 2393–2394, 2396f–2397f
 - supination-inversion, 2393, 2393f–2394f
 - treatment of, 2400–2401, 2401f–2403f
 - supination-plantar flexion, 2393, 2394f–2395f
 - treatment of, 2397, 2400f
 - Tillaux, 2405–2410, 2407f–2410f
 - closed reduction of, 2405, 2407f
 - mechanism of, 2405, 2407f
 - open reduction of, 2407, 2409f–2410f
 - prognosis in, 2407, 2410
 - radiographic features of, 2405, 2408f
 - treatment of, 2395–2403, 2400f–2403f
 - anesthesia in, 2395
 - operative, indications for, 2396

- Ankle (*Continued*)

Salter-Harris type I and II distal fibular, 2396–2397

Salter-Harris type I tibial, 2397

Salter-Harris type II tibial, 2397–2398, 2400f–2401f

Salter-Harris type III distal tibial, 2399–2400, 2402f

Salter-Harris type IV distal tibial, 2401–2402, 2402f–2403f

Salter-Harris type V distal tibial, 2401–2403

triplane, 2410–2414, 2410f–2417f

closed reduction of, 2411–2412

mechanism of, 2410–2411, 2410f–2411f

open reduction of, 2412, 2414, 2415f–2417f

prognosis in, 2414

radiographic features of, 2411, 2412f–2414f

in cerebral palsy. *See Cerebral palsy, foot and ankle in.*

in myelomeningocele, 1262–1269. *See also Myelomeningocele, foot and ankle in.*

in poliomyelitis. *See Poliomyelitis, foot and ankle in.*

inversion of, 44, 44f

kinematics of

in cerebral palsy, 1128, 1128f

in sagittal plane, 79f

kinetics of, during gait, 80f, 81

motion of, limitation of, bone blocks for, 1395

plantar flexion of, 43, 43f

and gait, 77, 78f

push-off power of, in cerebral palsy, 1129, 1129f

range of motion of, 43–44, 43f–45f

stability of

assessment of, before triple arthrodesis, 1378

operative procedures for, history of, 1388t

syndesmosis of, 2391, 2392f

transfixation of, avoidance of, in treatment of congenital tibial pseudarthrosis, 877, 877f

valgus deformity of

after fractures, 2405, 2407f

after intramedullary rod fixation, for congenital tibial pseudarthrosis, 876, 876f

from congenital fibular pseudarthrosis, 884, 885f–886f, 887

hemiepiphiodesis in, 1113f, 1268

in cerebral palsy, 1153, 1158

in hereditary multiple exostosis, varus closing wedge osteotomy for, 1924, 1925f

in myelomeningocele, 1265, 1268f, 1269, 1270p–1271p

varus deformity of, after fractures, 2405, 2406f

Ankle clonus, 56

Ankle fusion, Chuinard, for triceps surae muscle paralysis, 1372

Ankle-foot orthoses (AFOs)

for cerebral palsy, 1131–1132, 1132f–1133f

for leg length inequality, 1058, 1061f

for myelomeningocele patient, 1292, 1293f

ground-reaction, for myelomeningocele, 1292, 1293f

Ankylosing spondylitis, 1821–1822, 1822f

back pain in, 103

Annular ligament, repair of, in reduction of Monteggia fractures, 2221, 2223f–2224f

Anoxia, perinatal, and cerebral palsy, 1123

Antalgic gait, 83

in Legg-Calvé-Perthes disease, 672f

“Anteater’s nose,” in calcaneonavicular coalition, 980, 980f

Anterior cruciate ligament (ACL)

anatomy of, 2354, 2354f, 2361f, 2361–2362

injuries to, 2361f–2365f, 2361–2366

diagnosis of, 2362–2364, 2363f–2364f

mechanism of, 2362

treatment of, 2364–2365, 2365f

Anterior drawer test, 2363, 2363f

Antibiotic(s)

for diskitis, 1862–1863

for Lyme disease, 1868

for osteomyelitis, 1858–1859

prophylactic, preoperative, 109

Anticholinesterase drug(s), for myasthenia gravis, 1501

Antiseizure medication(s), and rickets, 1690

AO external fixator, for open tibial fractures, 2383, 2384f–2385f

AO tension band, for olecranon fractures, 2200, 2203f

Aorta

ascending, dilation of, in Marfan’s syndrome, 1587, 1590

protection of, in anterior instrumentation, for idiopathic scoliosis, 268p–269p, 271p

Aortic arch, components of, 1754f

Apert’s syndrome, 466–467, 1615

anesthesia in, 120t

with tarsal coalition, 976, 977f

Apical ectodermal ridge (AER), and limb development, 379, 1745, 1746f

Apical ectodermal ridge maintenance factor (AERMF), and syndactyly, 433

Apical vertebra(e), wedging of, in Scheuermann’s kyphosis, 324, 327f

Aponeurotic fibroma, of upper extremities, 502

Apophyseal ring, fractures of, 99

Aquaplast, for splints, 388f, 389

Arachnodactyly

contractural, 1594f–1595f, 1594–1596

vs. Marfan’s syndrome, 1588, 1590

in Marfan’s syndrome, 1586–1587, 1587f

Arm. *See also Forearm; Upper extremity(ies).*

amputation through, 2018p–2021p

carrying angle of, 2165, 2167f

Arnold-Chiari malformation, 1253, 1254f

and torticollis, 193

Arteriovenous fistula(s), in Klippel-Trénaunay syndrome, 1670, 1672

Arthritis. *See also Osteoarthritis; under specific site, e.g., Knee.*

gonococcal, 1835–1836

pauciarticular juvenile, limping from, 88

psoriatic, 1822–1823, 1823f, 1823t

rheumatoid. *See Rheumatoid arthritis.*

septic, 1841–1859

clinical features of, 1845–1846

complications of, 1859, 1860f

definition of, 1841

diagnosis of, 85–87, 88f

differential diagnosis of, 1850–1852

epidemiology of, 1841

etiology of, 1841

from hip fractures, 2300

Arthritis (*Continued*)

gonococcal, 1864–1865

historical aspects of, 1841

imaging studies in, 1846–1848, 1847f–1852f

in infants, 1846

laboratory studies in, 1846

of hand, 478–479

of sacroiliac joint, 1863

pathogens in, 1844t, 1844–1845

pathophysiology of, 1843–1844

treatment of, 1852–1860, 1854p–1857p

antibiotics in, 1858–1859

aspiration in, 1852–1853, 1854p–1857p

surgical, 1853, 1854p–1857p, 1858

vs. acute transient synovitis of hip, 1824

syndromes in, 1823t

tuberculous, 1826–1831, 1828f–1830f, 1865–1868, 1866f–1867f

clinical features of, 1828–1829

incidence of, 1826

pathophysiology of, 1826, 1828, 1828f

radiographic features of, 1829f–1830f, 1829–1831

treatment of, 1831

Arthrochaliasis multiplex congenita, 1640

Arthrodesis

atlantoaxial

Brooks technique for, 187f

for Klippel-Feil syndrome, 185–187, 187f

Gallie technique for, 187f

Magerly technique for, 187f

distraction-compression, with ankle fusion, in poliomyelitis, 1395, 1400p–1405p

extra-articular subtalar, of foot and ankle, in poliomyelitis, 1388, 1396p–1399p

for hemophilic arthropathy, 1889

Grice extra-articular, for pes valgus, in cerebral palsy, 1152, 1154p–1157p, 1158f

metatarsophalangeal, for hallux valgus, in cerebral palsy, 1158, 1166f

occipitocervical

Dewar’s technique for, 185, 186f, 187

for Klippel-Feil syndrome, 185–187, 186f

of ankle, in poliomyelitis, 1373–1395, 1388f–1394f, 1390p–1393p

distraction-compression, 1395, 1400p–1405p

extra-articular subtalar, 1388, 1396p–1399p

triple, 1383–1387, 1388f–1389f, 1390p–1393p

of hip

for avascular necrosis, from slipped capital femoral epiphysis, 753, 756p–757p

for paralytic dislocation, 1350–1351, 1356f

for proximal focal femoral deficiency, 1770, 1773f–1774f

of knee, with foot amputation, for proximal focal femoral deficiency, 1762f, 1762–1763, 1764p–1767p

of subtalar joint, for pes planovalgus, 915, 915f

pantalar, with ankle fusion, in poliomyelitis, 1388–1389, 1394–1395

subtalar, for pes valgus, in cerebral palsy, 1152

triple

for Charcot-Marie-Tooth disease, 1440, 1443f

Volume 1, pgs. 1–710; Volume 2, pgs. 711–1502; Volume 3, pgs. 1503–2438.

- Arthrodesis (Continued)**
- for equinovarus deformity, in cerebral palsy, 1146–1147
 - for pes cavus, 1003, 1010–1011, 1010f–1011f
 - for pes planovalgus, 915, 916f
 - for pes valgus, in cerebral palsy, 1153, 1165f
 - for talipes equinovarus, with residual deformity, 951, 953f
 - for valgus overcorrection, 942–943, 944f of foot and ankle, in poliomyelitis, 1373–1395, 1388f–1389f, 1390p–1393p, 1394f
- Arthrography**
- during percutaneous *in situ* pinning of slipped capital femoral epiphysis, 722
 - in developmental dysplasia of hip, 532–533, 534f
 - after closed reduction, 544f
 - in Legg-Calvé-Perthes disease, 682, 685f
- Arthrogryposis**, 1647–1662, 1648f–1652f, 1656p–1659p, 1660f–1662f
- and teratologic hip dislocation, 646, 650, 650f
 - clinical features of, 1648–1649, 1649f–1651f
 - differential diagnosis of, 1650
 - etiology of, 1647
 - genetic basis of, 1647
 - hand in, 1655, 1660
 - pathophysiology of, 1649–1650
 - rotation flap for, 444, 447f
 - scoliosis in, 1660
 - treatment of, 1649f–1651f, 1651–1660, 1656p–1659p, 1660f–1661f
 - elbow in, 1654–1655, 1656p–1659p, 1660f–1662f
 - foot in, 1653–1654
 - goals of, 1651–1652
 - hip in, 1651–1652
 - knee in, 1652–1653
 - principles of, 1654
 - timing of, 1651
 - vs. Beals' syndrome, 1595
 - wrist in, 1655
- Arthrogryposis multiplex congenita**, anesthesia in, 120t
- Arthropathy**
- in hemophilia, 1880–1881, 1881f–1882f, 1882t. *See also* Hemophilia, arthropathy in.
 - neuropathic, 1825–1826, 1826f–1827f
- Articular cartilage**
- abnormalities of, in Legg-Calvé-Perthes disease, 669, 671f
 - limb lengthening and, 1087–1088
 - pathology of, in chondrolysis, from slipped capital femoral epiphysis, 749
 - thickening of, in Legg-Calvé-Perthes disease, 668, 671f
- Ascending aorta**, dilation of, in Marfan's syndrome, 1587, 1590
- ASIA (American Spinal Injury Association)**, evaluation form of, 2103, 2104f
- Asphyxiating thoracic dysplasia**, 1682f, 2153
- Aspirin**, for rheumatoid arthritis, 1820
- Asthma**, and anesthesia, 110
- Asymmetric tonic neck reflex**, 53, 54f
- Ataxia**, Friedreich's, 1246–1248, 1247f
- Athetosis**, in cerebral palsy, 1124
- in upper extremity, 1211, 1212f
- Athletic activity(ies)**
- and back pain, 95–96
 - and muscle strain injuries, of back, 98
 - restriction of, in spondylolysis, 100
- Atlanta Scottish Rite brace**, for Legg-Calvé-Perthes disease, 696, 698f
- Atlantoaxial arthrodesis**
- Brooks technique for, 187f
 - for Klippel-Feil syndrome, 185–187, 187f
 - Gallie technique for, 187f
 - Mageryl technique for, 187f
- Atlantoaxial instability**
- in Down syndrome, 1618
 - spinal fusion for, 1619–1621, 1620f
 - in Morquio syndrome, 1577, 1578f
 - in pseudoachondroplasia, 1524
 - traumatic, 2092, 2094f–2095f
- Atlantoaxial rotatory displacement**, 187–191, 188f–193f
- classification of, 189
 - clinical features of, 187, 188f, 189
 - pathophysiology of, 187, 188f
 - radiographic features of, 188f, 189
 - treatment of, 189–191, 190f–192f
- Atlanto-dens interval (ADI)**
- in atlantoaxial rotatory displacement, 188f
 - in cervical spine fractures, 2090, 2092, 2094f
 - in Down syndrome, 1617–1618
 - in Klippel-Feil syndrome, 177–178, 181–182
- Atlas**
- fractures of, 2092
 - ossification centers of, 2087, 2088f
- Attention deficit disorder (ADD)**, and Legg-Calvé-Perthes disease, 663–664
- Autologous blood donation**, 116–117
- Avascular necrosis (AVN)**
- after hip dislocation, 2280–2281, 2282f
 - vs. Legg-Calvé-Perthes disease, 690
 - after open hip reduction, 548
 - after Pavlik harness treatment, 536–537
 - and Köhler's disease, 899
 - corticosteroids and, 1698, 1699f
 - diagnosis of, computed tomography in, 146f
 - Frejka pillow and, 538, 538f
 - from hip fractures, 2288, 2289f, 2293–2296, 2294f–2297f
 - classification of, 2294, 2294f–2296f
 - incidence of, 2293
 - radiographic features of, 2294, 2294f–2296f
 - risk factors for, 2293–2294
 - treatment of, 2294–2296
- from humeral fractures, lateral condylar, 2185
- from intramedullary nailing, 2314, 2318f from slipped capital femoral epiphysis, 744–745, 750–753, 750f–755f, 756p–757p
- and extrusion of screws, 753, 754f
 - pathophysiology of, 750
 - prevention of, 751, 753
 - radiographic features of, 750f–752f, 751 treatment of, 751, 753, 754f–755f, 756p–757p
 - hip arthrodesis for, 753, 756f–757f
 - hip fusion for, 753, 755f
- from talar fractures, 2420
- in developmental dysplasia of hip, 569f, 582–584, 582f–586f, 587t
- after traction, 539–540
- classification of, 582–586, 582f–586f, 587t
- Dega osteotomy for, 631–632
- diagnosis of, 582
- Avascular necrosis (AVN) (Continued)**
- intertrochanteric double osteotomy for, 584, 597f, 612p–613p
 - pathophysiology of, 582, 582f
 - Salter innominate osteotomy for, 596, 618–621, 621f, 630f–633f, 631, 634p–641p. *See also* Salter innominate osteotomy, for avascular necrosis.
 - trochanteric advancement for, 584, 598p–611p
 - with lateral closing wedge valgus osteotomy, 584, 614p–617p
 - trochanteric epiphysiodesis for, 584, 588p–595p
 - valgus deformity from, 583, 583f
 - in Gaucher's disease, 1643–1645, 1644f
 - in Legg-Calvé-Perthes disease, 666, 666f
 - in sickle cell disease, 1894
 - of metatarsal head, and Freiberg's infarction, 901
 - AVN. *See* Avascular necrosis (AVN).
 - Avulsion fracture(s)**, in slipped vertebral apophysis, 374
 - Avulsion injury(ies)**, pelvic, treatment of, 2259, 2259f–2260f, 2260
 - Avulsive cortical irregularity**, 132, 133f
 - Axial fixator**, dynamic, for limb lengthening, 1092
 - Axillary freckling**, in neurofibromatosis, 1625, 1627f
 - Axillary nerve**, sensory distribution of, 2130f
 - Axis**, ossification centers of, 2087, 2088f
 - Axonotmesis**, in brachial plexus palsy, 480
- Babinski's reflex**, 56
- Back pain**
- differential diagnosis of
 - age and, 96
 - bone scan in, 98
 - choice of, 97, 97t
 - computed tomography in, 98
 - single-photon emission, 98
 - history in, 95–96
 - laboratory studies in, 98
 - magnetic resonance imaging in, 98
 - neurologic assessment in, 97
 - physical examination in, 96–97, 96f–97f
 - radiography in, 97–98
 - from athletic activities, 95–96
 - from developmental disorders, 99–102, 100f–102f
 - from disk herniation, lumbar, 371
 - in infectious/inflammatory disease, 102–104, 103f
 - in Marfan's syndrome, 304–305
 - in neoplastic disease, 104–105, 104f–105f
 - in Scheuermann's kyphosis, 325
 - in scoliosis, 221
 - in adults, 216
 - in slipped vertebral apophysis, 374
 - in spondylolisthesis, 354
 - incidence of, 95
 - intra-abdominal causes of, 105
 - intrathoracic causes of, 105
 - leg length inequality and, 1041
 - mechanical, 98–99, 99f
 - nature of, 95
 - psychosomatic, 96, 105–106
 - Backprojection, in computed tomography, 138, 139f

- Baclofen, intrathecal, for spasticity, in cerebral palsy, 1133
 Bado classification, of Monteggia fractures, 2219
 BAI (basion-axial interval), in occipito-atlantal instability, 1618, 1619f
 Bailey-Dubow rod, for fixation of long bone deformities, in osteogenesis imperfecta, 1733–1738, 1734f–1737f
 Baker procedure, for equinus deformity, in cerebral palsy, 1136, 1139f
 Ballet dancer(s), os trigonum injury in, 894, 895f
 Bankart lesion, 2129, 2129f
 Banks and Coleman hemipelvectomy, for osteosarcoma, 1964p–1971p
 Barlow's test, 63–64, 64f
 for developmental dysplasia of hip, 524, 524f
 Barr erector spinae transfer, for gluteus maximus muscle paralysis, 1336–1337, 1348, 1348f
 Base-of-neck osteotomy, for slipped capital femoral epiphysis, 737f, 737–738, 742p–743p
 Basic fibroblast growth factor (bFGF), and fibrodysplasia ossificans progressiva, 1636, 1638
 Basilar impression
 in Klippel-Feil syndrome, 176
 in osteogenesis imperfecta, 1739–1740
 Basion-axial interval (BAI), in occipito-atlantal instability, 1618, 1619f
 Battered child syndrome. *See Child abuse.*
 Baumann's angle, 2144, 2145f
 Bayne classification, of ulnar deficiency, 412f, 413, 1790, 1791f
 "Beak" triple arthrodesis, for pes cavus, 1010, 1011f
 Beals' syndrome, 1594f–1595f, 1594–1596
 vs. arthrogryposis, 1595
 vs. Larsen's syndrome, 1604
 vs. Marfan's syndrome, 1588, 1590
 Becker muscular dystrophy. *See Muscular dystrophy, Becker.*
 Beckwith-Wiedemann syndrome, 1039–1040
 anesthesia in, 120t
 Bedrest
 for Legg-Calvé-Perthes disease, 692
 for lumbar disk herniation, 373
 Bell Thompson ruler, for measurement of midthigh amputation level, 1982p–1991p
 Bell-Tawse technique, of annular ligament repair, in Monteggia fractures, 2224, 2224f
 Below-knee amputation, 1794, 1795f
 for osteochondroma, 1996p–1999p
 prostheses for, 1804
 Below-knee orthosis, plantar flexion-assist, 1326, 1326f
 Benign congenital hypotonia, 1313t
 Benik glove, 394, 395f
 Ben's splint, 393, 393f
 Benzodiazepine(s), for postoperative pain management, 113
 Berndt and Hardy classification, of osteochondral talar fractures, 2421, 2421f
 Bernese periacetabular osteotomy, for acetabular repositioning, 633, 642, 643f
 bFGF (basic fibroblast growth factor), and fibrodysplasia ossificans progressiva, 1636, 1638
 Biceps brachii muscle transfer, for supination contractures of forearm, 1425–1427
 Biceps brachii tendon, rerouting of, for brachial plexus palsy, with elbow deformities, 487, 498p–501p
 Biceps femoris muscle, lengthening of for iliotibial band contractures, in poliomyelitis, 1331, 1332p–1335p
 in hamstring lengthening, 1170p–1171p
 Biceps femoris tendon, transfer of, for quadriceps femoris paralysis, 1351, 1356–1358, 1357f
 Biceps reflex, 56
 Bigelow circumduction, in hip dislocation, 2275, 2278f
 Bikini incision
 for open reduction of hip, using anterolateral approach, 548, 560p–561p
 for Salter innominate osteotomy, 634p–635p
 Bilhaut-Cloquet procedure, for radial polydactyly, 426, 426f–427f
 Billings radiographic view, modified, in slipped capital femoral epiphysis, 719–720
 Biofeedback, for scoliosis, idiopathic, 232
 "Biologic capping," after amputation, 1792f, 1794
 "Biologic plasticity," 656
 Biophysical profile, low scores on, and risk of cerebral palsy, 1121
 Biopsy. *See also under specific location, e.g., Muscle, biopsy of; under specific tumor, e.g., Osteosarcoma, biopsy of.*
 incisional, of musculoskeletal tumors, 1899t, 1899–1900
 Bipartite navicular, 896, 898f
 Bipartite patella, 811
 Bipolar latissimus dorsi transfer, for elbow flexion, in arthrogryposis, 1655, 1660f–1662f
 Bipolar pectoralis major transfer, for elbow flexion, in arthrogryposis, 1654–1655, 1660f
 Birch classification, of fibular deficiency, 1776, 1777t
 Birmingham brace, for Legg-Calvé-Perthes disease, 696, 696f
 Birth history, in orthopaedic history, 23–24
 Bite wound(s), 478
 Bivalved cast(s), after tendon transfer, for poliomyelitis, 1329
 Bladder, paralysis of, from myelomeningocele, 1258–1259
 Blastomycosis, 1870, 1870f
 Bleeding
 during scoliosis surgery, 116
 hemophilia and, 1879–1880. *See also Hemophilia.*
 Block test
 for pes cavus, 986, 988f
 in Charcot-Marie-Tooth disease, 1436, 1439f
 Block vertebra, and congenital scoliosis, 284, 284f, 288f
 Blood donation, autologous, 116–117
 Blood transfusion(s). *See Transfusion(s).*
 Blood volume, calculation of, 2071
 Blount brace, 845–847, 847f
 Blount's disease. *See also Tibia vara, infantile.*
 adolescent, treatment of, epiphysiodesis in, 1113, 1114f
 with tibial varus deformity, 1104f, 1107, 1109f
 "Blowout" lesion, in aneurysmal bone cyst, 1909, 1912f
 Blue sclerae, in osteogenesis imperfecta, 1724–1725
 B-mode ultrasonography, 163
 BMP(s) (bone morphogenic proteins), in limb development, 1747
 BMP-4 (bone morphogenic protein-4), in fibrodysplasia ossificans progressiva, 1636
 Body position. *See Position.*
 Body righting reflex, 55
 Body size, proportionate, evolution of, 5, 5f, 10
 Body weight, in relation to age, 10, 13f–18f
 Bohler's angle, 2423
 Bone
 accessory, in foot, 895f
 atrophy of, in tuberculous arthritis, 1829, 1829f–1830f
 demineralization of, in hypophosphatasia, 1709f–1710f, 1710
 density of, in pyknodysostosis, 1565
 formation of
 as reaction to tumor, 1898–1899
 in Legg-Calvé-Perthes disease, 667, 667f–670f
 limb lengthening and, 1089f, 1090
 fragility of, in osteogenesis imperfecta, 1720, 1720t, 1722f, 1724
 infarction of, in sickle cell crisis, 1892–1893
 long. *See Long bone(s).*
 metastatic disease in, vs. osteopoikilosis, 1555
 new periosteal formation of, in neonate, 131, 131f
 overgrowth of, 2060–2062, 2062f
 after amputation, 1793–1794, 1793f–1794f, 1796
 plastic deformation of, 2059, 2060t
 regenerate, fracture of, after limb lengthening, 1091
 remodeling of, 2060–2062, 2062f
 after treatment of slipped capital femoral epiphysis, 758
 in fibrodysplasia ossificans progressiva, 1634
 subperiosteal formation of, in hypervitaminosis A, 1707, 1708f
 Bone age, delayed, in Legg-Calvé-Perthes disease, 661–662, 664f
 Bone age score, Oxford, in slipped capital femoral epiphysis, and prediction of contralateral slip, 747
 Bone block(s), for limitation of ankle motion, in pes calcaneus, 1395
 Bone crises, in Gaucher's disease, 1644
 Bone cyst(s), 1901–1906, 1902f–1906f
 aneurysmal, 104, 105f, 1906–1909, 1907f–1913f
 and cervical spine instability, 208f, 209
 clinical features of, 1908
 common sites of, 1906, 1907f
 etiology of, 1908
 pathology of, 1908, 1908f
 phases of, 1908–1909
 radiographic features of, 1908–1909, 1910f
 solid, 1908, 1908f
 treatment of, 1909, 1912f–1913f
 unicameral
 clinical features of, 1903
 common locations of, 1901
 differential diagnosis of, 1903
 etiology of, 1901, 1903
 pathology of, 1903, 1903f
 radiographic features of, 1902f, 1903
 treatment of, 1903–1906, 1905f–1906f
 autologous bone marrow injection in, 1905

- Bone cyst(s) (*Continued*)
 corticosteroids in, 1904–1905,
 1904f–1905f
 curettage and bone grafting in, 1906,
 1906f
 drilling in, 1905–1906
 Bone dysplasia, 1505–1583. *See also under*
specific disorder, e.g., Achondroplasia.
 classification of, 1506f, 1506–1510,
 1507t–1512t
 conditions associated with, 1506
 coxa vara in, 766–767
 diagnosis of, 1505
 differential diagnosis of, radiography in, 1513
 genetic transmission of, 1505
 Bone graft(s)
 after curettage of bone cyst
 aneurysmal, 1909, 1913f
 unicameral, 1906, 1906f
 bypass, for congenital tibial pseudarthrosis,
 869, 871f, 873
 for epiphysiodesis, for leg length inequality,
 1064p–1065p, 1070p–1071p
 for fibrous dysplasia, 1913
 for Grice extra-articular arthrodesis, 1152,
 1156p–1157p
 for open tibial fractures, 2383, 2386, 2387f
 for Salter innominate osteotomy, 638p–641p
 for scoliosis surgery, 240–241, 256p–257p
 with anterior instrumentation, 274p–275p
 for shelf procedure, 645f, 648f
 for subtalar extra-articular arthrodesis,
 1398p–1399p
 in Pemberton osteotomy, 628p–629p
 with compression wiring, for spondylolisthesis,
 359, 359f
 Bone graft arthrodesis, distraction-compression,
 of ankle, in poliomyelitis, 1395,
 1400p–1405p
 Bone graft epiphysiodesis, for slipped capital
 femoral epiphysis, 731, 732p–735p, 736
 Bone island(s), enlargement of, differential
 diagnosis of, 129
 Bone marrow
 autologous, injection of, for unicameral bone
 cyst, 1905
 Gaucher cells in, 1643, 1643f–1644f
 magnetic resonance imaging of, 159
 signal intensity in, 154f, 154–155, 156f
 Bone marrow space, vascular fibrous tissue of,
 in Caffey's disease, 1560, 1560f
 Bone marrow transplantation
 for Hurler's syndrome, 1573
 for osteogenesis imperfecta, 1729
 for osteopetrosis, 1551
 Bone morphogenic protein(s) (BMPs), in limb
 development, 1747
 Bone morphogenic protein-4 (BMP-4), in
 fibrodysplasia ossificans progressiva, 1636
 Bone plug, removal of, in greater trochanteric
 epiphysiodesis, for avascular necrosis,
 590p–593p
 Bone scan(s). *See also under specific disorder.*
 in differential diagnosis of back pain, 98
 indications for, 167
 radionuclide, in osteomyelitis, 1847
 Tc^{99m}
 in accessory navicular, 896, 898f
 in slipped capital femoral epiphysis, 721
 Bone screw(s), placement of, in anterior
 instrumentation, for idiopathic scoliosis,
 272p–273p
- Bone spur(s), from humeral fractures, lateral
 condylar, 2183, 2185f
 Bone transfer, free vascularized, microsurgery
 for, 508, 509f–511f, 512
 Bone tumor(s). *See also Musculoskeletal*
tumor(s); under specific type, e.g.,
Osteosarcoma.
 benign, of upper extremities, 503
 common anatomic sites of, 1896, 1896t
 limping from, 93
 malignant, 1955–2046
 Bone turnover, in osteogenesis imperfecta, 1722
 Boston brace, for scoliosis, idiopathic, 230, 231f
 adolescent, 230, 231f
 infantile, 277
 Botulinum toxin
 for cerebral palsy, in upper extremity, 1209
 for spasticity, in cerebral palsy, 1133–1134
 Boutonniere deformity, of proximal
 interphalangeal joint, treatment of, 476
 Bowlegs. *See Genu varum.*
 Boyd technique, for reduction of Monteggia
 fractures, 2221, 2222f
 Brace(s)
 Atlanta Scottish Rite, for Legg-Calvé-Perthes
 disease, 696, 698f
 Birmingham, for Legg-Calvé-Perthes disease,
 696, 696f
 for congenital scoliosis, 287
 for developmental dysplasia of hip, in young
 child, 537–538, 538f
 for idiopathic juvenile osteoporosis, 1717
 for idiopathic scoliosis, 229–232, 230f–233f
 compliance with, 232
 contraindications to, 229
 effectiveness of, 229
 hours of wear and, 231
 indications for, 229
 protocols for, 231–232, 233f
 types of, 229–231, 230f–231f, 233f
 for Marfan's syndrome, 305
 for osteogenesis imperfecta, 1730
 Milwaukee. *See Milwaukee brace.*
 Newington, for Legg-Calvé-Perthes disease,
 696, 697f
 pattern-bottom, for Legg-Calvé-Perthes
 disease, 658f
 prophylactic, after laminectomy, 310, 312
 Tachdjian, for Legg-Calvé-Perthes disease,
 696, 697f
 Toronto, for Legg-Calvé-Perthes disease,
 695f, 695–696
 Braces, Boston, for scoliosis, idiopathic, 230,
 231f
 adolescent, 230, 231f
 infantile, 277
 Brachial artery
 dissection of, in disarticulation of elbow,
 2022p–2023p
 injury to, from supracondylar humeral
 fractures, 2148f, 2162, 2163f, 2163–2165
 Brachial plexus, dissection of, in forequarter
 amputation, 2010p–2011p
 Brachial plexus palsy, obstetrical, 110p–117p,
 118f, 479–487, 482f–486f
 classification of, 480
 clinical features of, 480
 differential diagnosis of, 480–481
 etiology of, 479–480
 incidence of, 479
 initial treatment of, 481
 later treatment of, 481
 natural history of, 480
- Brachial plexus palsy (*Continued*)
 prognosis in, 480
 with residual deformities
 of elbow, 481
 management of, 486–487, 498p–501p
 of forearm, management of, 486–487,
 498p–501p
 of hand, 483–484
 management of, 487
 of shoulder, 481, 482f–483f, 483t
 management of, 484–486, 488p–495p,
 496f–497f
 of wrist, 481, 483, 484f
 Brachydactyly, in pseudohypoparathyroidism,
 1703f, 1704
 Brachytherapy, for soft tissue sarcoma, 2045,
 2045f
 Bracing
 for myelomeningocele, 1291. *See also*
Myelomeningocele, management of,
orthoses in.
 for scoliosis
 in cerebral palsy, 1225
 in Duchenne muscular dystrophy, 1471
 in myelomeningocele, 1281
 prophylactic, in congenital tibial
 pseudarthrosis, 868–869, 870f–872f
 Brain
 disorders of, 1121–1245. *See also under*
specific disorder, e.g., Cerebral palsy.
 levels affected by, 1121, 1122t
 pathology of, in myelomeningocele, 1253,
 1254f
 Breast(s)
 development of, in females, 19, 19f
 shielding of, during radiography, 137
 Breast cancer, radiation and, 224
 Breech presentation, and developmental
 dysplasia of hip, 515, 515f
 Brodie's abscess, 155f
 Brooks technique, for atlantoaxial arthrodesis,
 187f
 Brooks-Seddon pectoralis major muscle
 transfer, in poliomyelitis, 1418,
 1420p–1423p
 Broomstick plasters, for Legg-Calvé-Perthes
 disease, 656, 659f
 Brown procedure
 for hip stabilization, in proximal focal
 femoral deficiency, 1770, 1772f
 for tibial deficiency, 1780–1782
 Brucellar osteomyelitis, 1869
 Bryant's traction
 for developmental dysplasia of hip, 540f
 for femoral shaft fractures, 2307, 2307f
 Bucholz-Ogden classification system, for
 avascular necrosis, 582–583, 582f–586f
 Buck-Gramcko "stiletto flap," for fingertip
 reconstruction, in syndactyly, 441, 441f
 Buckle fracture(s), 2059–2060, 2060f–2061f
 of distal forearm, treatment of, 2237, 2238f
 of immature skeleton, 2059–2060,
 2060f–2061f
 of olecranon, 2202f
 of radius, 2060f–2061f
 Buddy strap(s), for finger splints, 392, 392f
 Bulbocavernosus reflex, 2103
 Bunion(s). *See Hallux valgus.*
 dorsal. *See Dorsal bunion.*
 Bunnell sternoclidomastoid muscle transfer, in
 poliomyelitis, 1418–1419, 1424f
 Bupivacaine, epidural administration of,
 114–115

- Burn injury, of hand, 400, 401f
- Burst fracture(s)
of cervical spine, 2096–2097
treatment of, 2105–2106
- Buschke-Ollendorff syndrome, 1555
- Butler procedure, for varus fifth toe, 1024, 1028f–1029f
- Butterfly vertebra(e), in myelomeningocele, 1289f
- Bypass bone graft, for congenital tibial pseudarthrosis, 869, 871f, 873
- C1–2 transarticular screw fixation, for atlantoaxial rotatory displacement, 191, 192f
- “Cable cast,” for talipes equinovarus, 937
- Cadence parameter(s), 75–76, 76t
- Café-au-lait spots
in fibrous dysplasia, 1911
in neurofibromatosis, 1625, 1626f
- Caffey’s disease, 1561–1565, 1562f–1563f, 1564t
clinical features of, 1562
diagnosis of, 1564
differential diagnosis of, 1564, 1564t
genetic features of, 1562
orthopaedic management of, 1565
pathology of, 1562, 1562f
prognosis in, 1565
radiographic features of, 1563f, 1564
treatment of, 1564
vs. hypervitaminosis A, 1566t
vs. progressive diaphyseal dysplasia, 1556
- Calcaneal apophysis, accessory, in Larsen’s syndrome, 1604, 1606f
- Calcaneal neck, osteotomy of, with bone graft, for pes planovalgus, 915, 917f
- Calcaneal osteotomy
for equinovarus deformity, in cerebral palsy, 1146
for pes cavovarus, in Charcot-Marie-Tooth disease, 1441f–1442f
for pes cavus, 995, 999, 1000p–1001p, 1002f–1003f
for pes valgus, in cerebral palsy, 1153
for talipes equinovarus, with residual deformity, 948, 950, 950f
for valgus overcorrection, 943, 944f
sliding, for pes planovalgus, 916, 919f–920f
- Calcaneocavus deformity, of foot, from triceps surae muscle paralysis, 1365, 1378f
- Calcaneocuboid capsulotomy, for talipes equinovarus, 940, 951f
- Calcaneocuboid joint, release of, for talipes equinovarus, 935–936, 937f
- Calcaneofibular release, for talipes equinovarus, 937, 938f
- Calcaneonavicular coalition, 978f–980f
limping from, 92, 92f
treatment of, 980–981
- Calcaneovalgus deformity, vs. vertical talus, 961
- Calcaneus
abnormalities of, in talipes equinovarus, 923–924, 924f–925f
anatomy of, 2423, 2423f
cyst in, 1902f
fractures of, 2423f–2424f, 2423–2425
classification of, 2423–2424, 2424f
complications of, 2425
diagnosis of, 2424–2425
mechanism of, 2423
treatment of, 2425
- Calcaneus (*Continued*)
lengthening of, for pes valgus, in cerebral palsy, 1152–1153, 1160p–1163p, 1164f
skeletal fixation of, for limb lengthening, 1079, 1079f
transfer of tibialis anterior tendon to, 1264–1265, 1266p–1267p
- Calcaneus deformity
from Achilles tendon lengthening, for cerebral palsy, 1138, 1142f
from congenital posteromedial tibial bowing, 883, 883f
in myelomeningocele, 1264–1265, 1265f, 1266p–1267p
in poliomyelitis, operative procedure for, 1365–1373, 1373f, 1379f, 1380p–1385p, 1386f–1387f
- Calcaneus gait, after talipes equinovarus surgery, muscle transfers for, 946, 947f
- Calcification
epiphyseal, in chondrodysplasia punctata, 1545f, 1546
of knee, in dermatomyositis/polymyositis, 1494, 1494f
of muscle, in myositis ossificans, 1497, 1497f
of soft tissue, in hypervitaminosis D, 1704, 1705f
- Calcium
deposition of, failure of, in rickets, 1686, 1686f
metabolism of, 1685, 1686f
- Calf
atrophy of, in pes cavus, 987, 989f
pseudohypertrophy of, in Becker muscular dystrophy, 1478, 1478f
- Callotasis, for limb lengthening, 1083–1086, 1084f–1085f
complications of, 1088, 1089f, 1090
- Callus(es)
hyperplastic, in osteogenesis imperfecta, 1726–1727, 1728f
plantar, in pes cavus, 985, 987f
- Calmodulin, in platelets, and scoliosis, 218
- Campanacci’s disease, 1914–1917, 1917f–1919f
clinical features of, 1915
differential diagnosis of, 1916–1917
etiology of, 1915
pathology of, 1915, 1917f
radiographic features of, 1916, 1918f–1919f
treatment of, 1917
- Camptomelic dwarfism, vs. osteogenesis imperfecta, 1728
- Camptomelic dysplasia, 1677–1679, 1678f
- Camurati-Engelmann disease, 1555f, 1555–1556
- Cannulated screw(s)
for fixation of slipped capital femoral epiphysis, unstable, 744
for slipped capital femoral epiphysis
in situ open pinning of, 729–731, 730f
percutaneous fixation of, 724p–727p
- Cantilever technique, with spinal fusion, for Scheuermann’s kyphosis, 334, 335f
- Capital femoral epiphysis, slipped. *See Slipped capital femoral epiphysis (SCFE).*
- Capitellum
fractures of, 2205
osteochondritis dissecans of, 2207–2208, 2208f
- Capsulectomy, for surgical drainage of hip, 1856p–1857p
- Capsuloplasty
for correction of vertical talus, 972p–973p
in open reduction of hip, using anterolateral approach, 566p–567p
- Capsulotomy, in open reduction of hip, using anterolateral approach, 562p–565p
- Carbon monoxide, exposure to, and congenital spinal anomalies, 280–281
- Cardiac surgery, neonatal, and risk of cerebral palsy, 1123
- Cardiovascular disease
in Marfan’s syndrome, 1588, 1589t, 1590
in muscular dystrophy
Becker, 1478
Duchenne, 1469
- Carpal bone
accessory ossification centers of, in Larsen’s syndrome, 1604, 1606f
duplication of, in mirror hand, 425
growth of, delayed, in Legg-Calvé-Perthes disease, 661–662, 664f
- Carpal coalition, in Ellis-van Creveld syndrome, 1680, 1681f
- Carpal tunnel syndrome
macrodactyly and, 454
with Hurler’s syndrome, 1575
- Carpectomy, proximal row, in arthrogryposis, 1655
- Carpenter’s syndrome, anesthesia in, 120t
- Carpotarsal osteolysis, 1572f
- Carpus, centralization of, for radial deficiency, 1787
- Carrying angle
of arm, 2165, 2167f
of elbow, 26, 26f
- Cartilage
abnormalities of, in diastrophic dysplasia, 1535
destruction of, in hemophilia, 1880
- Cartilage oligomeric matrix protein (COMP), and pseudoachondroplasia, 1523
- Cartilage-hair hypoplasia, McKusick type, 1549
- Cartilaginous cap, in solitary osteochondroma, 1917–1919
- Cassette(s), radiographic, 133
- Cast(s)
bivalved, after tendon transfer, for poliomyelitis, 1329
changing of, after treatment of talipes equinovarus, 941–942
- for humeral fractures, supracondylar, nondisplaced, 2149
- for idiopathic scoliosis, infantile, 277
- for Scheuermann’s kyphosis, 329, 330f
- Petrie, for Legg-Calvé-Perthes disease, 656, 659f, 695
- spica. *See Spica cast(s).*
- Casting
after closed hip reduction, 546, 548f
after correction of vertical talus, 966–967
after open hip reduction, 548–549
using anterolateral approach, 566p
using medial approach, 558p–559p
- for ankle fractures
Salter-Harris type I and II distal fibular, 2397
Salter-Harris type II distal tibial, 2397
- for brachial plexus palsy, with shoulder deformity, 485
- for closed hip reduction, 542p–543p
- for distal physeal fractures, 2328, 2329f
- for Köhler’s disease, 900
- for patellar dislocation, 2352
- for reduction of spondylolisthesis, 366, 367f
- for talipes calcaneovalgus, 907
- for talipes equinovarus, 927–928, 931–934
- for tibial diaphyseal fractures, 2379

Casting (*Continued*)
 for tibial spine fractures, 2355
 for vertical talus, 966
 of foot, and prevention of pressure sores, 1262, 1262f
 of radial and ulnar fractures, 2227, 2229f
 of supracondylar humeral fractures, 2156
 principles of, 2077, 2077f–2078f
 Cataract(s), in chondrodysplasia punctata, 1546
 Catheterization, clean intermittent, in myelomeningocele, 1258–1259
 Catterall classification, of Legg-Calvé-Perthes disease, 683–684, 686f
 Cauda equina, encroachment on, in low-grade spondylolisthesis, 361
 Caudal regression syndrome, 1305–1310, 1306f–1310f
 classification of, 1306, 1520f–1523f
 clinical features of, 1306–1310, 1310f
 pathology of, 1305–1306
 treatment of, 1310
 “Cauliflower ear,” in diastrophic dwarfism, 1535, 1536f
 Cavovarus deformity
 in lipomeningocele, 1302, 1303f
 in pes cavus, 985, 986f
 Cavus foot. *See* Pes cavus.
 Cellulitis, from puncture wounds, 1863
 Center post screw(s), for Texas Scottish Rite Hospital instrumentation, 246p–247p
 Center-edge angle, in developmental dysplasia of hip, 531, 532f
 Central core disease, anesthesia in, 120t
 Central nervous system (CNS), tumors of, and torticollis, 193, 193f
 Cerebellar level, neuromuscular disorders at, and motor function, 1121, 1122t
 Cerebral blood flow, induced hypotension and, 116
 Cerebral gigantism, anesthesia in, 125t
 Cerebral hypotonic diplegia, 1313t
 Cerebral palsy, 1122–1234
 ambulation in, prognosis for, 1130
 and fractures, 1230, 1232
 assessment of, 1124–1130, 1126f–1129f
 deep tendon reflexes in, 1126
 electromyography in, 1128–1129, 1129f
 gait analysis in, 1126f–1129f, 1126–1130
 history in, 1124
 motor milestones in, 1124–1125
 muscle tone in, 1126
 classification of, 1123–1124, 1124f–1125f
 definition of, 1122
 epidemiology of, 1122–1123
 etiology of, 1123
 fingers in, surgical procedures on, 1211, 1216p–1219p
 foot and ankle in, 1134–1158, 1135f–1147f, 1140p–1141p, 1148p–1149p, 1154p–1157p
 dorsal bunion of, 1158, 1166f
 equinovarus deformity of, 1142f–1147f, 1142–1150, 1148p–1149p
 calcaneal osteotomy for, 1146
 gait analysis in, 1142–1143, 1143f
 posterior tibialis tendon lengthening for, 1143, 1143f
 split anterior tibialis tendon transfer for, 1145–1146, 1148p–1149p
 split posterior tibialis tendon transfer for, 1143–1145, 1144f–1147f
 triple arthrodesis for, 1146–1147

Cerebral palsy (*Continued*)
 equinus deformity of, 1134–1142, 1135f–1139f, 1140p–1141p
 Achilles tendon lengthening for, 1136–1137, 1137f–1139f, 1140p–1141p
 and midfoot break, 1136, 1136f
 anterior Achilles tendon transposition for, 1137
 assessment of, 1135–1136, 1135f–1136f
 hallux valgus of, 1158, 1166f
 pes valgus of, 1150f–1151f, 1150–1152, 1154p–1157p, 1158f, 1164f–1166f
 calcaneal lengthening for, 1152–1153, 1160p–1163p, 1164f
 calcaneal osteotomy for, 1153
 clinical features of, 1150, 1150f
 Grice extra-articular arthrodesis for, 1152, 1154p–1157p, 1158f
 peroneal tendon lengthening for, 1152
 radiographic features of, 1150–1151, 1150f–1151f
 subtalar arthrodesis for, 1152
 triple arthrodesis for, 1153, 1165f
 valgus deformity of, 1153, 1158
 gait in, 84
 hip in, 1173–1203, 1178f–1179f, 1180p–1183p, 1185f–1186f, 1188f–1190f, 1192p–1195p, 1196f–1198f, 1200p–1201p, 1202f, 1204p–1207p, 1208f–1210f
 adduction contracture surgery of, 1173, 1178–1179, 1179f, 1180p–1183p, 1184, 1185f
 flexion contracture surgery of, 1173, 1178f–1179f
 subluxation/dislocation of
 anterior, 1203, 1211f
 clinical features of, 1185
 in nonreconstructable hip, 1197–1201, 1207f–1209f
 incidence of, 1182
 pathophysiology of, 1182, 1184f, 1185, 1186f
 radiographic features of, 1185, 1187f
 surgical treatment of, 1184–1199, 1186f–1190f, 1192p–1207p
 complications of, 1199, 1208f
 femoral osteotomy in, 1191, 1196, 1196f
 indications for, 1187, 1189
 pathophysiologic factors in, 1197, 1197f
 pelvic osteotomy in, 78–81, 79f–80f, 82p–83p, 84f, 86p–89p, 90f
 phases of, 1187, 1189
 soft tissue release in, 1180p–1183p, 1189–1193, 1192p–1195p
 histopathologic findings in, 1124
 knee in, 1158–1159, 1166f–1168f, 1166–1173, 1170p–1171p, 1172f–1178f
 hamstring spasticity from
 hamstring lengthening for, 1159, 1168f, 1170p–1171p
 measurement of, 1159, 1167f
 posture from, 1159, 1168f
 with fixed knee flexion contracture, 1159, 1167f
 rectus femoris transfer for, 1167–1169, 1174p–1175p
 rotational osteotomy for, 1169–1173, 1178f

Cerebral palsy (*Continued*)
 limping from, 88
 management of, 1130–1203, 1132f–1211f
 goals of, 1130
 medical treatment of spasticity in, 1132–1134
 orthoses in, 1131–1132, 1132f–1133f
 outcome assessment in, 1232
 physical therapy in, 1130–1131
 rhizotomy in, 1231–1232, 1232f
 surgical, 1132f–1211f, 1134–1203. *See also under specific area, e.g., Cerebral palsy, knee in.*
 anesthesia for, 1134
 general principles of, 1134
 timing of, 1134
 scoliosis in, 1222–1229, 1223f–1228f
 spinal fusion for, 1223–1229, 1226f–1228f
 anterior, 1225, 1228
 complications of, 1228f, 1228–1229
 Dunn-McCarthy technique of, 1227, 1229f
 Galveston technique of, 1227, 1230f
 Luque technique of, 1224f–1225f
 with spondylolisthesis, 1228
 with spondylosis, 1228
 spine in, 1224f–1230f, 1225–1230
 survival rate in, 1130
 thumb-in-palm deformity in, surgical treatment of, 1213, 1222, 1222f–1224f
 upper extremity in, 1211–1222, 1212f–1213f, 1214p–1221p, 1220f–1225f
 and growth, 1209
 and sensory impairment, 1209
 treatment of, 93–95, 1212f–1213f, 1214p–1221p, 1222f
 wrist in, surgical procedures on, 1211, 1212p–1219p
 Cervical artery(ies), and blood supply to femoral head, 660f
 Cervical dysraphism, and torticollis, 182f
 Cervical kyphosis, in Larsen’s syndrome, treatment of, 1604, 1607, 1607f
 Cervical line, posterior, 132, 132f
 Cervical spine
 anatomy of, 2087, 2088f
 disorders of, 171–210. *See also under specific disorder(s), e.g., Torticollis.*
 classification of, 171
 extension of, 34–35, 35f
 facet dislocation of, 2098
 flexion of, 34–35, 35f
 fractures of, 2087–2099, 2088f–2098f. *See also under specific segment, e.g., Odontoid, fractures of.*
 diagnosis of, 2087–2091, 2088f–2090f
 magnetic resonance imaging in, 2090–2091
 physical examination in, 2088
 radiography in, 2088–2090, 2088f–2090f
 initial management of, 2091, 2091f
 late complications of, 2098–2099
 mechanism of injury in, 2087
 pedicle, at C2, 2096, 2097f
 subaxial, 2096–2098, 2098f
 fusion of. *See Spinal fusion, cervical.*
 heterotopic ossification of, in fibrodysplasia ossificans progressiva, 1634, 1635f–1636f
 instability of, 206–210, 207f–209f
 in spondyloepiphyseal dysplasia congenita, 1528, 1528f–1529f
 ligamentous, 207
 occipital-atlantal, nontraumatic, 209

Cervical spine (*Continued*)
 occipital-atlantal anomalies and, 206, 209
 odontoid anomalies and, 209
 os odontoideum and, 206, 207f
 subaxial instability and, 207, 209f, 210
 kyphosis of, 195f–205f, 195–206. *See also*
 Kyphosis, cervical.
 lateral bending of, 35, 35f
 osteoblastoma of, 1936, 1937f
 posterior ligamentous instability of, 2090,
 2090f
 pseudosubluxation of, 131–132, 132f, 206f,
 2089–2090, 2090f
 range of motion of, 32–35, 34t, 35f
 rheumatoid arthritis of, 1819
 rotation of, 35
 spina bifida occulta of, in diastrophic
 dysplasia, 1536, 1537f
 spondylolysis of, in scoliosis, from cerebral
 palsy, 1231
 upper, instability of, in Down syndrome,
 1617f, 1617–1618
 treatment of, 1619–1621, 1620f
 Cervicothoracic incision, for forequarter
 amputation, 2000p–2001p
 Cervicothoracic spine, abnormalities of, in
 congenital scoliosis, 286f
 Chamberlain's line, in Klippel-Feil syndrome,
 176–177, 182f
 Charcot joint, 1827–1828, 1828f–1829f
 from congenital insensitivity to pain, 1448,
 1451f–1452f, 1827–1828, 1828f–1829f
 in lipomeningocele, 1303, 1304f
 Charcot-Marie-Tooth disease
 type 1, 1435–1445, 1436f–1447f
 claw toe in, 995f, 998f, 1438f–1439f,
 1440–1441
 clinical features of, 1435, 1436f
 diagnosis of, 1435–1436, 1437f
 genetic basis of, 1435
 hip dysplasia in, 1441, 1444f
 pes cavovarus in, 1436, 1438f–1442f, 1440
 pes cavus in, 995f, 998f, 1002f
 scoliosis in, 1441, 1445f–1446f
 treatment of, triple arthrodesis in, 1440,
 1443f
 type 2, 1445
 upper extremity weakness in, 1441, 1444,
 1447f
 Charleston brace, for scoliosis, idiopathic, 231,
 231f
 Check rein procedure, for talipes equinovarus,
 940, 941f
 Cheilectomy, for Legg-Calvé-Perthes disease,
 701
 Chemical shift artifact(s), on magnetic
 resonance imaging, 155
 Chemotherapy
 for congenital fibrosarcoma, 2042
 for osteosarcoma, 1962–1963
 for rhabdomyosarcoma, 2040–2041
 for soft tissue sarcoma, 2046
 Chevron osteotomy, for hallux valgus, 1016,
 1017f
 Chiari malformation(s), 1253, 1254f
 Chiari osteotomy
 for acetabular augmentation, 642, 644f, 646
 for avascular necrosis, 596
 for hip arthrodesis, for proximal focal
 femoral deficiency, 1770, 1773f–1774f
 for Legg-Calvé-Perthes disease, 701, 703f
 for subluxation/dislocation of hip, in cerebral
 palsy, 1197–1198, 1198f

"Chicken wire" calcification, in
 chondroblastoma, 1929, 1930f
 Chief complaint, in orthopaedic history, 23, 71
 Child(ren)
 cooperation milestones in, 381, 381t
 distraction of, 398
 Child abuse, 2079, 2081
 diagnosis of, bone scan in, 167, 167f
 femoral fractures from, 2302
 vs. osteogenesis imperfecta, 1728–1729
 Childbirth, clavicular fractures during, 2116,
 2119
 Chondroblastoma, 1929–1930, 1930f–1931f
 Chondrocyte(s), disordered, in slipped capital
 femoral epiphysis, 717f, 717–718
 Chondrodiatasis, for limb lengthening,
 1082–1083
 Chondrodysplasia, metaphyseal, 1547–1549,
 1548f–1549f
 Jansen type, 1547
 Schmid type, 1547–1548, 1548f–1549f
 Chondrodysplasia punctata, 1544f–1546f,
 1545–1546
 clinical features of, 196, 197f
 Chondroectodermal dysplasia, 1679f–1681f,
 1679–1682
 clinical features of, 1679–1680, 1679f–1680f
 genetic basis of, 1679
 radiographic features of, 1680–1681,
 1680f–1681f
 treatment of, 1681f, 1681–1682
 Chondrolysis
 in slipped capital femoral epiphysis,
 747–750, 749f
 clinical features of, 747–748, 749f
 epidemiology of, 748–749
 etiology of, 749
 from pin penetration, 730–731
 natural history of, 749–750
 pathology of, 749
 treatment of, 750
 limping from, 91, 91f
 Chondromatosis, synovial, 1944, 1946f
 Chondromyxoid fibroma, 1930–1933, 1932f
 Chondro-osseous tissue, in osteopetrosis, 1550,
 1551f
 Chondrosarcoma, 2038–2040
 progression of hereditary multiple exostosis
 to, 1924, 1926
 progression of multiple enchondromatosis
 to, 1928–1929
 progression of solitary osteochondroma to,
 1920, 1921t
 Chorionic villus sampling (CVS), and limb
 defects, 1755
 Chotzen's syndrome, anesthesia in, 120t
 Christmas disease, 1879
 Chronic renal failure, and slipped capital
 femoral epiphysis, 716
 Chuinard ankle fusion, for triceps surae muscle
 paralysis, 1372
 Chylothorax, in Gorham's disease, 1569f
 Cincinnati incision, for open reduction of
 vertical talus, 966, 968p–971p
 Ciprofloxacin, for puncture wounds, 1865
 Circulation, collateral, around elbow, 2163f
 Circumduction, during ambulation with
 prosthesis, 1802
 Circumflex artery(ies)
 anatomy of, 2284, 2284f–2285f
 medial, blood flow in, and Legg-Calvé-
 Perthes disease, 661f

CK (creatinine kinase), in muscular dystrophy,
 1465–1466
 Clamping mechanism, for Texas Scottish Rite
 Hospital instrumentation, 243, 243f,
 246p–247p
 Clasped thumb, 447
 rotation flap for, 444, 447f
 Clavicle
 anatomy of, 2f–3f, 2115
 "duplicated," 2115, 2117f
 hypoplasia/absence of, in cleidocranial
 dysostosis, 1568f–1569f, 1568–1570
 injuries to, 2115–2121, 2116f–2120f
 complications of, 2121
 diagnosis of, 2116, 2118, 2118f
 during childbirth, 2116, 2119
 mechanism of, 2116, 2118f
 midshaft, 2116, 2118, 2118f
 physeal, 2115, 2116f, 2118
 radiographic features of, 2118–2119, 2119f
 treatment of, 2119–2120, 2119f–2120f
 sectioning of, in forequarter amputation,
 2008p–2009p
 "Claw," placement of, in Texas Scottish Rite
 Hospital instrumentation, 254p–255p
 Claw toe
 in Charcot-Marie-Tooth disease, 995f, 998f,
 1438f–1439f, 1440–1441
 in pes cavus, 991, 995f, 998f
 Cleft foot, 983f, 983–984
 Cleft hand, with central polydactyly, 423, 423f
 Cleidocranial dysostosis, 1568f–1569f,
 1568–1570
 clinical features of, 1568f, 1569
 genetic features of, 1568–1569
 orthopaedic management of, 1569f, 1570
 radiographic features of, 1569f, 1569–1570
 Cleidocranial dysplasia, differential diagnosis of,
 1564t
 Clinodactyly, in Down syndrome, 1616f
 Closing wedge osteotomy
 Dwyer, for pes cavus, 995, 999,
 1000p–1001p, 1002f
 for angular deformities of leg, 1107, 1110f
 for cubitus varus, from supracondylar
 humeral fractures, 2168, 2169f–2171f
 for physiologic genu valgum, 860–861, 863f
 of midcarpus, in arthrogryposis, 1655
 plantar, for dorsal bunion, 944, 945f
 valgus, of proximal femur, with distal-lateral
 advancement of greater trochanter,
 614p–617p
 varus, for ankle valgus, in hereditary
 multiple exostosis, 1924, 1925f
 Clotting factor, for hemophilia, 1885–1887
 "Cloverleaf" skull, in thanatophoric dwarfism,
 1521
 Clubfoot. *See* Talipes equinovarus.
 Clubhand, radial
 free joint transfer in, 511, 511f
 surgical treatment of, 382
 CNS (central nervous system), tumors of, and
 torticollis, 193, 193f
 Coagulopathy, and Legg-Calvé-Perthes disease,
 656–657
 Cobb method, for measurement of curve in
 scoliosis, 225, 225f
 Cobb periosteal elevator(s), for Texas Scottish
 Rite Hospital instrumentation, 250p–251p
 Coccidioidomycosis, 1870, 1870f
 "Cock robin" appearance, in congenital
 muscular torticollis, 171, 173f
 Cock-up splint, for wrist, 394, 395f–396f, 396

- Codman's triangle, 1898
in osteosarcoma, 1959f
- Codman's tumor, 1929–3430, 1930f–1931f
- Cognitive function
in myelomeningocele, 1255
in neurofibromatosis, 1628–1629
- Cold abscess(es), in tuberculous arthritis, 1832–1833, 1833f
- Coleman block test
for pes cavus, 986, 988f
in Charcot-Marie-Tooth disease, 1436, 1439f
- Collaboration, multidisciplinary
in treatment of myelomeningocele, 1260
in treatment of rheumatoid arthritis, 467
- Collagen, metabolism of, 1718, 1719f
in osteogenesis imperfecta, 1718–1722
- Collagen vascular disease, anesthesia in, 120t
- Collateral circulation, around elbow, 2163f
- Collimator(s), 133
- Color Doppler ultrasonography, 163
- Communication
with parents, during physical examination, 72–73
with referring physicians, 73
- COMP (cartilage oligomeric matrix protein), and pseudoachondroplasia, 1523
- Compartment syndrome, 2075–2077, 2077f–2078f
diagnosis of, 2076, 2077f
from femoral shaft fractures, 2319
from Monteggia fractures, 2225
from open tibial fractures, 2386, 2386f
from proximal tibial phyeal fractures, 2351
from radial/ulnar fractures, 2233
from tibial tuberosity fractures, 2343
in elbow, 2141
of forearm, 403
of hand, 404
pathophysiology of, 2075–2076
prevention of, 2076
- Composite graft, for fingertip reconstruction, in syndactyly, 441, 441f
- Compression fracture(s)
in osteoporosis, juvenile idiopathic, 1716f
of cervical spine, 2097
treatment of, 2105
- Compression neuropathy, of ulnar nerve, in Volkmann's contracture, 408, 410f
- Compression pad(s), for securing total contact molding inserts, 390, 391f, 392
- Compression rod(s), for Scheuermann's kyphosis, 331, 335–336
- Compression wiring, with bone graft, for spondylolysis, 359, 359f
- Compressive force, in developmental coxa vara, 768, 769f
- Computed radiography (CR), 134–135
- Computed tomography (CT), 137–150, 138f–149f, 142t–143t. *See also under specific disorder, e.g., Scoliosis.*
backprojection in, 138, 139f
clinical applications of, 143f–149f, 145
electron beam, 139, 141f
history of, 137
in differential diagnosis of back pain, 98
in measurement of femoral anteversion, 784
patient preparation for, 142–143, 145
principles of, 137–140, 138f–142f, 142t
contrast resolution as, 140, 142t
slip ring technology in, 139, 140f–141f
spatial resolution as, 140, 142f
- Computed tomography (CT) (*Continued*)
radiation exposure from, 145
scanners for, 138–139, 138f–141f
technical considerations in, 140–143
- Computed tomography myelography, in lumbar disk herniation, 372f, 372–373
- Concentric muscle contraction(s), during gait, 79, 79t
- Confusion test, 1142, 1143f
- Congenital analgia, 1448, 1449t
- Congenital band syndrome, 461–466
and congenital tibial pseudarthrosis, 867–868
and limb absence, 1752, 1752f
classification of, 462–463, 463f
clinical features of, 461–462, 463f
etiology of, 461–466
foot deficiencies from, 1784, 1785f
pathology of, 461, 462f
treatment of, 463–466, 464f–467f
acute, 463–464
complications of, 466
reconstructive, 464f–465f, 464–466
results of, 466, 466f
- Congenital digitus minimus varus. *See Varus fifth toe.*
- Congenital snapping knee, 825, 826f
- Congruence angle, in patellar dislocation, measurement of, 796, 798f
- Connective tissue disorder(s), and scoliosis, 218
- Conradi-Hünermann syndrome, 1545
- Conradi's syndrome
clinical features of, 196, 197f
radiographic features of, 197f, 199
- Conscious sedation, 111
- Consolidation period, after corticotomy with callotasis, 1083, 1085f, 1088
- Contact dermatitis, from latex allergy, 117
- Containment treatment, for Legg-Calvé-Perthes disease, 655–656, 659f. *See also Legg-Calvé-Perthes disease, treatment of, containment.*
- Continuous passive motion (CPM), for talipes equinovarus, 929
- Contractural arachnodactyly, 1594f–1595f, 1594–1596
vs. Marfan's syndrome, 1588, 1590
- Contracture(s). *See also under specific body part, e.g., Elbow; under specific muscle, e.g., Sternocleidomastoid muscle.*
abduction, of hip, Ober's test for, 40, 41f
acquired, 26–27
assessment of, 27
congenital, 26
flexion, physiologic, in neonates, 3, 5f, 36, 38f
in Duchenne muscular dystrophy, 1468–1469
surgical correction of, 1469–1470, 1470t, 1471f, 1472p–1475p
in melorheostosis, 1560–1561
in poliomylitis, development of, 1325, 1325f
in vertical talus, 961, 964f
myostatic, 45
- Contract resolution, in computed tomography, 140, 142t
- Conversion reaction, back pain in, 105–106
- Cooperation, from pediatric patient, 71
- Cooperation milestone(s), in children, 381, 381t
- Cornelia de Lange's syndrome, 1663–1664, 1664f
- Coronoid, fractures of, 2205
- Coronoid line, 2145, 2147f
- Cortical hyperostosis, infantile. *See Caffey's disease.*
- Cortical irregularity, avulsive, 132, 133f
- Corticospinal level, neuromuscular disorders at, and motor function, 1121, 1122t
- Corticosteroid(s)
and avascular necrosis, 1698, 1699f
anesthesia and, 109
epidural, for lumbar disk herniation, 373
for bone cysts, unicameral, 1904–1905, 1904f–1905f
for dermatomyositis/polymyositis, 1495
for Duchenne muscular dystrophy, 1477
for progressive diaphyseal dysplasia, 1556
for spinal cord injury, 2106–2107
intra-articular, for rheumatoid arthritis, 469
of hand, 474, 474f–475f
- Corticotomy, with callotasis, for limb lengthening, 1083–1086, 1084f–1085f
incomplete, 1088, 1089f
- Cotrel-Dubousset instrumentation, for idiopathic scoliosis, 242–243, 243f
- Coventry and Johnston classification, of fibular deficiency, 1775–1776
- Coventry apparatus, internal fixation with, in oblique osteotomy of proximal femur, intertrochanteric, 1350, 1352p–1355p Lloyd-Roberts, 578p–571p
- Coxa valga, 25
- Coxa vara, 25
after hip fractures, 2296, 2298f
classification of, 765, 766t
congenital
treatment of, 770–778, 772f–774f, 776p–778p
goals of, 770
indications for, 770–771
with genu recurvatum, 826f
- developmental, 765–778
biomechanical features of, 768–770, 769f–771f
clinical features of, 765–766
hereditary factors in, 765
incidence of, 765
pathophysiology of, 768, 768f
treatment of
Pauwels' intertrochanteric Y-osteotomy in, 773, 774f, 775, 776p–779p
radiographic features of, 766f–767f, 768–768
recurrence of, 772–773
treatment of
greater trochanteric epiphysiodesis in, 772–773
valgus osteotomy in, 772f–773f, 772–775
- in bone dysplasia, 766–767
- in cleidocranial dysostosis, 1567f, 1568
- in osteopetrosis, 1551
- in proximal focal femoral deficiency, 1772f
- in spondyloepiphyseal dysplasia congenita, 1528–1529, 1528f–1529f
limping from, 89, 90f
- CPM (continuous passive motion), for talipes equinovarus, 929
- CR (computed radiography), 134–135
- Craig splint, for developmental dysplasia of hip, in child ages 1–6 months, 538, 538f
- Craniocarpotarsal dysplasia, 1662, 1663f
- Craniocervical stenosis, in achondroplasia, 1513–1514
- Cranioplast, in physeal bar resection, 2071, 2072f

- Craniosynostosis, anesthesia in, 120t, 123t
 Crankshaft phenomenon, 239f, 239–240, 1225
 C-reactive protein, in acute hematogenous osteomyelitis, 1846
 Creatine kinase (CK)
 in dermatomyositis/polymyositis, 1494
 in muscular dystrophy, 1465–1466
 facioscapulohumeral, 1481
 in muscular dystrophy, Emery-Dreifuss, 1479
 Cremasteric reflex, 56
 Cri-du-chat syndrome, anesthesia in, 120t
 Crossed extension reflex, 51, 53f
 Cross-leg sign, in Marfan's syndrome, 1587
 Crouch gait, in cerebral palsy, 1127, 1127f, 1158–1159, 1166f
 correction of, 1157, 1166f, 1170p–1171p
 Crouzon's disease, anesthesia in, 120t
 CT. *See* Computed tomography (CT).
 Cubitus valgus, 25, 26f
 from humeral fractures
 lateral condylar, 2185, 2188f
 supracondylar, 2165–2168
 Cubitus varus, 25
 from humeral fractures
 lateral condylar, 2183, 2185f
 supracondylar, 2165–2168, 2166f–2174f
 transphyseal, 2165, 2176f
 Cuboid
 abnormalities of, in talipes equinovarus, 923–924, 924f–925f
 position of, in vertical talus, 961, 966f
 Cup arthroplasty, for avascular necrosis, after hip fracture, 2296
 Curly toe(s), 1027, 1032f
 Cutaneous neurofibroma(s), 1625, 1627f
 CVS (chorionic villus sampling), and limb defects, 1755
 Cyst(s)
 bone. *See* Bone cyst(s).
 calcaneal, 1902f
 epidermoid inclusion, of upper extremities, 497
 metaphyseal, in Legg-Calvé-Perthes disease, 677–678, 678f
 popliteal, 828f–831f, 829–831, 832p–833p.
 See also Popliteal cyst(s).
 synovial, of upper extremities, 496–497
 Cystathione, deficiency of, and homocystinuria, 1596, 1597f
 Cytokine(s), and acute hematogenous osteomyelitis, 1843
- Dactylitis
 in psoriatic arthritis, 1822, 1823f
 sickle cell, 1864
 tuberculous, 1866, 1867f
 Dancer(s), os trigonum injury in, 894, 895f
 Dantrolene sodium, for malignant hyperthermia, 118
 Darrach resection, for synovitis, of wrist, 471–472, 472f
 DDH (developmental dysplasia of hip). *See* Hip, developmental dysplasia of.
 Deafness, in osteogenesis imperfecta, 1725
 Decubitus ulcer(s). *See* Pressure sore(s).
 Deep tendon reflex(es), 56
 in cerebral palsy, 1126
 Deformation(s), and growth and development, 5
 Deformity(ies), 25–27, 26f
 and contractures, 26–27. *See also* Contracture(s).
 and spasticity, 27. *See also* Spasticity.
- Deformity(ies) (*Continued*)
 angular, 25–26, 26f. *See also* Leg, angular deformities of.
 Dega osteotomy
 for avascular necrosis, 631–632
 for subluxation/dislocation of hip, in cerebral palsy, 1198–1199, 1202f, 1204p–1207p
 Degenerative joint disease (DJD), of hip, Legg-Calvé-Perthes disease and, 703–704
 Dejerine-Sottas disease, 1445, 1447
 Delbet classification, of hip fractures, 2285, 2286f–2287f
 Delee classification, of talar fractures, 2416, 2419f
 Delivery
 clavicular fractures during, 2116, 2119
 fetal position during, and brachial plexus palsy, 479–480
 Delta phalanx, 449, 1032–1033, 1036f
 in Rubinstein-Taybi syndrome, 1665, 1665f
 Deltoid, malignant schwannoma of, 2045f
 Deltoid muscle
 and shoulder motion, 1395
 dissection of, in shoulder disarticulation, 2014p–2017p
 paralysis of, in poliomyelitis, Saha trapezius transfer for, 1406f, 1406t, 1406–1407
 Demineralization, in hypophosphatasia, 1709f–1710f, 1710
 Denis classification, of thoracic and lumbar spine injuries, 2103, 2106t
 Dentinogenesis imperfecta, 1725, 1725f
 Denver Developmental Screening Test, 3, 4f
 Dermatitis, contact, from latex allergy, 117
 Dermatofibrosis lenticularis disseminata, 1557
 Dermatomyositis/polymyositis, 1492–1495, 1493f–1494f
 clinical features of, 1493–1494, 1493f–1494f
 diagnosis of, 1495
 etiology of, 1492–1493
 laboratory features of, 1494
 pathology of, 1493
 prognosis in, 1495
 treatment of, 1495
 Derotation maneuver, in Cotrel-Dubousset instrumentation, 243
 Derotation osteotomy, with femoral shortening, for open hip reduction, 570p–571p
 Derotation spondylodesis instrumentation, Zielke, for idiopathic scoliosis, 259–260, 260f
 Development. *See also* Growth and development.
 definition of, 3
 delayed, differential diagnosis of, 3, 5
 during puberty, Tanner's stages of, 19, 19f
 milestones in, 3, 4f, 19–21, 20t–21t
 for fine motor skills, 20t, 20–21
 for gross motor skills, 19–20, 20t
 for social skills, 21, 21t
 for verbal skills, 21, 21t
 Developmental disorder(s), back pain in, 99–102, 100f–102f
 Developmental dysplasia of hip (DDH). *See* Hip, developmental dysplasia of.
 Developmental reflex(es), 46–56, 49t, 50f–55f.
 See also Reflex(es).
 Dewar-Galeazzi procedure, for recurrent patellar dislocation, 797, 800p–805p
 Dewar's technique, for occipitocervical arthrodesis, 185, 186f, 187
 Dextroscoliosis, 147f
- Diaper(s), triple, for developmental dysplasia of hip, 538
 Diaphragm, dissection of, in anterior instrumentation, for thoracolumbar scoliosis, 266p–267p
 Diaphyseal dysplasia, progressive, 1553f, 1553–1554
 Diaphysis (diaphyses), cellular anatomy of, and acute hematogenous osteomyelitis, 1842, 1843f–1844f
 Dias-Tachdjian modification of Lauge-Hansen classification, of ankle fractures, 2392–2393, 2393f
 Diastematomyelia, 282, 282f, 1303–1305, 1304f–1305f
 clinical features of, 1303
 diagnosis of, magnetic resonance imaging in, 160f
 in congenital scoliosis, 288f
 in myelomeningocele, 1258
 radiographic features of, 1303, 1305, 1305f
 treatment of, 1305
 Diastrophic dysplasia, 1535f–1541f, 1535–1542
 clinical features of, 196, 196f, 1535–1537, 1535f–1537f
 differential diagnosis of, 1538–1539
 genetic features of, 1535
 “hitchhiker's thumb” in, 449, 1535, 1536f
 magnetic resonance imaging in, 160f
 orthopaedic management of, 1539f–1541f, 1539–1542
 pathology of, 1535
 prenatal diagnosis of, 1538
 radiographic features of, 199, 1537–1538, 1538f
 Digit(s). *See also* Finger(s); Toe(s).
 athetosis of, in cerebral palsy, 1212, 1212f
 reconstruction of in amniot disruption sequence, 1786f, 1786–1787
 Digital fibroma, infantile, of upper extremities, 502
 Digital gigantism. *See* Macrodactyly.
 Digital nerve, resection of, for macrodactyly, 457–458, 459f
 Digital radiography (DR), 135
 Digitus minimus varus, congenital. *See* Varus fifth toe.
 “Dimple effect,” in Legg-Calvé-Perthes disease, 679, 682
 Diplegia
 definition of, 44
 hypotonic, cerebral, 1313t
 in cerebral palsy, 1124, 1125f
 spastic, gait in, 81f
 Diplomyelia, 282f
 Discoid lateral meniscus, 814–815, 816p–821p, 822
 clinical features of, 814–815
 complete, 814
 diagnosis of, 815, 815f
 magnetic resonance imaging in, 159f
 treatment of, 815, 816p–821p, 822f
 partial resection in, 815, 822f
 total resection in, 815, 816p–821p
 anatomic considerations for, 818p–819p
 dissection for, 816p–817p, 820p–821p
 positioning for, 816f
 postoperative care and, 820
 incomplete, 814
 limping from, 91
 pathogenesis of, 814, 814f
 Wrisberg ligament type, 814, 814f

Diskectomy
anterior, with spinal fusion, for infantile idiopathic scoliosis, 277–278, 278f–279f
for lumbar disk herniation, 373–374
thoracoscopic, with spinal fusion, for Scheuermann's kyphosis, 334
with anterior instrumentation, for idiopathic scoliosis, 270p–271p

Diskitis, 1862–1863
and torticollis, 193–194
back pain from, 102–103, 103f
diagnosis of
imaging studies in, 102, 103f
magnetic resonance imaging in, 1851f
limping from, 87–88, 89f
treatment of, 103
vs. vertebral osteomyelitis, 87–88

Disruption(s), in growth and development, 5

Distal metatarsal articular angle (DMAA), in hallux valgus, 1014

Distraction, of children, 398

Distraction lengthening, of hand, 1783, 1785f

Distraction osteogenesis, with external fixation, for congenital tibial pseudarthrosis, 878, 879f–880f, 881

Distraction period, after corticotomy with callotasis, 1083, 1085f

Distraction-compression bone graft arthrodesis, of ankle, in poliomyelitis, 1395, 1400p–1405p

DJD (degenerative joint disease), of hip, Legg-Calvé-Perthes disease and, 703–704

DMAA (distal metatarsal articular angle), in hallux valgus, 1014

Dome osteotomy
for genu recurvatum, in poliomyelitis, 1359, 1360f
midtarsal, for pes cavus, 999, 1003
with release of Vickers' ligament, for Madelung's deformity, 419–420, 420f

Doppler ultrasonography, 163

Dorsal bunion
in cerebral palsy, 1158, 1166f
in triceps surae muscle paralysis, 1372–1373, 1378f–1379f, 1386f–1387f
plantar, closing wedge osteotomy for, 944, 1171f

Dorsal flap, for syndactyly, 437, 437f–439f
thumb-index web, 444, 446f

Dorsal wedge osteotomy, for pes cavus, 999, 1004p–1005p

Dorsiflexion, muscles producing, 1361

"Dotted veil appearance," in myositis ossificans, 1497

Double inclinometer test, of thoracolumbar spine flexion, 36, 37f

Double-button wire(s), for Wisconsin segmental spinal instrumentation, 242, 242f, 244p–245p

Down syndrome, 1616–1623
and qualification for Special Olympics, 1618
anesthesia in, 121t
clinical features of, 1616f, 1616–1617
foot disorders in, 1623
genetic basis of, 1616
hip dysplasia in, treatment of, 1621f–1622f, 1621–1623
instability of upper cervical spine in, 1617f, 1617–1618
treatment of, 1619–1621, 1620f
patellofemoral instability in, 1623
prenatal screening for, 1616

DR (digital radiography), 135

Dressing(s), for hand, 382–387, 385f–387f. *See also* Hand, dressings for.

Drug(s)
and rickets, 1690
prenatal exposure to, and cerebral palsy, 1123

Duchenne muscular dystrophy. *See* Muscular dystrophy, Duchenne.

Duncan Ely test, 1166, 1172f

Dunlap radiographic view, in slipped capital femoral epiphysis, 719–720

Dunn procedure, 736–737, 737f, 740p–741p

Dunn-McCarthy instrumentation, for spinal fusion, in spinal muscular atrophy, 1317f

Dunn-McCarthy technique
for spinal fusion, in cerebral palsy, 1227, 1229f
of lumbar kyphectomy, for myelomeningocele, 1280, 1282p–1287p

Dwarfism. *See also* Achondroplasia.
camptomelic, *vs.* osteogenesis imperfecta, 1728
diastrophic. *See* Diastrophic dysplasia.
metatropic, 1673, 1674f–1677f, 1677
pituitary, 1711–1712
thanatophoric, 1521–1523, 1522f

Dwyer closing wedge osteotomy, for pes cavus, 995, 999, 1000p–1001p, 1002f

Dwyer instrumentation, for idiopathic scoliosis, 259, 259f

Dwyer technique, for talipes equinovarus, with residual deformity, 948, 950, 950f

Dynamic axial fixator, for limb lengthening, 1092

Dysautonomia
anesthesia in, 124t
familial, 1449t, 1453

Dysostosis, classification of, 1508t–1509t

Dysplasia(s), and growth and development, 5

Dysplasia epiphysealis hemimelica, 1946, 1947f–1948f

Dysplastic spondylolisthesis, 351–352, 352f

Dystonia, in cerebral palsy, 1123
in upper extremity, 1212

Dystrophin, amount of, in muscular dystrophy, 1463, 1465
Becker, 1477

Ear, "crumpled" appearance of, in Beals' syndrome, 1594, 1594f

Eaton's angle, 408, 409f

Eccentric muscle contraction(s), during gait, 79t, 79–80

Echo time (TE), in magnetic resonance imaging, 152, 152t

Ectopia lentis, in Marfan's syndrome, 1587

Eddy current distortion, on magnetic resonance imaging, 155

Eggers procedure, for hamstring transfer to femoral condyles, 1159, 1168f

Ehlers-Danlos syndrome, 1639–1640, 1639f–1641f, 1642t
anesthesia in, 121t
classification of, 1640, 1642t
clinical features of, 1639, 1639f–1641f

Elbow. *See also* Humerus; Radius; Ulna.
carrying angle of, 26, 26f
collateral circulation around, 2163f
deformities of, from brachial plexus palsy, 481
management of, 486–487, 498p–501p

Elbow (*Continued*)
disarticulation of, 2022p–2025p
dislocation of, 2143f, 2191–2194, 2191f–2194f
anatomic considerations in, 2191
and radial neck fractures, 2194, 2195f
anterior, 2192
complications of, 2193–2194, 2193f–2194f
diagnosis of, 2192
in Larsen's syndrome, 1602, 1604f, 1611
mechanism of, 2191, 2191f
recurrent, 2193–2194, 2194f
treatment of, 2192f, 2192–2193
extension contractures of, posterior release of, 1654, 1656p–1659p
flexion contractures of, in cerebral palsy, 1212
flexion of, procedures for, in arthrogryposis, 1654, 1660f–1662f
fractures about, 2139–2208, 2140f–2208f
hyperextension of, 31, 33f
in poliomyelitis, 1408–1419, 1409f, 1410p–1426p. *See also* Poliomyelitis, elbow in
ischemic, realignment of, 2147, 2149, 2150f
"Little League," 2208, 2209f
neurovascular anatomy of, 2139, 2142f
nursemaid's, 2206f, 2206–2207
osteochondrosis of, 2207–2208, 2208f
range of motion of, 31, 33f
secondary ossification centers of, 2139, 2139f
splints for, 396–398, 398f
ulnar deficiency in, treatment of, 414, 414f

Elbow release, with tricepsplasty, for arthrogryposis, 1654, 1656p–1659p

Electrical stimulation
for congenital tibial pseudarthrosis, 878
for scoliosis, idiopathic, 232

Electromyography (EMG)
for predicting outcome of rectus femoris transfer, 1166–1167
in cerebral palsy, 1128–1129, 1129f
in dermatomyositis/polymyositis, 1494
in muscular dystrophy, 1467

Electron beam computed tomography, 139, 141f

Ellis-van Creveld syndrome, 1679–1680, 1679f–1680f
clinical features of, 1679–1680, 1679f–1680f
genetic basis of, 1679
radiographic features of, 1680–1681, 1680f–1681f
treatment of, 1681f, 1681–1682

Ely test, 1166, 1172f

Embolism
fat, from femoral shortening, 1076
pulmonary, after spinal cord injury, 2107–2108

Embryonal rhabdomyosarcoma, 2040

Emery-Dreifuss muscular dystrophy, clinical features of, 1464t

EMG. *See* Electromyography (EMG).

Encephalocele, 1249

Enchondral ossification, failure of, and achondroplasia, 1510

Enchondroma
of femur, angular deformity from, 1104f, 1106, 1107f–1108f
solitary, 1926f, 1926–1927

Enchondromatosis
multiple, 1927–1929, 1928f–1929f
of upper extremities, 503
Enders rod(s), for femoral shaft fractures, 2305, 2306f, 2313f, 2316f
Endobone(s), in osteopetrosis, 1551, 1552f
Endocrine disorder(s)
and slipped capital femoral epiphysis, 715–716
of bone, 1685–1740. *See also under specific disorder, e.g., Hypoparathyroidism.*
Energy expenditure, during ambulation with prosthesis, 1803
Enneking classification, of musculoskeletal tumors, 1897t, 1898
Enzyme replacement therapy, for Gaucher's disease, 1645
Eosinophilic granuloma, 104, 1938f–1941f, 1938–1942
and torticollis, 194
clinical features of, 1938–1939
differential diagnosis of, 1940–1941
multifocal, with extraskeletal involvement, 1942
pathology of, 1938, 1938f
radiographic features of, 1939–1940, 1939f–1941f
treatment of, 1941–1942
Epidermoid inclusion cyst(s), of upper extremities, 497
Epidermolysis bullosa, anesthesia in, 121t
Epidural analgesia, 114–115
after scoliosis surgery, 241
Epiphyseal arrest, for macrodactyly, 457
Epiphyseal cartilage, abnormalities of, in Legg-Calvé-Perthes disease, 669, 671f
Epiphyseal dysplasia, multiple, vs. Legg-Calvé-Perthes disease, 689–690, 694f
Epiphyseal nerve, injury to, from radial/ulnar fractures, 2233
Epiphyseal stapling, for leg length inequality, 1063, 1073–1074, 1073f–1074f
Epiphysiodesis
bone graft
for leg length inequality, 1064p–1065p, 1070p–1071p
for slipped capital femoral epiphysis, 731, 732p–735p, 736
for angular deformities of leg, 1113, 1114f
for idiopathic scoliosis, infantile, 278–279
for leg length inequality, 1058–1063, 1062f, 1064p–1071p, 1072f
for valgus deformity, in cerebral palsy, 1153, 1158
lateral, for adolescent tibia vara, 851–853, 854f
tibial, for valgus deformity, 2376, 2376f
timing of
Moseley straight line graphs in, 1051–1053, 1056p–1057p, 1059f
Rotterdam straight line graphs in, 1060f
trochanteric. *See Trochanteric epiphysiodesis.*
Epiphysiolysis
for limb lengthening, 1082–1083
for Madelung's deformity, 419
proximal humeral, 2128, 2128f
Epiphysis (epiphyses)
calcification of, in chondrodysplasia punctata, 1545f, 1546
chondroblastoma in, 1930, 1931f
growth of, disturbance of, in avascular necrosis, 583, 583f–584f

Epiphysis (epiphyses) (*Continued*)
impaired ossification of, in
pseudoachondroplasia, 1525, 1526f
of lower extremities
closure of growth plates in, age at, 9f
secondary ossification centers in, age at
appearance of, 7f
of upper extremities
closure of growth plates in, age at, 7f
secondary ossification centers in, age at
appearance of, 6f
ossification of, disturbance of, in multiple epiphyseal dysplasia, 1531–1533, 1533f
pressure, 1045
“ringing” of, in scurvy, 1706, 1706f
sclerosis of, in melorheostosis, 1560, 1560f
traction, 1045
Equiangular spiral, movement of hand in, 405, 407f
Equinovarus deformity, in lipomeningocele, 1302, 1303f
Equinovalgus deformity, of foot
from anterior tibial muscle paralysis, 1364, 1365f
in Larsen's syndrome, 1610–1611, 1612f
Equinovarus deformity. *See under specific site, e.g., Ankle.*
Equinus deformity. *See under Talipes.*
Equinus prosthesis, 1802
Erlenmeyer flask deformity, in Gaucher's disease, 1643, 1644f
Ertl procedure, 1794, 1795f
Erythema nigrans, in Lyme disease, 1868
Erythrocyte sedimentation rate (ESR), in acute hematogenous osteomyelitis, 1846
Erythropoietin, with autologous blood donation, 116
ESR (erythrocyte sedimentation rate), in acute hematogenous osteomyelitis, 1846
Ethnicity, and myelomeningocele, 1249–1250
Etidronate, for fibrodysplasia ossificans progressiva, 1638
Evans procedure, for recurrent talipes equinovarus, 948, 949f
Eversion
definition of, 27t
muscles producing, 1361
Ewing's sarcoma, 2030f–2037f, 2030–2038, 2032t
back pain from, 105
biopsy of, 2034–2035
clinical features of, 2033
common sites of, 2032, 2032t
femoral, 1848f
metastatic, 2038
pathology of, 2032f, 2032–2033
prognosis in, 2035
radiographic features of, 2030f–2031f, 2033–2034, 2034f
staging of, 2034
treatment of, 2035, 2036f–2037f, 2038
Examination, 25–56, 70–73, 71t, 72f. *See also under specific part and condition.*
communication with parents during, 72–73
form for, 58
guidelines for, 71t
history in, 71
neurologic, 46–56, 49t, 50f–55f. *See also Neurologic assessment.*
of muscle strength, 44–46, 45t–48t

Examination (*Continued*)
of range of motion, 27t, 27–44, 28f–45f, 34t, 38t. *See also Range of motion.*
orthopaedic, 63–73
focused, 65–70, 65f–70f
screening, 63–65, 64f
recognition of deformities on, 25–27, 26f
Excessive lateral pressure syndrome, patellar, 810
Exercise
for poliomyelitis
during chronic phase, 1325–1326
during convalescent phase, 1324
for tenosynovitis, of wrist, 470
stretching
for congenital muscular torticollis, 172, 175f
for pes planovalgus, with Achilles tendon contracture, 912–913, 913f
Exostosis
hereditary multiple. *See Hereditary multiple exostosis.*
subungual, 1032, 1033f
EXT gene(s), and hereditary multiple exostosis, 1921
Extension. *See also under specific joint.*
definition of, 27t
Extension blocking splint, for wrist, 394, 396f
Extension contracture(s), of knee, in myelomeningocele, 1269, 1272
Extensor carpi radialis brevis tendon, transfer of extensor carpi ulnaris tendon to, 1213, 1214p–1217p
Extensor carpi ulnaris tendon, transfer to extensor carpi radialis brevis tendon, 1213, 1214p–1217p
Extensor digitorum longus muscle, paralysis of, in poliomyelitis, 1363–1364
Extensor hallucis longus muscle, paralysis of, in poliomyelitis, 1363–1364
Extensor thrust reflex, 52
External fixator(s)
for distal metaphyseal femoral fractures, 2325–2326
for femoral shaft fractures, 2308, 2310, 2310f–2311f
for pelvic fractures, 2264–2266, 2266f
for tibial fractures
diaphyseal, 2379, 2381f
open, 2383, 2384f–2385f
External oblique abdominal muscle, for iliopsoas transfer, 1336
External oblique muscle transfer, for subluxation/dislocation of hip, in myelomeningocele, 1274, 1276f
Extra-articular subtalar arthrodesis, of foot and ankle, for poliomyelitis, 1388, 1396p–1399p
Extracapsular ring, anatomy of, and Legg-Calvé-Perthes disease, 659–660
Extrapyramidal level, neuromuscular disorders at, and motor function, 1121, 1122t
Extremity(ies)
hemihypertrophy of
in neurofibromatosis, 1628, 1631f
macrodactyly with, 451–452, 453f, 458–459
lower. *See Lower extremity(ies).*
severed, replantation of, 503–504
upper. *See Upper extremity(ies).*
Eyebolt(s), placement of, in anterior instrumentation, 274p–275p

- Eyebolt-nut system, for Texas Scottish Rite Hospital instrumentation, 243, 243f
- Eyelid(s), heliotropic, in dermatomyositis/polymyositis, 1493f, 1494
- Ezaki classification, of polydactyly, 422t
- Ezeform, for splints, 388, 388f
- Facial diplegia, congenital, anesthesia in, 123t
- Facies
- in achondroplasia, 1511, 1511f
 - in Apert's syndrome, 467, 1615
 - in bone dysplasia, 1505
 - in cleidocranial dysostosis, 1568f, 1569
 - in Down syndrome, 1616, 1616f
 - in fibrous dysplasia, 1911, 1914f
 - in Hurler's syndrome, 1574, 1574f
 - in Kniest's dysplasia, 1542
 - in Larsen's syndrome, 1602
 - in myotonic dystrophy, 1487, 1488f
 - in osteogenesis imperfecta, 1724
 - in pseudoachondroplasia, 1524f
- Faciocapulohumeral muscular dystrophy. *See* Muscular dystrophy, faciocapulohumeral.
- Factor V Leiden mutation(s), and Legg-Calvé-Perthes disease, 657
- Factor VIII
- administration of, for hemophilia, 1885–1887
 - deficiency of, and hemophilia, 1879–1880
- Fairbanks apprehension test, 794
- in recurrent patellar subluxation, 799
- Familial dysautonomia, 1449t, 1453
- anesthesia in, 124t
- Familial hypophosphatemic rickets. *See* Rickets, vitamin D-resistant.
- Familial periodic paralysis, anesthesia in, 121t
- Family history, in orthopaedic history, 23
- Fanconi's pancytopenia syndrome, 1752
- Fascia lata, dissection of, in trochanteric transfer, for avascular necrosis, 598p–601p
- Fascia graft, in Steindler flexorplasty, 1408–1409, 1409f
- Fasciotomy, for compartment syndrome, 2076, 2077f–2078f, 2078
- of forearm, 403, 403f
- Fat, infiltration of nerves by, in macrodactyly, 451f, 452
- Fat embolism, from femoral shortening, 1076
- "Fat pad sign," in supracondylar humeral fractures, 2144, 2147f
- Fatigue, in poliomyelitis, during convalescent phase, 1324
- FBNI gene, mutations in, and Marfan's syndrome, 1587–1588
- Feeding skill(s), development of, 20
- Felon, 477, 477f
- Femoral artery, dissection of, in midthigh amputation, 1984p–1985p
- Femoral condyle
- medial, osteochondritis dissecans of, 791f
 - transfer of hamstrings to, Eggers procedure for, 1159, 1168f
- Femoral cutaneous nerve, lateral, protection of, in open reduction of hip, using anterolateral approach, 562p–563p
- Femoral deficiency, proximal focal, 1756–1773, 1758f–1762f, 1764p–1769p, 1770f–1778f
- classification of, 1756–1760, 1758f–1760f
 - Aitken, 1756, 1758, 1758f–1759f
 - Fixsen and Lloyd-Roberts, 1760
 - Gillespie, 1760, 1760f
 - Gillespie and Torode, 1759–1760
- Femoral deficiency, proximal focal (*Continued*)
- Hamanishi, 1758–1759, 1760f
 - clinical features of, 1760–1761, 1761f
 - diagnosis of, ultrasonography in, 164, 165f
 - prostheses for, 1611f, 1763, 1804
 - treatment of, 1762f, 1762–1763, 1764p–1767p, 1770–1773, 1770f–1778f
 - amputation in, 1762f, 1762–1763, 1764p–1767p
 - femoral pseudarthrosis stabilization in, 1763, 1772f
 - hip stabilization in, 1763, 1770, 1773f–1775f
 - limb lengthening in, 1769–70, 1773, 1773f–1774f
 - options for, 1762
 - rotationplasty in, 1763, 1768f–1769f
 - surgical principles in, 1770–1771, 1773t, 1776f
- Femoral epiphysis, growth of, 2283–2284, 2284f
- Femoral head
- arterial circulation in, and Legg-Calvé-Perthes disease, 658f, 658–661, 660f–662f
 - avascular necrosis of. *See* Avascular necrosis (AVN).
 - blood supply to, and Legg-Calvé-Perthes disease, 658f
 - displacement of, in cerebral palsy, 1184f
 - double infarcts in, and Legg-Calvé-Perthes disease, 661, 662f, 669–670
 - flattening of, in multiple epiphyseal dysplasia, 1533f–1534f
 - ossification of, delayed, in diastrophic dysplasia, 1538, 1538f
 - osteochondritic lesions in, after treatment of Legg-Calvé-Perthes disease, 701, 704f
 - penetration of fixation devices into, 720, 721f
 - reduction of, in open reduction of hip, medial approach to, 558p–559p
 - reossification in, and Legg-Calvé-Perthes disease, 675, 675t, 677f
 - seating of, in evaluation of closed hip reduction, 544–546, 544f–547f
 - shape of, and prognosis in Legg-Calvé-Perthes disease, 702–704
 - varus deformity of, in developmental coxa vara, 766f
 - venous drainage of, in Legg-Calvé-Perthes disease, 661, 663f
- Femoral head-neck angle, measurement of, Southwick method of, 714, 714f
- Femoral head-neck osteoplasty, for residual deformity from slipped capital femoral epiphysis, 745, 746f
- Femoral hypoplasia-unusual facies syndrome, 1753f, 1754
- Femoral metaphysis, radiographic changes in, in Legg-Calvé-Perthes disease, 675–678, 678f–679f
- Femoral neck
- fractures of. *See* Hip, fractures of.
 - length of, and mechanical stress in coxa vara, 770, 771f
 - venous drainage of, in Legg-Calvé-Perthes disease, 661, 663f
- Femoral neck-shaft angle, in developmental coxa vara, 767–768, 767f–768f, 770, 770f
- Femoral nerve
- injury to, from Salter innominate osteotomy, 621
 - lateral cutaneous, protection of, during Salter innominate osteotomy, 634p–635p
- Femoral nerve (*Continued*)
- palsy of, in hemophilia, 1883–1884
 - protection of, in Barr erector spinae transfer, 1338p–1341p
- Femoral osteotomy
- for Legg-Calvé-Perthes disease, 696–698, 699f–700f
 - for pseudoachondroplasia, 1524
 - varus derotation, for hip subluxation/dislocation, in cerebral palsy, 1191, 1196f
 - with traction, for limb lengthening, 1079, 1079f
- Femoral physes, radiographic changes in, in Legg-Calvé-Perthes disease, 678–679, 680f
- Femoral shaft, fractures of, 2301f–2318f
- anatomic considerations in, 2301f, 2301–2302
 - and leg length inequality, 1040
 - classification of, 2302
 - displacement patterns in, 2303, 2304f
 - mechanism of, 2302, 2303f
 - nonunion of, 2319
 - pathologic, 2302, 2303f
 - physical examination in, 2302
 - radiographic features of, 2302–2303, 2303f–2304f
 - recurrence of, 2308, 2311f
 - treatment of, 2303–2319, 2305f–2318f
 - external fixation in, 2308, 2310, 2310f–2311f
 - in child 6 to 10 years, 2305, 2306f
 - in child 7 months to 5 years, 2304–2305, 2306f
 - in child 11 years to skeletal maturity, 2305
 - in infant 0 to 6 months, 2304, 2305f
 - intramedullary fixation in, 2313–2314, 2313f–2316f
 - intramedullary nailing in, 2314, 2317f–2318f, 2319
 - open reduction/internal fixation in, 2310, 2312f, 2313
 - spica cast in, 2308, 2308f–2309f
 - traction in, 2305, 2307f, 2307–2308
- Femoral varus derotation osteotomy (VDRO), for hip subluxation/dislocation, in cerebral palsy, 1191, 1196, 1196f
- Femoral vein, dissection of, in midthigh amputation, 1984p–1985p
- Femoral vessel(s), retraction of, in open reduction of hip, medial approach to, 552p–555p
- Femoral/innominate osteotomy, for Legg-Calvé-Perthes disease, 699
- Femoral-tibial angle
- measurement of, 26f
 - normal, 1103, 1106f
- Femur
- anatomic axis in, 1103, 1106f
 - anteversion of, 783–786, 784f–786f
 - and intoeing, 66f, 66–67, 783, 785f
 - in cerebral palsy, 1169, 1178f
 - clinical features of, 783, 785f–786f
 - conditions associated with, 784
 - definition of, 783
 - degree of, 783, 784f
 - differential diagnosis of, 783
 - measurement of, 784, 784f
 - natural history of, 783–784
 - prognosis in, 783–784
 - treatment of, 784–786

Femur (*Continued*)
 avulsive cortical irregularities of, 132, 133f
 blood supply to, 2301f, 2301–2302
 bone cyst in, 1902f
 chondromyxoid fibroma of, 1932f
 derotational osteotomy of, 785
 distal, epiphysiodesis of, for leg length inequality, 1064p–1067p
 distal epiphyseal fractures of, 2321–2325, 2322f–2324f
 anatomic considerations in, 2321–2322, 2322f
 classification of, 2322–2323, 2322f–2323f
 complications of, 2322–2324, 2334f–2335f
 diagnosis of, 2325
 mechanism of, 2323–2325, 2324f
 radiographic features of, 2325
 distal metaphyseal fractures of, treatment of, 2325–2327, 2326f–2328f
 closed reduction and internal fixation in, 2326, 2326f
 external fixation in, 2325–2326
 open reduction and internal fixation in, 2326–2327, 2327f–2328f
 distal physial fractures of, treatment of, 2327–2332, 2329f–3833f
 casting in, 2328, 2329f
 closed reduction in, 2328–2330, 2329f–2331f
 goals of, 2327–2328
 open reduction and internal fixation in, 2332, 2333f
 principles of, 2327
 division of, in midthigh amputation, 1988p–1989p
 enchondroma of, angular deformity from, 1104f, 1106, 1107f–1108f
 Erlenmeyer flask deformity of, in Gaucher's disease, 1643, 1644f
 Ewing's sarcoma of, 1848f
 fibrous dysplasia of, 1915f
 fractures of. *See also* Femoral shaft, fractures of.
 and physal arrest, 2066f–2067f, 2069f
 with ipsilateral tibial fractures, 2389f, 2389–2390
 hereditary multiple exostosis of, 1924f
 intramedullary fixation of, Bailey-Dubow rod for, 1733–1734, 1735f–1737f
 lengthening of
 fixation over intramedullary rods for, 1093–1094, 1094f
 for leg length inequality, 1081–1082, 1082f
 complications of, 1089f
 results of, 1091–1092
 timing of, 1056p–1057p
 mechanical axis in, 1103, 1106f
 osteosarcoma of, 1958f–1959f, 1961f–1962f
 prediction of remaining growth in, 1048–1053, 1049t, 1050f–1055f, 1052t, 1056p–1057p
 Anderson-Green-Messner charts for, 1048–1051, 1049t, 1050f–1055f, 1052t
 for timing of epiphysiodesis, 1053–1058, 1056p–1057p, 1058f
 Menelaus method for, 1051, 1055f
 Moseley straight line graphs for, 1051–1053, 1056p–1057p, 1059f
 proximal
 closing wedge valgus osteotomy of, for avascular necrosis, 614p–617p

Femur (*Continued*)
 intertrochanteric oblique osteotomy of, and internal fixation with Coventry apparatus, 1350, 1352p–1355p
 Lloyd Roberts intertrochanteric oblique osteotomy of, and internal fixation with Coventry apparatus, 578p–581p
 osteonecrosis of, from osteomyelitis, 1859, 1860f
 resection of, with interposition arthroplasty, for nonreconstructable hip in cerebral palsy, 1203, 1209f
 valgus osteotomy of, for nonreconstructable hip in cerebral palsy, 1203, 1210f
 pseudarthrosis of, stabilization of, for proximal focal femoral deficiency, 1763, 1772f
 rotational osteotomy of, for cerebral palsy, in knee, 1169–1173, 1178f
 shortening of
 for hip dislocation, in Larsen's syndrome, 1608, 1610f–1611f
 for leg length inequality, 1075f–1076f, 1075–1077
 with open hip reduction, 549, 549f, 568–569, 568f–569f, 570p–581p
 acetabular procedures for, 568–569, 568f–569f
 maximal age for, 569, 582
 results of, in older children, 568f–569f, 569
 solitary osteochondroma of, 1919, 1919f
 vascular anatomy of, 2284, 2284f
 "Fenestrated syndactyly," 1784, 1785f
 Fentanyl, epidural administration of, 114–115
 Fetal movement, decreased, and arthrogryposis, 1647
 Fetal position
 and congenital dislocation of knee, 822–823
 and congenital torticollis, 171
 and developmental dysplasia of hip, 515–516, 515f–516f
 and talipes calcaneovalgus, 905
 during delivery, and brachial plexus palsy, 479–480
 Fetal surgery, for myelomeningocele, 1257
 FGF. *See* Fibroblast growth factor (FGF).
 FGFR (fibroblast growth factor receptor) gene, and achondroplasia, 1508
 Fibroblast culture(s), in mucopolysaccharidosis, 1572
 Fibroblast growth factor (FGF)
 and congenital anomalies of upper extremities, 411, 411t
 and physal growth, 1047
 in limb development, 1745
 Fibroblast growth factor receptor (FGFR) gene, and achondroplasia, 1510
 Fibrocartilaginous anlage, in ulnar deficiency, 413
 Fibrocartilaginous dysplasia, focal, tibia vara from, 855, 857f–858f, 858t
 Fibrodysplasia ossificans, anesthesia in, 123t
 Fibrodysplasia ossificans progressiva, 1632–1638, 1633f–1637f
 clinical features of, 1632–1634, 1633f–1637f
 etiology of, 1632
 histologic features of, 1636–1638
 radiographic features of, 1633f–1637f, 1634
 treatment of, 1638

Fibroma
 aponeurotic, of upper extremities, 502
 chondromyxoid, 1930–1933, 1932f
 nonossifying, 1942–1944, 1942f–1945f
 clinical features of, 1943
 differential diagnosis of, 1943
 natural history of, 1943
 pathology of, 1942, 1942f
 radiographic features of, 1943, 1943f
 treatment of, 1943–1944, 1945f
 vs. chondromyxoid fibroma, 1933
 ossifying. *See* Campanacci's disease.
 Fibrosarcoma, congenital, 2042, 2043f
 Fibrotic tissue, retractive, and talipes equinovarus, 923
 Fibrous ankylosis, of hip, from hemophilic arthropathy, 1882f
 Fibrous cortical defect. *See* Fibroma, nonossifying.
 Fibrous dysplasia, 1909–1913, 1913f–1916f and congenital tibial pseudarthrosis, 867–868
 classification of, 1909, 1911
 clinical features of, 1911, 1914f
 congenital. *See* Campanacci's disease.
 differential diagnosis of, 1911
 etiology of, 1911
 pathology of, 1911, 1913f
 radiographic features of, 1911, 1915f
 treatment of, 1912–1913, 1916f
 vs. Campanacci's disease, 1917
 Fibrous popliteal cyst(s), 830f, 831
 Fibrous tumor(s), of upper extremities, 502
 Fibula
 anatomy of, 2371
 congenital pseudarthrosis of, 883–884, 884f–886f, 887
 distal, fractures of. *See* Ankle, fractures of.
 epiphysiodesis of, for leg length inequality, 1068p–1071p
 fractures of, 2371–2373, 2372f
 diaphyseal, 2376–2381, 2378f–2382f
 classification of, 2376
 diagnosis of, 2377, 2378f
 mechanism of, 2376–2377
 treatment of, 2377–2381, 2381f–2382f
 hemimelia of
 bilateral, 1780–1782
 partial, 1776–1777
 ossification of, 893f
 osteofibrous dysplasia of, 1914–1917, 1917f–1919f. *See also* Campanacci's disease.
 sectioning of, in below-knee amputation, 1998p–1999p
 synostosis to tibia, for tibial deficiency, 1783f
 tenodesis of Achilles tendon to, for valgus ankle deformity, 1268
 Fibular deficiency, 1775–1778, 1776f–1779f, 1777t
 classification of, 1775–1776, 1776f, 1777t
 clinical features of, 1776, 1777f–1779f
 prostheses for, 1802–1803
 treatment of, 1776–1778, 1779f
 Fibular graft(s)
 for free vascularized bone transfer, in forearm, 510f
 vascularized, for congenital tibial pseudarthrosis, 877–878, 878f
 FIM (Functional Independence Measure), 2105f
 Fine motor skill(s), developmental milestones in, 4f, 20t, 20–21

- Finger(s)
 clawing of, in Charcot-Marie-Tooth disease, 1444, 1447f
 flexor tendons of, fractional lengthening of, 1218p–1221p
 fractures of, 2249–2250
 radiography in, 400, 402f
 in cerebral palsy, surgical procedures on, 1213, 1218p–1221p
 in mirror hand, 425f, 426
 in pseudoachondroplasia, 1524, 1524f
 index, duplication of, 448
 multiple enchondromatosis of, 1928f
 nerve injuries to, acute care of, 400
 partial, augmentation of, 1783, 1785f
 skin injuries to, acute care of, 400, 401f
 splinting of, 392–393, 392f–394f
 synovitis of, 472–473, 473f
 tendon injuries in, acute care of, 398–400, 399f–400f
 Fingertip, separation of, in congenital band syndrome, 465f
 Fishtail deformity, from humeral fractures, lateral condylar, 2185, 2188f
 Fixation device(s), penetration into femoral head, 720, 721f
 Fixsen and Lloyd-Roberts classification, of femoral deficiency, 1760
 Flail knee, in poliomyelitis, 1360–1361, 1361f
 Flatfoot
 assessment of, 67f, 67–68
 flexible. *See* Pes planovalgus.
 in Down syndrome, 1623
 peroneal spastic. *See* Tarsal coalition.
 Flexible flatfoot. *See* Pes planovalgus.
 Flexion. *See also under specific joint.*
 definition of, 27t
 Flexion contracture(s). *See also under specific joint, e.g., Knee.*
 physiologic, in neonates, 3, 5f, 36, 38f
 Flexion deformity(ies), of hip, assessment of, 38–39, 39f
 Flexor hallucis brevis tendon, transfer to first metatarsal, for dorsal bunion, 1372–1373, 1379f, 1386f–1387f
 “Floating knee,” 2389f, 2389–2390
 Fluid(s), postoperative, 111
 Fluoroscopic imaging, for placement of sacroiliac screws, 2257, 2268f
 Folate deficiency, and myelomeningocele, 1251
 Foot
 accessory bones in, 895f
 alignment of, tibial torsion and, 1378, 1388f
 amputation of
 prostheses for, 1803
 with knee arthrodesis, for proximal focal femoral deficiency, 1762f, 1762–1763, 1764p–1767p
 arthrodesis of, in poliomyelitis, 1383–1395, 1388f–1389f, 1390p–1393p, 1394 triple, 1383–1387, 1388f–1389f, 1390p–1393p
 calcaneocavus deformity of, from triceps surae muscle paralysis, 1365, 1378f
 calcaneus deformity of, from congenital posteromedial tibial bowing, 883, 883f
 casting of, and prevention of pressure sores, 1262, 1262f
 cavus. *See* Pes cavus.
 cleft, 983f, 983–984
- Foot (*Continued*)
 deficiencies of
 congenital, 901–984, 1782, 1783f. *See also* Talipes.
 postural, 901–907, 902f–906f
 prostheses for, 1803
 equinovarus deformity of, in diastrophic dysplasia, 1534, 1535f, 1537, 1537f
 functions of, 891
 growth of, 891, 893f
 in cerebral palsy. *See* Cerebral palsy, foot and ankle in.
 in myelomeningocele, 1262–1269. *See also* Myelomeningocele, foot and ankle in.
 injuries to, 2414–2434, 2417f–2433f. *See also* Calcaneus; Talus.
 from lawnmowers, 2432, 2434
 movements of, 891
 neurogenic abnormalities of. *See also* Pes cavus, 984–1011, 984f–1011f, 992p–999p
 normal anatomic variations in, 891–899
 ossification of, 891, 892f–893f
 passive stretching of, for metatarsus adductus, 993, 994f–995f
 poliomyelitis and. *See* Poliomyelitis, foot and ankle in.
 pronation of, 44, 45f
 prosthetic, solid ankle-cushioned, for leg length inequality, 1058, 1058f
 puncture wounds of, 1863, 2434
 range of motion of, 43–44, 43f–45f
 stability of, operative procedures for, history of, 1388t
 supination of, 44, 45f
 Foot frame(s), complications of, 953, 957f
 Foot grasp reflex, 46, 50f
 Foot-progression angle, 79, 79f, 863, 865
 Forearm
 compartment syndrome of, 403
 deformities of, from brachial plexus palsy, management of, 486–487, 498p–501p
 finger and wrist flexors in, fractional lengthening of, 1218p–1221p
 fractures of, 2218–2241, 2220f–2240f. *See also* Radius; Ulna.
 both-bone. *See* Radius, fractures of, with ulnar fractures.
 distal, 2233–2241, 2235f–2240f
 anatomic considerations in, 2233
 classification of, 2235, 2235f
 complications of, 2238–2241, 2239f
 diagnosis of, 2233–2235
 mechanism of, 2233
 radiographic features of, 2235
 treatment of, 2235–2238, 2236f–2237f
 greenstick, 2059f
 Monteggia, 2218–2225, 2220f–2224f. *See also* Monteggia fracture(s).
 nonunion of, in congenital insensitivity to pain, 1450f
 in poliomyelitis, 1424–1427
 muscles of, dissection of, in elbow disarticulation, 2022p–2025p
 postoperative position of, in radioulnar synostosis, 415–416
 pronation contractures of, in cerebral palsy, 1212, 1213f
 pseudarthrosis of, free vascularized bone transfer for, 508, 509f–511f, 512
 range of motion of, 31, 33f
 splints for, 396, 397f
 ulnar deficiency in. *See* Ulnar deficiency.
- Forearm (*Continued*)
 volar fasciotomy of, 2076f
 Volkmann’s contracture of, 408, 409f–410f, 410. *See also* Volkmann’s ischemic contracture.
- Forefoot
 equinus deformity of, in pes cavus, 986, 987f
 plantar flexion deformity of, in pes cavus, 985, 985f
 Foreign body(ies), in upper extremities, 497
 Forequarter(s), amputation of, posterior approach to, 2000p–2013p
 Forward bending test, Adams’ in Scheuermann’s kyphosis, 324, 328f
 in scoliosis, 222
 Four-flap Z-plasty, for thumb-index web syndactyly, 444, 445f
 Fracture(s). *See also under specific site, e.g., Forearm, fractures of; under specific type, e.g., Pathologic fracture(s).*
 burst
 of cervical spine, 2096–2097
 treatment of, 2105–2106
 cerebral palsy and, 1230, 1232
 classification of, 2064–2065, 2064f–2065f
 compression. *See* Compression fracture(s).
 epidemiology of, 2059
 hangman’s, 2098, 2099f
 in pyknodysostosis, 1567
 in osteogenesis imperfecta, 1722f, 1723–1724
 Jefferson, 2094
 nonunion of
 in congenital insensitivity to pain, 1450f
 in osteogenesis imperfecta, 1731
 open, 2074t, 2074–2075
 risk of, myelomeningocele and, 1262, 1263f
 vascular injuries from, 2078f, 2078–2079
 Fracture callus, development of, for limb lengthening, 1083, 1085f
 Fracture table(s), for percutaneous in situ pinning of slipped capital femoral epiphysis, 722–723, 723f, 724p–727p
 Fragilitas ossium, anesthesia in, 124t
 Frantz and O’Rahilly classification of fibular deficiency, 1775
 of limb deficiencies, 1747–1748, 1747f–1749f
 Freckling, in neurofibromatosis, 1625, 1627f
 Free joint transfer, in radial clubhand, 511, 511f
 Free motorized muscle transfer, microsurgery for, 504, 507f
 Free tissue transfer, microsurgery for, 504, 505f–506f
 Free toe transfer, microsurgery for, 504, 508, 508f
 Free vascularized bone transfer, microsurgery for, 508, 509f–511f, 512
 Freeman-Sheldon syndrome, 1662, 1663f
 Freiberg’s infarction, 900–901, 901f
 Frejka pillow, 538, 538f
 French technique, of manipulation, for talipes equinovarus, 929, 933f–934f
 Friedreich’s ataxia, 1246–1248, 1247f
 anesthesia in, 121t
 Frog leg radiographic view, in slipped capital femoral epiphysis, 719, 720f
 “Frog-leg” deformity, in myelomeningocele, 1256
 Functional Independence Measure (FIM), 2105f

Functional training, for poliomyelitis, during chronic phase, 1326
 Fungal osteomyelitis, 1869–1870, 1870f

Gadolinium, for magnetic resonance imaging, 156
 GAG(s) (glycosaminoglycans), urinary, in mucopolysaccharidosis, 1572

Gait analysis of
 in adolescent, 85–86
 in cerebral palsy, 1126f–1129f, 1126–1130 with equinovarus deformity, 1142–1143, 1143f
 in school-age child, 85
 in toddlers, 85, 86f
 laboratories for, 81f, 82
 purpose of, 1130
 ankle dorsiflexion and, 77, 78f
 antalgic, 83
 in Legg-Calvé-Perthes disease, 672f
 cadence parameters in, 75–76, 76t
 calcaneus, after talipes equinovarus surgery, muscle transfers for, 946, 947f
 crouch, in cerebral palsy, 1158–1159, 1166f correction of, 1158–1159, 1168f, 1170p–1171p
 determinants of, 77, 77t
 developmental milestones in, 20
 energy for, 76–77, 77t
 function of, 76
 in coxa vara, 89, 90f
 in developmental dysplasia of hip, 89, 90f
 in Duchenne muscular dystrophy, 1467, 1468f
 in femoral anteversion, 783, 785f
 in hypertrophic interstitial neuritis, 1445
 in Legg-Calvé-Perthes disease, 89–90, 91f
 in myelomeningocele, level of lesion and, 1256–1257
 in spastic diplegia, 81f
 kinematics and, 77–79, 78f–79f
 in coronal plane, 77–78, 78f
 in sagittal plane, 77, 78f–79f
 in transverse plane, 78–79, 79f
 kinetics and, 80f, 81
 leg length inequality and, 84, 1040–1041
 limping. *See Limp.*
 muscle activity and, 79t, 79–81, 80f
 neurologic control of, 76
 pathologic, 81, 81f
 problems with, from lower extremity prostheses, 1805–1806
 short-limb, 84
 spastic, 84
 stance phase of, 75, 76f
 muscle activity during, 80–81
 swing phase of, 75, 76f
 muscle activity during, 80–81
 temporal aspects of, 75–76, 76t
 toe-in, persistent, with talipes equinovarus, supramalleolar osteotomy for, 950–951, 952f
 Trendelenburg. *See Trendelenburg gait.*
 with proximal muscle weakness, 84, 85f
 β -Galactosidase, deficiency of, in Morquio syndrome, 1577
 Galant's reflex, 52, 54f
 Galeazzi fracture(s), 2235, 2238
 Galeazzi procedure, for patellar dislocation, 2353, 2353f

Galeazzi sign, in developmental dysplasia of hip, 525–526, 525f–526f
 Gallie fusion for atlantoaxial arthrodesis, 187f for atlantoaxial instability, 2092, 2094f–2095f
 Galveston technique, for spinal fusion, in cerebral palsy, 1227, 1230f
 Gamma rays(s), in nuclear imaging, 166
 Ganglion(s), of upper extremities, 496–497
 Ganglioside, for spinal cord injury, GM₁, 2106–2107
 Gangliosidosis, anesthesia in, 125t
 Gangrene, in purpura fulminans, 1792
 Ganz osteotomy, for acetabular repositioning, 633, 642, 643f
 Gardner's syndrome, anesthesia in, 121t
 Gastrocnemius muscle bursa of, popliteal cyst in, 829, 829f recession of, for equinus deformity, in cerebral palsy, 1136, 1137f–1139f
 Gastrosoleus muscle botulinum toxin injection of, for spasticity, in cerebral palsy, 1133–1134 contraction of, during gait, 79–80 pseudohypertrophy of, in Duchenne muscular dystrophy, 1468, 1468f
 Gaucher's disease, 1642–1645, 1643f–1644f clinical features of, 1643 diagnosis of, 1643 genetic basis of, 1642 orthopaedic manifestations of, 1643–1645, 1644f pathology of, 1642–1643, 1643f treatment of, 1645
 Gender and incidence of slipped capital femoral epiphysis, 711–712 and interpretation of radiographs, 127–128 and natural history of scoliosis, 215
 Gene therapy for Duchenne muscular dystrophy, 1477 for hemophilia, 1879
 Generator(s), for radiography, 133
 Genital(s), development of, in females, 19, 19f
 Genitourinary tract, injuries to, with pelvic fractures, 2255–2256
 Genu recurvatum acquired, 825, 828–829 after biceps femoris tendon transfer, 1357 in poliomyelitis, 1358–1360, 1359f–1360f
 Genu valgum, 855–863, 859f–866f degree of, measurement of, 26 hypoplastic tibial epiphysis and, 1680, 1680f in diastrophic dysplasia, 1537, 1538f in metaphyseal dysplasia, 863 in Morquio syndrome, 1579, 1579f in proximal tibial fracture, 861, 864f in pseudoachondroplasia, 1524, 1524f in rickets, 861, 863 in spondyloepiphyseal dysplasia, 861, 865f physiologic, 855–863 hemiphiodesis for, 855, 858–860, 860f–861f stapling in, 859–860, 862f timing of, 855, 858
 osteotomy for, 860–861, 863f
 Genu varum, 839–855, 840f–858f. *See also Tibia vara.*
 and tibia vara, 840–855, 840f–858f. *See also Tibia vara.*
 degree of, measurement of, 26 in achondroplasia, surgical treatment of, 1517, 1518f

Genu varum (*Continued*)
 in metaphyseal chondrodysplasia, 1547–1548, 1548f
 in multiple enchondromatosis, 1929f
 in pseudoachondroplasia, 1524, 1524f
 in vitamin D-resistant rickets, 1691f–1693f physiologic, 839–840, 840f–842f vs. infantile tibia vara, 844
 Geographic location, and incidence of slipped capital femoral epiphysis, 711
 Gestation, multiple, and incidence of cerebral palsy, 1123
 Gestational age, and incidence of cerebral palsy, 1123
 Giant cell(s) in aneurysmal bone cyst, 1908, 1908f in nonossifying fibroma, 1942, 1942f
 Giant cell tumor(s), of tendon sheath, in upper extremities, 502
 Gigantism cerebral, anesthesia in, 125t digital. *See Macrodactyly.* infantile, anesthesia in, 120t localized, 1024, 1025f
 Gigli saw, placement of, in Salter innominate osteotomy, 636p–639p
 Gillespie and Torode classification, of femoral deficiency, 1759–1760
 Gillespie classification, of femoral deficiency, 1760, 1760f
 Glasgow Coma Scale, 2073, 2073t
 Glenohumeral joint range of motion of, assessment of, 28f, 29–31, 30f–31f traumatic dislocation of, 2128–2132, 2129f–2131f anatomic considerations in, 2128–2129, 2129f complications of, 2131–2132 diagnosis of, 2129–2130, 2130f mechanism of, 2129 treatment of, 2130–2131, 2130f–2131f
 Gloma, optic, in neurofibromatosis, 1625, 1628
 Glomus tumor(s) of toe, 1032 of upper extremities, 502
 Glucocerebrosidase, for Gaucher's disease, 1645
 Gluteus maximus muscle dissection of, in Banks and Coleman hemipelvectomy, 1968p–1969p injections into, and sciatic nerve palsy, 1455–1456
 paralysis of, in poliomyelitis, 1336–1337, 1348–1349, 1348f–1349f Barr erector spinae transfer for, 1336–1337, 1348, 1348f Hogshead fascia lata transfer to lumbodorsal fascia in, 1349, 1349f motor evaluation in, 1336
 Gluteus medius muscle dissection of in hip disarticulation, 1976p–1977p in trochanteric transfer, for avascular necrosis, 602p–603p, 606p–607p paralysis of, in poliomyelitis, 1336, 1338p–1347p
 Gluteus minimus muscle, dissection of, in hip disarticulation, 1976p–1977p
 Gluteus muscle(s), dissection of, in open reduction of hip, using anterolateral approach, 562p–563p
 Glycosaminoglycans (GAGs), urinary, in mucopolysaccharidosis, 1572

- GM₁ ganglioside, for spinal cord injury, 2106–2107
- Goldenhar's syndrome, anesthesia in, 121t
- Gonad(s), shielding of, during radiography, 135–137, 136f–137f
- Goniometry, 27t, 27–28
- in assessment of angular deformities, 25–26, 26f
- Gonococcal arthritis, 1835–1836, 1864–1865
- Gonosomal aneuploidy, anesthesia in, 122t
- Gorham's osteolysis, 1570–1571, 1571f
- Gotttron's sign, in dermatomyositis/
polymyositis, 1493, 1494f
- Gowers' sign, 84, 85f, 1463, 1464f
- in Duchenne muscular dystrophy, 1468
- Gracilis muscle
- for free toe transfer, 504, 508f
 - transfer of
 - for elbow flexion, in arthrogryposis, 1655
 - for Volkmann's contracture of forearm, 408, 409f
- Gracilis tendon, transfer of rectus femoris tendon to, 1174p–1177p
- Graf classification, of developmental dysplasia of hip, 527–530, 529f, 529t
- Granuloma
- eosinophilic. *See* Eosinophilic granuloma.
 - giant cell reparative, 1908, 1908f
 - pyogenic, of upper extremities, 502
- Grasp reflex
- palmar, 46, 50f
 - plantar, 46, 50f
- Great toe, malformations of, in fibrodysplasia ossificans progressiva, 1632, 1633f
- Greater trochanter, epiphysiodesis of, for developmental coxa vara, 772–773
- Green and Grice procedure, for posterior tibialis tendon transfer, for calcaneus deformity, 1380p–1385p
- Green modification, Phemister technique, of epiphysiodesis, for leg length inequality, 1058–1063, 1062f, 1064p–1071p, 1072f
- Green-Anderson growth remaining chart, 2068f
- Green's quadricepsplasty, for recurrent patellar dislocation, 798, 806p–809p
- Greenstick fracture(s), 2058, 2059f
- creation of, in closing wedge valgus osteotomy, 102p–103p
 - of distal forearm, 2059f
 - treatment of, 2235
- Grice procedure, 1396p–1399p
- for vertical talus, 967
- Grid(s), radiographic, 133
- Grieving process, stages of, 381
- Gross motor skill(s), developmental milestones in, 4f, 19–20, 20t
- Ground-glass appearance, of bone, in renal osteodystrophy, 1697, 1697f
- Ground-reaction ankle-foot orthoses
- for cerebral palsy, 1131
 - for myelomeningocele, 1292, 1293f
- Growth
- and head circumference, 10, 11f–12f
 - and height, 10, 13f–18f
 - in boys, 10, 13f, 15f, 17f
 - in girls, 10, 14f, 16f, 18f
 - and weight
 - in boys, 10, 13f, 15f, 17f
 - in girls, 10, 14f, 16f, 18f
- Growth (*Continued*)
- definition of, 3
 - disturbance of, from phyeal injuries, 2066–2067, 2066f–2070f, 2070
 - epiphyseal, 6–10f, 10
 - normal, 3, 5, 5f
 - of upper extremity, cerebral palsy and, 1211
 - patterns of, and prediction of leg length inequality, 1051–1053, 1058f–1060f
 - prediction of, Anderson-Green-Messner charts for, 1048–1051, 1049t, 1050f–1055f, 1052t
 - remaining, calculation of, 2070f
- Growth and development
- disorders of, 5
 - history of, 24
 - in Legg-Calvé-Perthes disease, 661–662, 664f
 - normal, 3–5, 5f
- Growth arrest, 2066–2067, 2066f–2069f
- after distal forearm fractures, 2238, 2239f
 - after femoral physeal fractures, 2334, 2335f
 - after hip fractures, 2297, 2300
 - after humeral lateral condylar fractures, 2185
 - from proximal tibial physeal fractures, 2351
 - in proximal femoral epiphysis, 2283–2284, 2284f
 - tibial, after ankle fractures, 2403–2405, 2404f, 2406f
- Growth arrest line(s), Harris, 2063–2064, 2064f
- Growth hormone
- and physeal growth, 104f
 - deficiency of, and slipped capital femoral epiphysis, 715–716
 - for achondroplasia, 1517, 1519f
 - for hypochondroplasia, 1521
 - for pituitary dwarfism, 1712
 - for vitamin D-resistant rickets, 1694
- Growth plate(s)
- capital femoral, closure of, from avascular necrosis, 583
 - destruction of, in developmental dysplasia of hip, 592p–593p
 - in epiphyses of lower extremities, closure of, age at, 9f
 - in epiphyses of upper extremities, closure of, age at, 7f
 - pathology of, in slipped capital femoral epiphysis, 717f, 717–718
 - removal of, in epiphysiodesis, for leg length inequality, 1066p–1067p, 1070p–1071p
 - V-shaped, in achondroplasia, 1512, 1512f
- Growth plate bar(s), 158f
- Growth potential, in scoliosis, preoperative assessment of, 239–240, 239f–240f
- Growth spurt, during puberty, 1048, 1050f
- Guide wire, placement of, for slipped capital femoral epiphysis
- during *in situ* open pinning, 728, 729f
 - during percutaneous *in situ* pinning, 722, 724p–727p
 - in bone graft epiphysiodesis, 731, 732p–733p
 - in Kramer and Barmáda osteotomy, 742p–743p
- Guiding neuromuscular unit(s), in
- poliomyelitis, 1322
- Guillain-Barré syndrome, 1454–1455
- anesthesia in, 122t
- Gunshot wound(s), 2075
- Gustilo and Anderson classification, of open tibial fractures, 2381
- Hallux flexus, after treatment of talipes equinovarus, 943–945, 945f
- Hallux rigidus, 1019, 1021, 1021f
- Hallux valgus, 1012f–1018f, 1012–1019
- angle in, 1014, 1014f
 - classification of, 1014
 - clinical features of, 1012–1013, 1012f–1013f
 - dorsal, after treatment of talipes equinovarus, 943–945, 945f
 - in cerebral palsy, 1158, 1166f
 - incidence of, 1012
 - interphalangeus, 1019, 1019f
 - pathologic anatomy of, 1013f, 1013–1014
 - radiographic features of, 1014, 1014f
 - treatment of, 1014–1019, 1015f–1018f
 - Akin procedure in, 1018
 - chevron osteotomy in, 1016, 1017f
 - choice of procedure in, 1014
 - fusion of first metatarsophalangeal joint in, 1018f, 1018–1019
 - metatarsal osteotomy in, 1016, 1017f, 1018
 - Mitchell osteotomy in, 1015–1016, 1016f
 - soft tissue realignment in, 1015
 - splinting for, 1014, 1015f
- Hallux varus, 1019, 1020f
- Halo cast
- for cervical kyphosis, 202, 202f
 - for Klippel-Feil syndrome, 183
 - for occipitoatlantal dislocation, 2092, 2093f
 - for odontoid fractures, 2096
- Halo vest, for atlantoaxial rotatory displacement, 189, 190f–192f
- Halo-Ilizarov distraction, for diastrophic dysplasia, 196f
- Hamanishi classification, of femoral deficiency, 1758–1759, 1760f
- Hamartoma
- of nerves, in upper extremity, 502
 - vs.* macrodactyly, 451–452
- Hammer toe(s), 1027, 1034p–1035p
- Hamstrings
- lengthening of
 - for iliotibial band contractures, in poliomyelitis, 1331, 1332p–1335p
 - in cerebral palsy, 1159, 1168f, 1170p–1171p
 - proximal, release of, for subluxation/dislocation of hip, in cerebral palsy, 1191, 1192p–1195p
- Spasticity of
- in cerebral palsy. *See Cerebral palsy, knee in, hamstring spasticity from.*
 - measurement of, 1159, 1167f
 - tightness of, in spondylolisthesis, 354
- Transfer to femoral condyles, Eggers procedure for, 1159, 1168f
- Weakness of, after femoral shortening, 1076–1077
- Hand
- acute care of, 398–404, 399f–403f
 - in bone injury, 400–401, 402f–403f
 - in nerve injury, 400
 - in skin injury, 400, 401f
 - in tendon injuries, 398–400, 399f–400f
 - in vascular injury, 401–404, 403
 - cleft, with central polydactyly, 423, 423f
 - compartment syndrome of, 404
 - deformities of
 - from brachial plexus palsy, 483–484
 - management of, 487
 - in Apert's syndrome, 1615

- Hand (Continued)**
- dressings for, 382–387, 385f–387f
 - gentle compression components of, 385–386, 386f
 - padding and skin protection for, 386, 386f–387f
 - primary, 385, 385f–386f
 - principles of, 382–383
 - rigid outer shell, 386–387, 387f
 - “Steri-strip stent,” 386f
 - fixed unit of, 405, 405f
 - fracture-dislocations of, 2249–2250
 - fractures of, treatment of, 400–401, 403f
 - growth of, 382, 382t
 - hemihypertrophy of, in Proteus syndrome, 1666, 1668f
 - hemophilic pseudotumor of, 1885f
 - hypertrophic scarring of, 382, 383f
 - in arthrogryposis, 1655, 1660
 - in macrodactyly, type IV, 456, 456f
 - in Morquio syndrome, 1578f, 1579
 - infections of, 476f–478f, 476–479
 - and felon, 477, 477f
 - and herpetic whitlow, 477, 477f
 - and osteomyelitis, 479
 - and paronychia, 476f, 476–477
 - and septic arthritis, 478–479
 - conditions mimicking, 479
 - deep space, 478
 - mycobacterial, 479
 - papillomavirus, 479
 - inflammatory conditions of, noninfectious, 476
 - injury to, 110p–117p, 118f–119f, 120p–123p, 479–487, 482f–486f
 - lengthening of, 1783, 1785f
 - mirror, 423–426, 425f
 - treatment of, 432
 - mobile compartments of, 405, 406f–407f
 - myoelectric prostheses for, 1800–1801
 - papillomavirus infections of, 479
 - reconstruction of, 404–410
 - anatomic considerations in, 405, 405f–407f
 - evaluation for, 407–408
 - in Apert’s syndrome, 467
 - physiology and, 404
 - planning of, 405–406
 - sensibility and, 404f, 404–405
 - rheumatoid arthritis of, 1817, 1817f
 - treatment of, 473–474, 474f–475f
 - skin incisions in, placement of, 383f
 - splints for, 387–398, 388f–398f
 - materials for, 387–389, 388f–389f
 - mechanical principles of, 387, 392–398, 392f–398f
 - involving fingers, 392–393, 392f–394f
 - involving thumb, 394, 395f–396f, 396
 - involving wrist, 394, 395f–397f, 396
 - securing, 390, 390f
 - total contact molding inserts with, 390f–391f, 390–392
 - with rheumatoid arthritis, 473–474
 - surgical procedures on
 - timing of, 380–382, 381t–384t, 383f
 - wound closure after, 384–385, 385f
 - tenosynovitis of, pyogenic, 478, 478f
 - trident, in achondroplasia, 1511, 1511f
 - ulnar deficiency in, treatment of, 413. *See also* Ulnar deficiency.
 - Hand grasp reflex, 46, 50f
 - Hand-foot syndrome, 1864
 - in sickle cell disease, 1892f–1893f, 1893
- Hand-heart syndrome**, 1752
- Hand-Schüller-Christian disease**, 1938, 1942
- anesthesia in, 122t
 - Hangman’s fracture, 2098, 2099f
 - in pyknodysostosis, 1567
 - Hardcastle classification, of Lisfranc’s fractures, 2426–2427, 2427f
 - Harrington rod(s)
 - for idiopathic scoliosis, 235–236, 236f–238f, 241f, 241–242
 - for Scheuermann’s kyphosis, 331, 335
 - for Wisconsin segmental spinal instrumentation, 242, 242f, 244p–245p
 - Harris growth arrest line(s), 2063–2064, 2064f
 - Hawkins sign, 2420
 - H-E (Hilgenreiner-epiphyseal) angle, in developmental coxa vara, 767f, 767–768
 - and need for treatment, 771
 - and risk of recurrence, 772–773
 - Head and neck, blood supply to, 1755f
 - Head circumference, changes in, 10, 11f–12f
 - Head injury, 2072, 2073f
 - Head tilt, in Klippel-Feil syndrome, 173–174, 178f
 - Head-at-risk sign(s), in Catterall classification of Legg-Calvé-Perthes disease, 683
 - Head-shaft angle, femoral, in developmental coxa vara, 767f, 767–768
 - Head-to-trunk proportion, evolution of, 5, 5f, 10
 - Healing index, after corticotomy with callotasis, 1083–1084
 - Hearing loss, in osteogenesis imperfecta, 1725
 - Heart disease, congenital, scoliosis from, 308, 309f
 - Heart-hand syndrome, anesthesia in, 122t
 - Heat
 - for tenosynovitis, of wrist, 470
 - moist, during acute phase of poliomyelitis, 1323–1324
 - Heel
 - equinus deformity of, in vertical talus, 964f
 - valgus deformity of, in vertical talus, 960f
 - Heel cord
 - lengthening of
 - for Achilles tendon contracture, 915
 - for correction of vertical talus, 966, 968p–969p
 - tight, 68
 - Height, in relation to age, 10, 13f–18f
 - for boys, 10, 13f, 15f, 17f
 - for girls, 10, 14f, 16f, 18f
 - Heliotropic eyelid(s), in dermatomyositis/polymyositis, 1493f, 1494
 - “Helmet head,” in osteogenesis imperfecta, 1724
 - Hemangioma, of upper extremities, 502
 - Hemarthrosis, in hemophilia, 1880
 - treatment of, 1886
 - Hemiepiphyseal stapling, for angular deformities of leg, 1111, 1112f–1113f
 - Hemiepiphysiodesis
 - convex, for congenital scoliosis, 287
 - for congenital fibular pseudarthrosis, 887
 - for genu valgum, 855, 858–860, 860f–861f
 - stapling in, 859–860, 862f
 - timing of, 855, 858
 - for idiopathic scoliosis, infantile, 278
 - for valgus ankle deformity
 - in cerebral palsy, 1153, 1158
 - in myelomeningocele, 1268
- Hemihypertrophy**
- and leg length inequality, 1039
 - prediction of remaining growth in, 1051, 1055f
 - of extremities
 - in neurofibromatosis, 1628, 1631f
 - macrodactyly with, 451–452, 453f, 458–459
- Hemimelia**, 1748, 1749t. *See also under specific bone, e.g., Fibula.*
- Hemimyelodysplasia**, in myelomeningocele, 1291f
- Hemiparesis**, and leg length inequality, 1042, 1042f
- Hemipelvectomy**, Banks and Coleman, for osteosarcoma, 1964p–1971p
- Hemiplegia**
- definition of, 44
 - in cerebral palsy, 1124, 1124f–1125f
 - spastic, with equinovarus deformity, 1142, 1142f
- “Hemi-spina bifida,” 1291f
- Hemivertebra(e)**
- in congenital scoliosis, 283–284, 284f, 286f
 - excision of, 1503f
 - in myelomeningocele, 1289f
- Hemodilution**, acute normovolemic, 117
- Hemoglobinopathy(ies)**, and Legg-Calvé-Perthes disease, 656–657
- Hemophilia**, 1879–1889, 1881f–1885f, 1882t
- arthropathy in, 1880–1881, 1881f–1882f, 1882t
 - chronic, 1887
 - classification of, 1880–1881, 1882t
 - clinical features of, 1880
 - differential diagnosis of, 1880
 - pathophysiology of, 1880
 - radiographic features of, 1880–1881, 1881f–1882f, 1882t
 - site of, 1880
 - subacute, 1886–1887
 - classification of, 1879
 - clinical features of, 1879–1880
 - dislocations in, 1884–1885
 - fractures in, 1884
 - gene therapy in, 1879
 - hemarthrosis in, 1880
 - treatment of, 1886
 - historical perspective on, 1879
 - incidence of, 1879
 - myositis ossificans in, 1885
 - nerve palsy in, 1883–1884
 - pseudotumor in, 1884, 1885f
 - treatment of, 1889
 - soft tissue bleeding in, 1881–1883, 1883f
 - treatment of, 1885–1889
 - medical, 1885–1887
 - surgical, 1887–1889
- Hemophilus influenzae** infection(s), and osteomyelitis, 1845
- Hemorrhage**
- in muscle, from hemophilia, 1882f, 1882–1883
 - treatment of, 1886
 - periventricular, in cerebral palsy, 1124
- Hereditary multiple exostosis**, 1920–1926, 1921f–1925f
- clinical features of, 1922–1923, 1922f–1924f
 - common sites of, 1922, 1922f
 - differential diagnosis of, 1923–1924
 - etiology of, 1921
 - pathology of, 1921
 - radiographic features of, 1921f–1924f, 1923

- Hereditary multiple exostosis (*Continued*)
 sarcomatous change in, 1924, 1926
 treatment of, 1924, 1925f
- Hereditary onycho-osteodysplasia, 1599f–1601f, 1599–1602
- Hereditary progressive ophthalmomiarthropathy, 1593
vs. Marfan's syndrome, 1590
- Hereditary spastic paraparesis, 1243–1244
- Heredopathia atactica polyneuritiformis, 1447–1448
- Herpetic whitlow, 477, 477f
- Heterotopic ossification, in fibrodysplasia ossificans progressiva, 1634, 1635f–1636f
- Heyman tenodesis, for genu recurvatum, in poliomyelitis, 1359
- Heyman-Herndon release, of tarsometatarsal joint, for metatarsus adductus, 903, 906f
- Hibbs' angle, in pes cavus, 988, 990f
- Hilgenreiner-epiphyseal (H-E) angle, in developmental coxa vara, 767f, 767–768
 and need for treatment, 771
 and risk of recurrence, 772–773
- Hilgenreiner's line, in developmental dysplasia of hip, 531, 532f
- Hinge abduction, of hip, in Legg-Calvé-Perthes disease, 682, 685f
- Hip
 abduction contractures of, Ober's test for, 40, 41f
 abduction of, 39–40, 40f
 hinge, in Legg-Calvé-Perthes disease, 682, 685f
 in Legg-Calvé-Perthes disease, 672, 673f
 abduction/external rotation contractures of, in myelomeningocele, 1272–1273
 abductor muscles of, contraction of, during gait, 80
 adduction contractures of, in cerebral palsy, surgical treatment of, 1173, 1178–1179, 1179f, 1180p–1183p, 1184, 1185f, 1187–1189
 adduction of, 40, 40f–41f
 arthrodesis of
 for avascular necrosis, 753, 756p–757p
 from slipped capital femoral epiphysis, 753, 756p–757p
 for paralytic dislocation, 1350–1351, 1356f
 for proximal focal femoral deficiency, 1770, 1773f–1774f
 aspiration of, in acute transient synovitis, 1825
 closed reduction of, 540–546, 541f, 542p–543p, 544f–548f
 arthrography in, 544f
 evaluation of, 544–546, 544f–547f
 technique of, 540–541, 542p–543p
 zone of safety for, 541, 541f, 544
 concentric reduction of, with Salter innominate osteotomy, 618
 degenerative joint disease of, Legg-Calvé-Perthes disease and, 703–704
 development of
 in developmental dysplasia, 520–522, 520f–523f
 normal, 518f–520f, 519–520
 developmental dysplasia of, 513–650
 clinical features of, 524–526, 524f–527f
 in infant, 525–526, 525f–526f
 in neonate, 524–525, 524f–525f
 in walking child, 526–527, 527f
 Hip (*Continued*)
 complications of
 acetabular dysplasia as, 587–596, 619f
 avascular necrosis as, 569f, 582–584, 582–586f, 587t. *See also* Avascular necrosis (AVN).
 inadequate reduction and redislocation as, 586–587, 618f
 conditions associated with, 517–519, 517f–519f
 definition of, 513
 diagnosis of
 arthrography in, 532–533, 534f
 Klasic test in, 526, 526f
 magnetic resonance imaging in, 533–534
 radiography in, 530–532, 532f–533f, 533t
 acetabular index and, 531, 532f, 533t
 acetabular teardrop figure and, 531–532, 533f
 center-edge angle and, 531, 532f
 Hilgenreiner's line and, 531, 532f
 Severin classification and, 532, 533t
 Von Rosen view and, 531
 ultrasonography in, 164, 164f, 527–530, 528f–531f, 529t
 advantages of, 527
 anatomic landmarks in, 528f
 Graf classification and, 527–530, 529f, 529t
 sensitivity of, 528–530, 530f
 etiology of, 514–517, 515f–517f
 ligamentous laxity in, 514–515, 515f
 postnatal positioning in, 516–517, 517f
 prenatal position in, 515–516, 515f–516f
 evolution of, 513
 history of, 513–514, 514f
 in child ages 6 months to 2 years
 closed reduction for, 540–546, 541f, 542p–543p, 544f–548f. *See also* Hip, closed reduction of.
 guidelines for, 539
 open reduction for, 546–568,
 550p–567p, 568f–569f, 570p–581p.
 See also Hip, open reduction of.
 traction for, 539–540, 539f–540f
 in Marfan's syndrome, 1590–1591
 in neonate
 Pavlik harness for, 534–536, 535f–537f, 535t. *See also* Pavlik harness.
 physical examination in, 524–525, 524f–525f
 in young child
 Pavlik harness for, 536–537, 538f
 splints for, 537–538, 538f
 in young child ages 1–6 months
 braces for, 537–538, 538f
 Frejka pillow for, 538, 538f
 triple diapers for, 538
 incidence of, 513, 514t
 natural history of, 522–524, 523f–524f
 after neonatal period, 523–524, 523f–524f
 with neonatal instability, 522–523
 pathophysiology of, 517f–523f, 519–522
 barriers to reduction in, 520–522, 521f–522f
 hip development in, 520–522, 520f–522f
 subluxation in, 523–524, 524f
- Hip (*Continued*)
 screening for, 63–65, 64f, 534
 treatment of
 complications of, 569f, 582f–586f, 582–596, 587t
 in child 2 years and older, 566f–567f, 568–569, 582
 in child ages 6 months to 2 years, 539–568, 539f–541f, 542p–543p, 544f–549f, 550p–567p, 568f–569f, 570p–581p
 in neonate, 534–536, 535f–537f, 535t
 in young child, 536–539, 538f
 noncompliance with, 630f
 reconstructive procedures for, 596–646, 618t, 622p–630p
 Pemberton osteotomy, 596–597, 618, 620f, 622p–630p
 Trendelenburg gait in, 84, 84f, 526, 527
 disarticulation of, for osteochondroma, 1972p–1981p
 dislocation of, 2273f–2282f, 2273–2283
 anatomic considerations in, 2273, 2273f
 anterior, in cerebral palsy, 1203, 1210f
 avascular necrosis after, 2280–2281, 2282f
 vs. Legg-Calvé-Perthes disease, 690
 bilateral, 526, 526f–527f, 569, 569f
 classification of, 2273, 2274f
 complications of, 2280–2283, 2282f
 diagnosis of, 2274–2275, 2275f
 high-riding, 522, 522f
 in arthrogryposis, 1651–1652
 in diastrophic dysplasia, 1539, 1541
 in hemophilia, 1884–1885
 in Larsen's syndrome, treatment of, 1608, 1608f–1611f, 1610
 mechanism of, 2273–2274
 paralytic, 1349–1351, 1352p–1355p
 arthrodesis for, 1350–1351, 1356f
 varization osteotomy for, 1349–1350, 1352p–1355p
 radiographic features of, 2275, 2276f–2277f
 recurrent, 2281
 teratologic, 138f, 513, 646, 650
 treatment of, 2275, 2278f–2281f, 2280
 Allis closed reduction in, 2275, 2278f, 2280f
 Bigelow circumduction in, 2275, 2278f
 open reduction in, 2280, 2281f
 Stimson gravity technique in, 2275, 2279f
 traction in, 2275, 2280
 docking of, 546
 dysplasia of
 in Charcot-Marie-Tooth disease, 1441, 1444f
 in Down syndrome, 1621f–1622f, 1621–1623
 in talipes calcaneovalgus, 905
 embryology of, 517–520, 518f–520f
 extension of, 38–39, 38f–39f
 and gait, 77, 78f
 fibrous ankylosis of, from hemophilic arthropathy, 1882f
 flexion deformities of
 assessment of, 38–39, 39f
 in myelomeningocele, 1273
 flexion of, 38–39, 38f–39f
 and gait, 77, 78f
 inadequate, in Pavlik harness, 535, 546f
 fractures of, 2283–2300, 2284f–2291f
 anatomic considerations in, 2283–2285, 2284f–2285f

Hip (*Continued*)
 cervicotrochanteric, 2285, 2287f
 treatment of, 2291–2292, 2292f
 classification of, 2285–2286, 2285f–2288f
 complications of, 2293–2400, 2294f–2299f
 avascular necrosis as, 2288, 2289f,
 2293–2306, 2294f–2307f
 coxa vara as, 2296, 2298f
 growth arrest as, 2297, 2300
 infection as, 2300
 nonunion as, 2296–2297, 2299f
 premature phseal arrest as, 2297,
 2300
 diagnosis of, 2286–2287
 intertrochanteric, 2285, 2287f
 treatment of, 2292, 2293f
 mechanism of, 2286
 radiographic features of, 2287
 stress, 2286, 2288f
 treatment of, 2292–2303
 transcervical, 2285, 2286f
 treatment of, 2291f
 transepiphyseal, 2285, 2286f
 treatment of, 2288–2289, 2289f
 treatment of, 2287–2293, 2289f–2293f
 heterotopic ossification of, in fibrodysplasia
 ossificans progressiva, 1634, 1636f
 in cerebral palsy, 1173–1203, 1178f–1209f,
 1180p–1207p. *See also* Cerebral palsy,
 hip in.
 in myelomeningocele, 1272–1277,
 1274f–1277f. *See also*
 Myelomeningocele, hip in.
 open reduction of, 546–568, 550p–567p,
 570p–571p
 anterolateral approach to, 548–549,
 560p–567p
 capsuloplasty in, 566p–567p
 capsulotomy in, 562p–565p
 casting after, 548–549, 566p
 dissection in, 560p–565p
 incision for, 548, 560p–561p
 incision in, 548, 560p–561p
 postoperative care in, 566p
 technical difficulties in, 548
 technique of, 564p–565p
 failure of, and avascular necrosis, 586–587,
 618f
 indications for, 546
 Lloyd Roberts intertrochanteric oblique
 osteotomy in, of proximal femur, and
 internal fixation with Coventry
 apparatus, 578p–581p
 medial approach to, 546–548, 550p–559p
 capsulotomy in, 556p–557p
 casting after, 558p–559p
 dissection in, 550p–555p
 femoral head reduction in,
 558p–559p
 incision for, 550p–551p
 with derotation osteotomy, 570p–571p
 with femoral shortening, 549, 549f,
 570p–581p
 in child 2 years and older, 549, 549f,
 568–569, 568f–569f, 570p–571p.
 See also Femur, shortening of.
 with innominate osteotomy, 549,
 568
 with intertrochanteric varus osteotomy,
 and fixation with blade plate,
 572p–577p
 dissection in, 572p–575p
 radiography during, 572p–573p

Hip (*Continued*)
 osteoarthritis of, femoral anteversion and,
 784
 painful
 examination of, 72, 72f
 from slipped capital femoral epiphysis,
 718–719
 pseudoadduction of, 1178
 pull-off power of, in cerebral palsy, 1129,
 1129f
 range of motion in, from slipped capital
 femoral epiphysis, 718, 718f
 range of motion of, 36–42, 38f–43f, 38t
 age and, 37–38, 38t
 redislocation of, after treatment of
 developmental dysplasia of hip,
 586–587, 618f
 rotation of, 40–41, 42f
 in femoral anteversion, 783, 786f
 screening examination of, 63–64, 64f
 sepsis of, posture in, 1845
 septic arthritis of, surgical drainage of, 1853,
 1854p–1857p, 1858
 stabilization of, in proximal focal femoral
 deficiency, 1763, 1770, 1773f–1775f
 subluxation of
 from femoral lengthening, 1089f
 in septic arthritis, 1848f
 subluxation/dislocation of
 in cerebral palsy. *See also* Cerebral palsy,
 hip in, subluxation/dislocation of.
 in myelomeningocele, 1273–1277,
 1274f–1277f. *See also*
 Myelomeningocele, hip in,
 subluxation/dislocation of.
 in spinal muscular atrophy, 1314,
 1314f–1315f
 synovitis of, acute transient, 1823–1825
 total joint arthroplasty of
 for cerebral palsy, 1587
 for osteopetrosis, 1553
 or rheumatoid arthritis, 1821
 Trendelenburg's test of, 41–42, 42f–43f
 Hip arthrodesis
 for avascular necrosis, in slipped capital
 femoral epiphysis, 753, 756p–757p
 for nonreconstructable hip in cerebral palsy,
 1201
 Hip fusion, for avascular necrosis, in slipped
 capital femoral epiphysis, 753, 755f
 Hip-knee-ankle-foot orthoses (HKAOs), for
 myelomeningocele, 1291–1292,
 1293–1294, 1294f
 Histiocytosis X, 104, 1938f, 1938–1942
 eosinophilic granuloma in, 1938f–1941f,
 1938–1942
 clinical features of, 1938–1939
 differential diagnosis of, 1940–1941
 pathology of, 1938, 1938f
 radiographic features of, 1939–1940,
 1939f–1941f
 treatment of, 1941–1942
 with Hand-Schüller-Christian disease, 1942
 with Letterer-Siwe disease, 1942
 History. *See* Orthopaedic history.
 "Hitchhiker's thumb," in diastrophic dysplasia,
 449, 1533, 1534f
 HKAO(s) (hip-knee-ankle-foot orthoses), for
 myelomeningocele, 1291–1292
 Hoffmann's reflex, 56
 Hogshead fascia lata transfer, to lumbodorsal
 fascia, for gluteus maximus paralysis, 1349,
 1349f
 Holt-Oram syndrome, 1752
 anesthesia in, 122t
 Home traction, for developmental dysplasia of
 hip, 540, 540f
 Homocystinuria, 1596–1598, 1597f, 1598t
 clinical features of, 1596–1597
 diagnosis of, 1597, 1598t
 pathophysiology of, 1596, 1597f
 treatment of, 1597–1598
 vs. Beals' syndrome, 1595
 vs. Marfan's syndrome, 1588
 Hook-rod system(s), for Scheuermann's
 kyphosis, 334–335, 335f
 Hormonal disorder(s), and macrodactyly, 452
 Horn cell(s), anterior, of spinal cord, in
 poliomyelitis, 1322
 Hounsfield unit(s), in computed tomography,
 138, 139f, 142t
 Hovnanian operation, 1419, 1426f–1427f
 Hughston and Norwood posterior drawer test,
 2363
 Hughston technique, for radiography of
 patellar dislocation, 796f
 Human bite wound(s), 478
 Human papillomavirus infection(s), of hand,
 479
 Humeral line, 2145, 2147f
 Humeroulnar joint, dislocation of, in Larsen's
 syndrome, 1602, 1604f
 Humerus
 Ewing's sarcoma of, 2034f
 lateral condylar fractures of, 2143f,
 2175–2185, 2178f–2188f
 anatomic considerations in, 2175, 2177
 classification of, 2177–2178, 2179f–2180f
 complications of, 2180, 2183–2185,
 2185f–2188f
 avascular necrosis as, 2185
 bone spurs as, 2183, 2185f
 cubitus valgus as, 2185, 2188f
 cubitus varus as, 2183, 2185f
 delayed union/nonunion as, 2183–2185,
 2186f–2188f
 fishtail deformity as, 2185, 2188f
 growth arrest as, 2185
 diagnosis of, 2177
 late displacement of, 2180, 2184f
 mechanism of, 2177
 minimally displaced, 2178, 2178f, 2180,
 2184f
 radiographic features of, 2177, 2178f
 treatment of, 2178, 2180, 2181f–2184f
 lateral epicondylar fractures of, 2206
 lateral rotation osteotomy of, for brachial
 plexus palsy, 486, 492p–495p
 medial condylar fractures of, 2202, 2205,
 2205f
 medial epicondylar fractures of, 2185–2191,
 2189f–2190f
 anatomic considerations in, 2185
 classification of, 2188
 complications of, 2190–2191
 diagnosis of, 2185
 entrapment in, 2185, 2189f, 2191
 mechanism of, 2185
 radiographic features of, 2187–2188, 2189f
 treatment of, 2188, 2190, 2190f
 medial rotation osteotomy of, for brachial
 plexus palsy, 486, 497f
 muscular insertions of, 2132f
 osteosarcoma of, 1956f, 1960f
 proximal, epiphysiolytic, 2128, 2128f

- Humerus (*Continued*)
- proximal metaphyseal/shaft fractures of, 2132–2134, 2132f–2134f
 - anatomic considerations in, 2132f, 2132–2133
 - complications of, 2134
 - diagnosis of, 2133
 - mechanism of, 2133, 2133f
 - treatment of, 2133–2134, 2134f
 - proximal phyeal fractures of, 2123–2128, 2124f–2128f
 - anatomic considerations in, 2123–2124, 2125f
 - associated conditions in, 2128, 2128f
 - classification of, 2125–2126
 - complications of, 2127–2128
 - diagnosis of, 2125, 2125f–2126f
 - mechanism of, 2124–2125
 - treatment of, 2126–2127, 2127f
 - sectioning of, in amputation through arm, 2020p–2021p
 - shortening of, after proximal phyeal fractures, 2127–2128
 - supracondylar fractures of, 2139–2168, 2140f–2175f
 - anatomic considerations in, 2139, 2140f–2142f
 - classification of, 2145–2146, 2147f–2149f
 - complications of, 2163–2168, 2166f–2175f
 - cubitus valgus/varus as, 2165–2168, 2166f–2174f
 - myositis ossificans as, 2168, 2175f
 - peripheral nerve injury as, 2165
 - vascular injury as, 2163–2165
 - Volkmann's ischemic contracture as, 2165
 - diagnosis of, 2141–2142, 2142f–2144f
 - differential diagnosis of, 2141, 2143f
 - extension type, 2140, 2140f
 - flexion type, 2140, 2141f
 - malreduced, 2162–2163, 2164f
 - mechanism of, 2140, 2140f–2141f
 - radiographic features of, 2142–2145, 2145f–2147f
 - treatment of, 2147–2160, 2150f–2159f
 - casting in, 2156
 - closed reduction in, 2149, 2151f–2153f, 2153
 - controversies in, 2160–2163, 2161f–2164f
 - displaced, 2149–2160, 2151f–2159f
 - emergent, 2147, 2149, 2150f
 - minimally displaced, 2160, 2161f
 - nondisplaced, 2149, 2150f
 - open reduction in, 2156, 2159f, 2159–2160
 - percutaneous pinning in, 2153–2156, 2154f–2158f
 - and ulnar nerve injury, 2153, 2155, 2155f, 2160–2162, 2162f
 - timing of, 2160
 - traction in, 2156, 2159f
 - with "pulseless pink hand," 2163f
 - T-condylar fractures of, 2202, 2204f
 - transphyseal fractures of, 2143f, 2168, 2175, 2176f–2177f
 - Humoral immunity, and dermatomyositis/polymyositis, 1493
 - Hunka classification, of complications after hip sepsis, 1859, 1860f
 - Hunter's syndrome, 1573t, 1576–1577
 - Hurler's syndrome, 1573t, 1574–1575, 1574f–1577f
 - anesthesia in, 122t
 - clinical features of, 1574–1575, 1574f–1575f
 - genetic features of, 1574
 - orthopaedic management of, 1575, 1577f
 - pathology of, 1574
 - radiographic features of, 1575, 1576f
 - treatment of, 1575
 - Hyaline cartilage, erosion of, in tuberculous arthritis, 1830, 1830f
 - Hydrocephalus, in myelomeningocele, 1253, 1257
 - Hydromyelia, in myelomeningocele, 1258
 - Hyperactivity, and Legg-Calvé-Perthes disease, 663–664, 671–672
 - Hypercalcemia, in hyperparathyroidism, 1702
 - Hyperextension, definition of, 27t
 - Hyperkalemia, and periodic paralysis, 1490–1491
 - Hyperlordosis
 - after rhizotomy, 309, 1232, 1232f
 - in myelomeningocele, 1289
 - in spondylolisthesis, 354, 355f - Hyperostosis
 - cortical, infantile. *See Caffey's disease.*
 - with macrodactyly, 451, 452f - Hyperparathyroidism, primary, 1701–1702
 - radiography in, 129, 129f
 - Hyperphosphatasia, 1711
 - Hyperplastic callus, in osteogenesis imperfecta, 1726–1727, 1728f
 - Hyperthermia, malignant, 118
 - Hypertrophic interstitial neuritis, 1445, 1447
 - Hypervitaminosis A, 1707, 1708f
 - vs. Caffey's disease, 1566t
 - Hypervitaminosis D, 1704–1705, 1705f
 - Hypochondroplasia, 1521
 - Hypokalemia, and periodic paralysis, 1490
 - Hypoparathyroidism, idiopathic, 1702–1703
 - Hypophosphatasia, 1708–1711, 1709f–1710f
 - clinical features of, 1709f–1710f, 1710
 - congenital, vs. osteogenesis imperfecta, 1727
 - differential diagnosis of, 1710–1711
 - genetic basis of, 1708
 - laboratory studies in, 1709–1710
 - pathology of, 1708–1709
 - prenatal diagnosis of, 1710
 - treatment of, 1711
 - Hypophosphatemic rickets
 - familial. *See Rickets, vitamin D-resistant.*
 - tumor-related, 1695 - Hypopituitarism, and dwarfism, 1711–1712
 - Hypotension, induced, for minimizing intraoperative blood loss, 116
 - Hypothyroidism, 1712–1713, 1713f–1714f
 - and slipped capital femoral epiphysis, 715–716
 - vs. Legg-Calvé-Perthes disease, 689, 693f
 - Hypotonia
 - congenital, benign, 1313t
 - in cerebral palsy, 1123 - Hypotonic child syndrome. *See also Spinal muscular atrophy (SMA).*
 - differential diagnosis of, 1313t - Hypotonic diplegia, cerebral, 1313t
 - Hypoxia, perinatal, and cerebral palsy, 1123
 - Hysterical scoliosis, 312
 - Idiopathic juvenile osteoporosis, 1715–1717, 1716f, 1716t
 - back pain in, 102
 - vs. osteogenesis imperfecta, 1728
 - Ilfeld splint, for developmental dysplasia of hip, in young child ages 1–6 months, 538, 538f
 - Iliac apophysis, dissection of, in open reduction of hip, using anterolateral approach, 562p–563p
 - Iliac horn(s), in nail-patella syndrome, 1599f, 1599–1600
 - Iliac spine, avulsion injury of, 2260f
 - Iliac wing(s)
 - fractures of, 2261, 2261f
 - in achondroplasia, 1512, 1512f - Iliacus muscle, dissection of, in iliopsoas muscle transfer, 1338p–1341p
 - Iliofemoral fusion, for hip stabilization, in proximal focal femoral deficiency, 1770, 1773f–1774f
 - Iliofemoral ligament, anatomy of, 2273, 2273f
 - Iliogluteal incision, in Banks and Coleman hemipelvectomy, 1968p–1969p
 - Ilioinguinal incision, in Banks and Coleman hemipelvectomy, 1964p–1965p
 - Iliopsoas muscle
 - contraction of, during gait, 79
 - dissection of, in Salter innominate osteotomy, 636p–637p - Iliopsoas muscle transfer, for gluteus medius paralysis, in poliomyelitis, 1336, 1338p–1347p
 - Iliopsoas tendon
 - as barrier to closed hip reduction, 520–521, 522f
 - dissection of, in open reduction of hip
 - anterolateral approach to, 562p–563p
 - medial approach to, 552p–555p
 - release of, for hip flexion contractures, 1173 - Iliopsoas tendon transfer, for hip subluxation/dislocation, in myelomeningocele, 1273–1274, 1274f–1275f
 - Iliotibial band
 - anatomy of, 1330, 1330f
 - contractures of, in poliomyelitis, 1329–2831, 1330f, 1332p–1335p, 1336, 1337f
 - dissection of
 - in excision of discoid lateral meniscus, 816p–817p
 - in quadricepsplasty, 806p–809p
 - release of, 787 - Ilium
 - exposure of, in Pemberton osteotomy, 622p–627p
 - osteotomy of, in Banks and Coleman hemipelvectomy, 1970p–1971p - Ilizarov apparatus
 - for angular deformities of leg, 1116f, 1117
 - for genu varum, in vitamin D-resistant rickets, 1693f
 - for limb lengthening, 1092–1093
 - in fibular hemimelia, 1781f
 - of residual tibia, 1796, 1797f - Ilizarov procedure, 878, 879f–880f, 881
 - for genu varum, in achondroplasia, 1517, 1518f
 - for talipes equinovarus, with residual deformity, 951, 953, 954f–957f
 - Imaging. *See under specific modality, e.g., Radiography.*
 - Imbrication, medial, of talonavicularcuneiform complex, for pes planovalgus, 916, 918f, 920f

- Imhauser osteotomy, for slipped capital femoral epiphysis, 736, 737f, 738
- Imiglucerase, for Gaucher's disease, 1645
- Immobilization, during radiography, 135
- Immune system, activation of, and chondrolysis, 749
- Immunization, against tetanus, 2073
- Immunosuppressive therapy, for dermatomyositis/polymyositis, 1495
- Incisional biopsy, of musculoskeletal tumors, 1897t, 1897–1898
- Inclinometer test, of thoracolumbar spine flexion, 36, 37f
- Incontinence, urinary, from myelomeningocele, 1258–1259
- Index finger, duplication of, 448
- Infant(s)
- development of locomotion in, 20
 - physical examination of, 72
- Infantile gigantism, anesthesia in, 120t
- Infantile muscular atrophy, 1313t
- Infection(s)
- back pain in, 102–104, 103f
 - limping from, 86–88, 87f–89f
 - of bone, 1841–1871. *See also* Osteomyelitis.
- Infectious spondylitis, and torticollis, 193–194
- Inflammation, cellular mediators of, and acute hematogenous osteomyelitis, 1842–1843
- Inflammatory disorder(s)
- back pain from, 102–104, 103f
 - limping from, 86–88, 87f–89f
- Inflammatory popliteal cyst(s), 830f, 831
- Infraspinatus muscle
- and shoulder motion, 1395
 - division of, for brachial plexus palsy, with shoulder deformities, 118f, 486
- INH (isoniazid), for tuberculous arthritis, 1831
- Injury Severity Score (ISS), 2073
- Inlet view, of pelvic fractures, 2257, 2257f
- Innominate osteotomy
- for Legg-Calvé-Perthes disease, 698–699, 701f–702f
 - for open reduction of hip, 549, 568
 - Salter, for avascular necrosis, 596, 618–621, 621f–633f, 631, 634p–641p. *See also* Salter innominate osteotomy.
- Instrumentation. *See under specific type, e.g., Harrington rod(s).*
- Intercalary limb deficiency(ies), 1747, 1747f
- Intergroup Rhabdomyosarcoma Study, 2040–2041, 2041t
- International Federation of Societies for Surgery of the Hand, classification of congenital anomalies, 410, 410t
- Interosseous membrane, anterior transfer of posterior tibial tendon through, in Duchenne muscular dystrophy, 1470, 1472p–1475p
- Interosseous nerve, palsy of, after Monteggia fractures, 2224–2225
- Interpedicular distance, in spine, achondroplasia and, 1513, 1513f
- Interposition arthroplasty, with proximal femoral resection, for nonreconstructable hip in cerebral palsy, 1203, 1209f
- Interposition material, for phyeal bar resection, 2071–2072, 2072f
- Intertrochanteric osteotomy
- double, for avascular necrosis, 612p–613p
 - in developmental dysplasia of hip, 584, 597f, 612p–613p
 - for avascular necrosis, after hip fracture, 1506, 1507f
- Intertrochanteric osteotomy (*Continued*)
- for femoral anteversion, 786
 - for slipped capital femoral epiphysis, 736, 737f, 738
 - oblique
 - Lloyd Roberts, of proximal femur, and internal fixation with Coventry apparatus, 578p–581p
 - of proximal femur, and internal fixation with Coventry apparatus, 1350, 1352p–1355p
 - repositioning, for residual deformity from slipped capital femoral epiphysis, 745, 746f
 - varus, and internal fixation with blade plate, for open hip reduction, 572p–577p
 - chisel placement in, 574p–575p
 - dissection in, 572p–575p
 - postoperative care in, 576p
 - radiography during, 572p–573p
- Intertrochanteric osteotomy, Pauwels', for developmental coxa vara, 773, 774f, 775, 776p–779p
- Intervertebral disk(s)
- calcification of, and torticollis, 194
 - herniation of, 98–99, 99f
 - diagnosis of, physical examination in, 96f in spondylolisthesis, 354, 356
 - lumbar, 369–374, 372f. *See also* Lumbar disk(s), herniation of.
 - removal of. *See* Discectomy.
- Intoeing
- assessment of, 65f–66f, 65–67. *See also* Femur, anteversion of.
 - from cerebral palsy, 1128f
 - from tibial torsion, 866
- Intorsion deformity, in talipes equinovarus, 924, 926f
- Intracapsular ring, anatomy of, and Legg-Calvé-Perthes disease, 659–660
- Intracranial pressure, increased, anesthesia and, 109
- Intramedullary canal, in osteopetrosis, 1551, 1551f
- Intramedullary rod(s)
- for congenital tibial pseudarthrosis, 872f–877f, 873–877
 - for femoral lengthening, 1093–1094, 1094f
 - for femoral shaft fractures, 2313–2314, 2313f–2316f, 2317f–2318f, 2319
 - for femoral shortening, in leg length inequality, 1076, 1076f
 - for long bone deformities, in osteogenesis imperfecta, 1733–1738, 1734f–1737f
 - for radial/ulnar fractures, 2230, 2231f
 - for tibial diaphyseal fractures, 2379, 2380f
 - for tibial shortening, in leg length inequality, 1077, 1078f
- Intramuscular hemorrhage, in hemophilia, 1882f, 1882–1883
- Intraosseous abscess(es), 1848, 1851f
- Intraspinal canal, deformity of, in scoliosis, 220, 220f
- Intravenous alarm-control (IVAC) pump, for intercompartmental pressure measurement, 2076
- Intravenous immune globulin (IVIG), for Guillain-Barré syndrome, 1455
- Inversion
- definition of, 27t
 - muscles producing, 1361
- Irwin wedge osteotomy, for genu recurvatum, in poliomyelitis, 1359, 1360f
- Ischial tuberosity, avulsion injuries to, 2259, 2261
- Ischial-bearing above-knee amputation, for osteosarcoma, 1982p–1991p
- Ischiopubic synchondrosis, 130f, 131
- Ischium, transfer of hip adductors to, 1184, 1185f
- ISO/ISPO international classification, of limb deficiencies, 1749–1750, 1750t–1751t
- Isola instrumentation, for idiopathic scoliosis, 243, 258, 258f
- Isoniazid (INH), for tuberculous arthritis, 1831
- Isotretinoin, for fibrodysplasia ossificans progressiva, 1638
- ISS (Injury Severity Score), 2073
- Isthmic spondylolisthesis, 352, 353f
- IVAC (intravenous alarm-control) pump, for intercompartmental pressure measurement, 2076
- IVIG (intravenous immune globulin), for Guillain-Barré syndrome, 1455
- J sign
- in patellar dislocation, 794
 - in recurrent patellar subluxation, 799
- Jansen metaphyseal chondrodysplasia, 1547
- Japas V tarsal osteotomy, for pes cavus, 999, 1006p–1009p
- Jefferson fracture(s), 2092
- Jeune's disease, 1682f, 1683
- Joe Cool splint, 393, 395f
- Joint(s)
- aspiration of, 1811, 1812f
 - in hemophilia, 1886
 - in septic arthritis, 87
 - hypermobility of, in Ehlers-Danlos syndrome, 1639, 1640f–1641f
 - infection of, 1841–1859
 - inflammation of, pathophysiology of, 1811
 - laxity of, in Marfan's syndrome, 1589, 1589f, 1593
 - motion of, types of, 27t
 - range of motion of. *See* Range of motion.
 - subluxation of, limb lengthening and, 1087–1088, 1089f, 1090
 - types of, 1811
- Jones classification, of tibial deficiency, 1779, 1782f–1784f
- Jones fracture(s), 2431, 2432f–2433f
- Jones transfer, 991, 994, 995f, 996p–997p
- for Charcot-Marie-Tooth disease, 1441f
- Juvenile idiopathic osteoporosis, 1715–1717, 1716f, 1717t
- Juvenile rheumatoid arthritis. *See* Rheumatoid arthritis.
- KAFO(s) (knee-ankle-foot orthoses). *See* Knee-ankle-foot orthoses (KAFOs).
- Kalamchi-MacEwen classification system, for avascular necrosis, 583, 587t
- Kaneda instrumentation, for thoracolumbar curves, in idiopathic scoliosis, 261, 261f
- Keith needle, insertion of, in greater trochanteric epiphysiodesis, for avascular necrosis, 590p–591p
- Keratan sulfate, urinary, in Morquio syndrome, 1577
- Ketorolac, after scoliosis surgery, 241
- Kidner procedure, for accessory navicular, 896

- Kinematics, 77–79, 78f–79f
 in coronal plane, 77–78, 79f
 in sagittal plane, 77, 78f–79f
 in transverse plane, 78–79, 79f
- Kinetic muscle strength, 44
- Kinetics, 80, 81f
- King procedure, for foot amputation with knee arthrodesis, for proximal focal femoral deficiency, 1763, 1764p–1767p
- Kingella kingae* infection(s), and osteomyelitis, 1845
- King-Moe classification, of curve patterns in idiopathic scoliosis, 234, 234f, 234t
 and fusion level, 235–236, 236f–238f
- Kirschner wire(s), placement of for arthrodesis of proximal interphalangeal joint, 1034p
 in correction of vertical talus, 970p–973p
 in metaphyseal distal forearm fractures, 2236–2237, 2237f
 in percutaneous pinning of supracondylar humeral fractures, 2153, 2155
 in trochanteric transfer, for avascular necrosis, 602p–603p, 606p–609p
- Klein's line, in slipped capital femoral epiphysis, 719, 719f
- Klinefelter's syndrome, anesthesia in, 122t
- Klippel-Feil syndrome, 173–187, 178f–187f
 anesthesia in, 122t
 anomalies occurring with, 175, 180f–181f
 clinical features of, 173–176, 178f–179f, 182f
 diagnosis of, 175–176
 genetic basis of, 174
 occipital atlantal assimilation in, 184f–185f, 185
 radiographic features of, 176–181, 178f, 182f–183f
 treatment of, 178f–179f, 181–187, 184f–187f
 fusion in, 178f–179f, 183, 185–187, 186f–187f
 halo in, 183
 with congenital scoliosis, cervicothoracic abnormalities in, 286f
- Klippel-Trénaunay syndrome, 1670–1672, 1671f
- Klippel-Trénaunay-Weber syndrome
 anesthesia in, 122t
 macrodactyly in, 1024, 1026f
 vs. Proteus syndrome, 1669
- Klusic test, 526, 526f
- Knee. *See also* Patella; *under* Genu.
 arthritis of, femoral anteverision and, 784
 arthrodesis of, with foot amputation, for proximal focal femoral deficiency, 1762f, 1762–1763, 1764p–1767p
 calcification of, in dermatomyositis/polymyositis, 1494, 1494f
 congenital dislocation of, 815, 822–825, 822f–825f
 classification of, 823, 825f
 clinical features of, 822–823, 823f–824f
 conditions associated with, 823
 diagnosis of, 823
 etiology of, 822, 822f
 pathology of, 823
 treatment of, 823, 825
 contractures of, in myelomeningocele, 1269, 1272
 disarticulation of, 1794
 for osteosarcoma, 1992p–1995p
 for tibial deficiency, 1780
 prostheses for, 1804
- Knee (*Continued*)
 dislocation of
 in arthrogryposis, 1652–1653
 in Larsen's syndrome, 825, 827f, 1603f–1604f
 treatment of, 1607–1608, 1608f–1611f
 in myelomeningocele, 1269
 effusion in, radiography of, 1846, 1847f
 extension of
 after tibial spinal fracture repair, 2357
 and gait, 77, 78f
 talipes equinus and, 1356f
 flexion contractures of
 from hamstring spasticity, in cerebral palsy, 1159, 1167f
 in arthrogryposis, 1653
 in Beals' syndrome, 1594, 1595f
 in poliomyelitis, 1358, 1358f
 in rheumatoid arthritis, 1820–1821
 in tibial deficiency, 1779, 1784f
 flexion of, and gait, 77, 78f
 “floating,” 2389f, 2389–2390
 goniometry of, 26, 26f
 in cerebral palsy, 1158–1159, 1166f–1168f, 1166–1173, 1170p–1171p, 1172f, 1174p–1177p, 1178f. *See also* Cerebral palsy, knee in.
 in poliomyelitis, 1351–1361, 1356f–1361f. *See also* Poliomyelitis, knee in.
 injuries to, 2321–2367, 2322f–2367f. *See also* Femur, distal epiphyseal fractures of; Patella, fractures of.
 ligamentous, 2361f–2365f, 2361–2366. *See also* Anterior cruciate ligament (ACL); Posterior cruciate ligament (PCL).
 instability of, in myelomeningocele, 1272
 osteochondritis dissecans of. *See* Osteochondritis dissecans.
 osteosarcoma of, 147f
 pain in, from slipped capital femoral epiphysis, 718
 range of motion of, 43, 43f
 snapping, congenital, 825, 826f
 stabilization of, after hip arthrodesis, 1350, 1356f
 subluxation of, from callotasis, 1089f
 swelling of, in rheumatoid arthritis, 1817, 1818f
 synovectomy of, for hemophilic arthropathy, 1888–1889
 total joint arthroplasty of, in rheumatoid arthritis, 1821
 valgus deformity of, in poliomyelitis, 1330
 vascular supply to, 2322
- Knee-ankle-foot orthoses (KAFOs)
 for Duchenne muscular dystrophy, 1470, 1476f
 for infantile tibia vara, 845–847, 847f
 for leg length inequality, 1058, 1061f
 for myelomeningocele, 1292–1293, 1293f–1294f
- Kniest's dysplasia, 1542–1545, 1543f
 clinical features of, 1542
 genetic features of, 1542
 orthopaedic management of, 1544
 pathology of, 1542
 radiographic features of, 1542, 1543f
 treatment of, 1544
- Knock-knee. *See* Genu valgum.
- Kocher-Langenbeck approach, to acetabular fractures, 2270, 2271f
- Köhler's disease, 899–900, 899f–900f
- Kramer and Barmada osteotomy, for slipped capital femoral epiphysis, 737f, 737–738, 742p–743p
- Kugelberg-Welander spinal muscular atrophy, 1312, 1312f, 1312t
- Kyphoscoliosis
 in diastrophic dysplasia, progression of, 1540f, 1541
 in Kniest's dysplasia, 1543f, 1544
 in metatropic dwarfism, 1677, 1677f
 in osteogenesis imperfecta, 1724, 1724f
- Kyphosis, 323–349
 after laminectomy, 308, 310f–311f, 337, 339, 340f–342f
 and neurologic disorders, 302
 cervical, 195f–205f, 195–206
 clinical features of, 195–198, 195f–198f
 congenital, 195
 in diastrophic dysplasia, 1538, 1538f–1539f, 1541
 postlaminectomy, 202, 203f, 204
 treatment of, 204–206, 205f
 progression of, 203f
 radiographic features of, 199
 treatment of, 199, 200f–203f, 202
 congenital, 291–297, 294f–295f, 323
 classification of, 291, 292f
 clinical features of, 294
 from failure of vertebral formation, 291, 293f
 from failure of vertebral segmentation, 291, 294f, 295–297
 magnetic resonance imaging in, 294
 natural history of, 291, 294
 radiographic evaluation of, 294, 296f
 treatment of, 294–297, 296f–298f
 failure of vertebral formation and, 294–295, 295f–298f
 failure of vertebral segmentation and, 295, 297
 mixed anomalies and, 297
 correction of, in anterior instrumentation, for idiopathic scoliosis, 270p–271p
 etiology of, 323, 325t
 from progressive noninfectious anterior fusion, 336–337, 339f
 in achondroplasia, 346
 in ankylosing spondylitis, 1822f
 in myelomeningocele, 1277–1280, 1278f–1279f
 and pressure sores, 1279, 1279f
 lumbar kyphectomy for, 1280, 1282p–1287p
 partial vertebrectomy for, 1279, 1279f
 patterns of, 1277, 1278f
 radiographic features of, 30f–31f
 in neurofibromatosis, 300–301, 302f
 in tuberculous spondylitis, 1834, 1835f
 lumbosacral, in high-grade spondylolisthesis, 363
- Marfan's syndrome and, 347–349, 348f
 measurement of, in scoliosis, radiographic, 226
- mucopolysaccharidosis and, 346–347, 347f
- physical examination in, 69, 70f
- postirradiation, 312, 339–346, 343f–346f
- pathophysiology of, 339, 343f–344f
- prevention of, 339
- treatment of
 prophylactic, 339–345
 strut grafts in, 343f, 345
- surgical, 345–346, 346f

- Kyphosis (*Continued*)
 postlaminectomy, 308, 310f–311f, 337, 339, 340f–342f
 postural, 323, 324f, 325t
 pseudoachondroplasia and, 346
 Scheuermann's, 101f, 101–102, 324–336, 326f–338f, 328t. *See also* Scheuermann's kyphosis.
 spondyloepiphyseal dysplasia and, 346–347
 thoracic, in neurofibromatosis, 145f
 thoracolumbar
 in achondroplasia, 1514–1515, 1514f–1515f
 in Hurler's syndrome, 1575, 1575f–1576f
 treatment of, rib strut graft for, 148f
 tuberculous, 1867f
- Lachman test, 2362–2363, 2363f
- Lag screw(s)
 for Lloyd Roberts intertrochanteric oblique osteotomy, of proximal femur, 578p–581p, 1352p–1355p
 placement of, in trochanteric transfer, for avascular necrosis, 606p–607p
- Lambrinudi operation, 1387
- Lamina(e), deformities of, in scoliosis, 220, 220f
- Laminar hook(s), for Texas Scottish Rite Hospital instrumentation, 246p–247p, 252p–255p
- Laminectomy
 for slipped vertebral apophysis, 374–375
 kyphosis after, 337, 339, 340f–342f
 cervical, 202, 203f, 204
 treatment of, 204–206, 205f
 osteoplastic, 339
 scoliosis after, 308–312, 310f–311f
 subaxial instability after, 207, 209f, 210
- Landau reflex, 54
- Langenskiöld classification, of infantile tibia vara, 844f, 844–845
 and type of treatment, 847–850, 848f–849f
- Langenskiöld greater trochanteric epiphysiodesis, for avascular necrosis, in developmental dysplasia of hip, 588p–595p. *See also* Trochanteric epiphysiodesis.
- Langerhans cell histiocytosis. *See* Histiocytosis X.
- Language, developmental milestones in, 4f
- Larsen-Johansson disease, 811, 812f
- Larsen's syndrome, 195f, 195–196, 1602–1614, 1603f–1614f
 anesthesia in, 122t
 clinical features of, 1602–1603, 1603f–1605f
 differential diagnosis of, 1604
 dislocation of elbow in, 1602, 1604f, 1611
 dislocation of knee in, 825, 827f, 1603f–1604f
 treatment of, 1608f–1611f
 neurologic assessment in, 1603–1604
 radiographic features of, 199, 1604, 1606f
 treatment of, 199, 200f, 1604–1614, 1607f–1614f
 for cervical kyphosis, 1604, 1607, 1607f
 for congenital knee dislocation, 1607–1608, 1608f–1611f
 for elbow/radial head dislocation, 1611
 for foot deformities, 1610–1611, 1612f
 for hip dislocation, 1608, 1608f–1611f, 1610
 for scoliosis, 1611–1614, 1613f–1614f
 priorities in, 1614
- Lasègue's sign, 98
- Lateral collateral ligament (LCL), injuries to
 diagnosis of, 2364
 treatment of, 2365–2366
- Lateral column lengthening, for pes valgus, in cerebral palsy, 1152–1153, 1160p–1163p, 1164f
- Lateral pillar B hip, in Legg-Calvé-Perthes disease, 681–682, 684f
- Lateral pillar classification, of Legg-Calvé-Perthes disease, 684–687, 687f–690f, 687t
 group A, 685, 687f–688f, 687t
 group B, 685, 687t, 689f
 group C, 685, 687t, 690f
- Latex, allergy to, 117–118
 in myelomeningocele, 1261
- Latissimus dorsi muscle
 dissection of, in forequarter amputation, 2002p–2003p, 2012p–2013p
 transfer to rotator cuff, for brachial plexus palsy, 108f, 110p–113p, 485
- Latissimus dorsi muscle transfer, in poliomyelitis, 1419, 1426f–1427f
- Latissimus dorsi tendon transfer, for subscapularis muscle paralysis, in poliomyelitis, 1407, 1409f
- Latissimus dorsi transfer, bipolar, for elbow flexion, in arthrogryposis, 1655, 1660f–1662f
- Lauge-Hansen classification, Dias-Tachdjian modification of, for ankle fractures, 2392–2393, 2393f
- Laurence-Moon-Biedl syndrome, anesthesia in, 122t
- Laurie technique, for radiography of patellar dislocation, 796, 797f
- Lawnmower(s), injuries from, 2075, 2432, 2434
- Lawrence-Gehan staging system, for rhabdomyosarcoma, 2040, 2040t
- LCL. *See* Lateral collateral ligament (LCL).
- Lead poisoning, metaphyseal bands in, 132f
- Leg
 angular deformities of, 1103–1117, 1104f–1116f, 1105t
 assessment of, 1103, 1105f
 correction of, 1107, 1110f–1112f, 1111
 epiphysiodesis in, 1113, 1114f
 external fixation in, 1114, 1116f
 hemiepiphyseal stapling in, 1111, 1112f–1113f
 osteotomy in, 1113–1117, 1115f–1116f
 etiology of, 1103, 1104f, 1105t
 from femoral shaft fractures, 2319
 in achondroplasia, 1517, 1518f
 in renal osteodystrophy, 1699
 physeal arrest and, 2069f
 compartments of, 2372, 2372f
 disorders of, 839–887. *See also* Genu; under Tibia.
 valgus deformity of, after intramedullary rod fixation, for congenital tibial pseudarthrosis, 876, 876f
- Leg length inequality, 1039–1103, 1041f–1094f and spinal deformity, 70f
 assessment of, 68, 69f, 1041–1045, 1042f–1046f
 clinical, 1041–1044, 1042f–1045f
 radiographic, 1043, 1043f–1046f
 etiology of, 1039–1040, 1040t, 1041f
 from femoral shaft fractures, 1040, 2319
 from intramedullary rod fixation, for congenital tibial pseudarthrosis, 876
 from open tibial fractures, 2386
- Leg length inequality (*Continued*)
 functional vs. structural, 1042–1043, 1042f–1043f
 gait in, 84, 1040–1041
 idiopathic hemihypertrophy and, 1039
 impact of, 1040–1041
 in Klippel-Trénaunay syndrome, 1672
 limping from, 92–83
 prediction of, 1045–1065, 1049t, 1050f–1055f, 1052t, 1056p–1060p
 normal skeletal growth and, 1045–1048, 1047f–1048f
 remaining growth in femur and tibia and, 1048–1053, 1049t, 1050f–1055f, 1052t, 1056p–1057p. *See also* Femur, prediction of remaining growth in.
 screening for, 1041–1042
 treatment of, 1058–1095, 1061f–1062f, 1064p–1071p, 1072f–1094f
 epiphyseal stapling in, 1062f, 1063–1095, 1064p–1071p, 1072
 epiphysiodesis in, 1082–1083
 indications for, 1058, 1061t
 knee-ankle-foot orthoses in, 1058, 1058f
 limb lengthening in, 1077–1097, 1079f–1094f
 acute, 1081–1082, 1081f–1082f
 by growth stimulation, 1077–1078
 complications of, 1088–1091, 1089f
 dynamic axial fixator for, 1092
 effect on articular cartilage, 1087–1088
 effect on muscle, 1087, 1087f
 effect on new bone formation, 1086f, 1086–1087
 effect on peripheral nerves, 1087
 gradual, 1082–1088, 1084f–1087f
 history of, 1078–1081, 1079f–1080f
 Ilizarov apparatus for, 1092–1093
 indications for, 1091
 internal and external fixation for, 1093–1094, 1094f
 psychosocial factors in, 1751
 results of, 1091–1092
 totally implantable devices for, 1094–1095
 limb shortening in
 acute, 1074–1077, 1075f–1078f
 epiphyseal stapling for, 1063, 1073–1074, 1073f–1074f
 epiphysiodesis for, 1058–1063, 1062f, 1064p–1071p, 1072f
 femoral shortening for, 1074–1077, 1075f–1076f
 tibial shortening for, 1077, 1077f–1078f
 orthoses in, 1058, 1061f
 Legg-Calvé-Perthes disease, 655–704
 arthrography in, 682, 685f
 bone scan in, 681–682, 683f–684f
 classification of, 683–689, 686f–692f, 687t
 by end results, 687, 691f
 Catterall, 683–684, 686f
 lateral pillar, 684–687, 687t, 687f–688f
 group A, 685, 687t, 687f–688f
 group B, 685, 687t, 689f
 group C, 685, 687t, 690f
 Mose, 687
 Stulberg, 687, 691f, 691t
 clinical course of, 673–674, 673t–674t
 clinical features of, 670–674, 671t, 672f–673f, 673t–674t
 age at onset and, 670
 during fragmentation stage, 673
 during healing stage, 673–674

Legg-Calvé-Perthes disease (*Continued*)
radiographic correlation with, 673,
674t
signs in, 672–673, 672f–673f
symptoms in, 670–672
computed tomography in, 683
definition of, 655
differential diagnosis of, 689–691, 692t,
693f–694f
etiology of, 656–665, 659t, 660f–665f
coagulopathy in, 656–657
environmental, 665
femoral head arterial circulation in, 658f,
658–661, 660f–662f
hereditary, 664
hyperactivity in, 663–664, 671–672
“predisposed child” theory in, 661–662,
664f
synovitis and, 665
trauma in, 662–663
unifying hypothesis in, 665, 665f
venous drainage of femoral head/neck in,
661, 663f
historical perspective on, 655–656,
656f–659f
containment treatment concepts and,
655–656, 659f
early treatment concepts and, 655,
658f–659f
incidence of, 655
limping from, 89–90, 91f
magnetic resonance imaging in, 679–681,
682f
natural history of, 674, 674t
pathology of, 665–670, 666f–671f
anatomic, 665–668, 666f–670f
articular cartilage thickening in, 668,
671f
double infarction in, 661, 662f, 669–670
in necrotic stage, 666, 666f
metaphyseal changes in, 670
new bone formation in, 667, 667f–670f
systemic findings in, 668–669, 671f
prognosis in
based on risk factors, 687–689, 692f, 692t
long-term, 702–704
radiography in, 675–679, 678f–682f
and acetabular changes, 679
and bilateral changes, 679
and limited changes, 679, 681f–682f
and metaphyseal changes, 675–678,
678f–679f
and physis changes, 678–679, 680f
radiography in, for staging, 675, 675t,
676f–677f
subchondral fractures in, 656, 659f, 699f
treatment of, 691–702, 693t, 694f–704f
approach to, 692, 695t
containment
experimental rationale for, 693–694
orthoses in, 694–695, 695–696,
695f–698f
surgical, 696–701, 699f–703f
cheilectomy in, 701
Chiari osteotomy in, 701, 703f
femoral osteotomy in, 696–698,
699f–700f
femoral/innominate osteotomy in,
699
innominate osteotomy in, 698–699,
701f–702f
shelf arthroplasty in, 700–701
valgus osteotomy in, 699–700

Legg-Calvé-Perthes disease (*Continued*)
history of, 655, 658f–659f
osteochondritic lesions in femoral head
after, 701, 704f
symptomatic, 692–694, 695f
bedrest in, 692
crutches in, 693
traction in, 692–693, 695f
ultrasonography in, 683
vs. acute transient synovitis of hip, 1824
vs. multiple epiphyseal dysplasia, 1534
Leg-straightening reflex, 51–52
Lens, dislocation of, in homocystinuria,
1597
Lesch-Nyhan syndrome, 1453–1454
anesthesia in, 122t
Letournel classification, of acetabular fractures,
2255
Letterer-Siwe disease, 1938, 1942
Leukemia
acute lymphocytic, back pain in, 104–105
limping from, 93
vs. osteomyelitis, 1851–1852
“Leukemia lines,” 128, 128f
Leukomalacia, periventricular, in cerebral palsy,
1124
Levator scapulae muscle, dissection of, in
forequarter amputation, 2004p–2005p
Levator scapulae tendon transfer, for
supraspinatus muscle paralysis, 1407,
1408f
Levoscoliosis, 145f
Lichtblau procedure, 1653
for recurrent talipes equinovarus, 948, 949f
Lichtblau's test, 903f
Lift(s), for leg length inequality, 1058, 1061f
Ligament of Wrisberg, anatomy of, 818p–819p
Ligamentous laxity, and developmental
dysplasia of hip, 514–515, 515f
Ligamentum teres, excision of, in open
reduction of hip
anterolateral approach to, 564p–565p
medial approach to, 556p–559p
Limb
acquired absence of, 1790–1796,
1790f–1796f. *See also* Amputation.
congenital absence of, 1751–1756,
1752f–1757f, 1754t–1755t
anomalies in, 1752, 1753f, 1754, 1754t
embryologic timing and, 1751–1752,
1752f
etiology of, 1752
psychosocial issues in, 1755–1756, 1757f
subclavian artery supply disruption
sequence hypothesis in, 1754–1755,
1754f–1756f, 1755t
development of, genetic regulation of, 1745,
1746f, 1747
embryology of, 1745, 1746f
lengthening of. *See also* Leg length inequality,
treatment of, limb lengthening in.
for hypochondroplasia, 1521
for proximal focal femoral deficiency,
1769–1771, 1773, 1774f–1776f
vs. amputation, for fibular hemimelia,
1776–1779
lower. *See* Lower extremity(ies).
phantom pain in, 1796
realignment of, in fracture management,
2078, 2078f
upper. *See* Upper extremity(ies).
Limb deficiency(ies)
chorionic villus sampling and, 1755

Limb deficiency(ies) (*Continued*)
classification of, 1747f–1749f, 1747–1750,
1750t–1751t
Frantz and O’Rahilly, 1747–1748,
1747f–1749f
ISO/ISPO, 1749–1750, 1750t–1751t
intercalary, 1747, 1747f
longitudinal, 1749–1750, 1750t
lower, 1756–1782, 1758f–1762f,
1764p–1767p, 1768f–1783f
femoral, 1756–1771, 1758f–1762f,
1764p–1767p, 1768f–1776f. *See also*
Femoral deficiency, proximal focal.
fibular, 1771–1778, 1776f–1779f, 1777t
of foot, 1782, 1783f
tibial, 1778–1782, 1780f–1782f
multiple, 1796–1799, 1797f–1798f
bilateral lower, 1798–1799
bilateral upper, 1796–1798, 1797f–1798f
bilateral upper and lower, 1799
bilateral upper/one lower, 1799
one upper/one lower, 1799
quadrimelic, 1799
terminal, 1747, 1747f–1748f
terminology related to, 1748–1749, 1749t
transverse, 1749–1750, 1750t
treatment of
long-term planning in, 1751
social factors in, 1751
timing of, 1750–1751
upper, 1782–1790, 1784f–1786f
transverse, 1782–1787, 1784f–1786f
amnion disruption sequence and, 1784,
1785f–1786f, 1786–1787
classification of, 1783, 1784f
etiology of, 1782–1783
radial, 1787, 1788f
treatment of, 1783–1784, 1784f–1785f
ulnar, 1787–1790, 1789f–1790f. *See also*
Ulnar deficiency.
Limb length inequality. *See* Leg length
inequality.
Limb salvage
contraindications to, 1792
for rotationplasty, 2028–2031, 2030f–2032f
in osteosarcoma, 1963
Limb-girdle muscular dystrophy, 1479–1481
clinical features of, 1464t
Limbus, anatomy of, 519, 519f–520f
Limp. *See also* Gait.
abductor
after varus femoral osteotomy, 697, 700f
in avascular necrosis, 583
differential diagnosis of, 84t
in adolescent, 85–86
in school-age child, 85
in toddlers, 85, 86f
evaluation of
imaging studies in, 83
physical examination in, 83
from cerebral palsy, 88
from chondrolysis, 91, 91f
from coxa vara, 89, 90f
from developmental dysplasia of hip, 89, 90f
from discoid meniscus, 91
from inflammatory/infectious disorders,
86–88, 87f–89f
from leg length inequality, 92–93
from Legg-Calvé-Perthes disease, 89–90, 91f,
670
from neoplastic disease, 93, 93f
from neurologic disorders, 88–89
from osteochondritis dissecans, 91

- Limp (*Continued*)

from overuse syndrome, 92

from slipped capital femoral epiphysis, 90–91, 91f

from tarsal coalition, 92, 92f
- Lipoma, of upper extremities, 502
- Lipomeningocele, 1302–1303, 1302f–1304f
- Lisch nodule(s), in neurofibromatosis, 1625
- Lisfranc's fracture, 2425–2428, 2426f–2429f

anatomic considerations in, 2426

classification of, 2426–2427, 2427f

complications of, 2428

diagnosis of, 2427, 2428f

mechanism of, 2426, 2426f

treatment of, 2427–2428, 2429f
- "Little League elbow," 2208, 2209f
- "Little League shoulder," 2128, 2128f
- Littlewood technique, of forequarter amputation, 2000p–2013p
- Lloyd Roberts intertrochanteric oblique osteotomy, of proximal femur, and internal fixation with Coventry apparatus, 578p–581p, 1352p–1355p
- LMB extension/flexion splint(s), 393, 394f
- LMB Finger-Hugger, 393, 393f
- Local anesthesia

intravenous, 115–116

side effects of, 115–116
- Locomotion, developmental milestones for, 20
- Long bone(s)

deformities of, in osteogenesis imperfecta, 1730–1731, 1731f. *See also* Osteogenesis imperfecta, long bone deformity in.

dumbbell shape of, in metatropic dwarfism, 1673, 1674f

fractures of, in osteogenesis imperfecta, 1730–1731, 1731f

growth of, longitudinal, 1045–1048, 1048f

of upper extremities, growth of, physis and, 10f

osteoblastoma of, 1936

"popcorn" calcification of, in osteogenesis imperfecta, 1725, 1726f
- Longitudinal arch, reconstitution of, in pes planovalgus, 910, 911f, 912
- Longitudinal epiphyseal bracket, 1027, 1032, 1033f
- Loose body(ies), in osteochondritis dissecans, 792, 792f
- Looser classification, of osteogenesis imperfecta, 1721
- Lorber's criteria, 1253
- Lordosis

lumbar

excessive, in developmental dysplasia of hip, 89, 90f

in poliomyelitis, 1331

measurement of, in scoliosis, radiographic, 226

thoracic, with lumbar kyphosis, in myelomeningocele, 1279f
- Low back pain, leg length inequality and, 1041
- Lower extremity(ies)

alignment of, normal, 1103–1107, 1106f–1109f

anatomic axis of, vs. mechanical axis, 2322f

angular deformities of. *See* Leg, angular deformities of.

contractures of, in metatropic dwarfism, 1673, 1674f
- Lower extremity(ies) (*Continued*)

epiphyses of, secondary ossification centers in, age at appearance of, 8f

epiphysis of, closure of growth plates in, age at, 9f

external rotation of, with talipes calcaneovalgus, 905

injuries to, 2251–2434. *See also under specific site, e.g., Pelvis, fractures of.*

lengthening of, for achondroplasia, 1515, 1517f

muscles of

innervation of, 48t

manual tests of, 59–61

prostheses for, 1801

gait problems with, 1801–1802

Lumbar curve(s), in idiopathic scoliosis

anterior instrumentation for, 262p–275p

diaphragm dissection in, 264p–271p

incision in, 262p–263p

positioning in, 262p–263p

protection of aorta and vena cava in, 268–269

solid rod, 260–261, 260f–261f

spinal fusion for, 236, 238f

Lumbar disk(s), herniation of, 369–374, 372f

clinical features of, 371

etiology of, 371

incidence of, 371

radiographic features of, 371–372, 372f

treatment of, 373–374
- Lumbar hyperlordosis, after rhizotomy, 309
- Lumbar kyphectomy, in myelomeningocele, 1280, 1282p–1287p
- Lumbar kyphosis

in myelomeningocele, 1278f–1279f, 1279

Scheuermann's, 102, 102f
- Lumbar lordosis

excessive, in developmental dysplasia of hip, 89, 90f

in poliomyelitis, 1331

measurement of, in scoliosis, radiographic, 226
- Lumbar spine. *See also* Thoracolumbar spine.

anatomy of, 2101, 2101f

congenital anomalies of, and disk herniation, 371

hyperlordosis of

after rhizotomy, 1232, 1232f

in pseudoachondroplasia, 1525f

in spondyloepiphyseal dysplasia congenita, 1527f

injuries to, 2101–2110, 2101f–2110f, 2106t

classification of, 2103, 2106t, 2107f

complications of, 2107–2108, 2109f

diagnosis of, 2102f, 2102–2103, 2104f

differential diagnosis of, 2103, 2105f

mechanism of, 2102

treatment of, 2103, 2105–2107, 2108f–2109f

without radiographic abnormality, 2110, 2110f
- Lumbar vertebra(e), spondylomegaly of, in Proteus syndrome, 1668f
- Lumbopelvic junction, in caudal regression syndrome, 1307, 1307f–1309f
- Lumbosacral agenesis. *See* Caudal regression syndrome.
- Lumbosacral junction, bony deformities of, computed tomography in, 146f
- Lumbosacral kyphosis, in high-grade spondylolisthesis, 363
- Luminescent screen-film combination(s), for radiography, 134
- Lung(s). *See also under Pulmonary.*

metastatic disease in, from osteosarcoma, 1957, 1960f
- Luque instrumentation

double L-rod segmented, for idiopathic scoliosis, 242, 242f, 244p–245p

for Scheuermann's kyphosis, 334

for spinal fusion

in cerebral palsy, 1226–1227

in myelomeningocele, 1286p–1287p, 1288f

in osteogenesis imperfecta, 1738–1739, 1739f

in spinal muscular atrophy, 1318f
- Luque-Galveston instrumentation, for spinal fusion, in spinal muscular atrophy, 1317f
- Lyme disease, 1868–1869
- Lytic lesion(s), in Gaucher's disease, 1645
- Macrodactyly, 450–461, 451f–460f, 1024, 1025f–1026f

classification of, 450–451, 451f–453f

clinical features of, 454–456, 454f–456f

differential diagnosis of, 450

etiology of, 451–452

historical perspective on, 450

imaging studies of, 456

in Proteus syndrome, 1666, 1668f

in von Recklinghausen's disease, 455f

of thumb, 458, 460f

pathology of, 452–454

site distribution of, 454, 454f

static subtype of, 454

treatment of, 456–459, 457f–460f

amputation in, 458, 460f

complications of, 459

digital nerve resection for, 457–458, 459f

epiphyseal arrest in, 457

goals of, 457

limitations of, 457

methods of, 457

nerve-stripping in, 457, 457f

palmar dog-ear from, 457, 458f

with hemihypertrophy of extremities, 451–452, 453f, 458–459

with neurofibromatosis. *See* Von Recklinghausen's disease.
- Madelung's deformity, 416–421, 417f–420f

clinical features of, 417f, 417–418

complications of, 421

etiology of, 416

historical perspective on, 416

imaging studies of, 418, 418f–419f

pathology of, 416–417, 417f

reverse, 418

treatment of, 418–421, 420f
- Maffucci's syndrome, 1927–1929

anesthesia in, 122t

vs. Proteus syndrome, 1669
- Magerl technique, for atlantoaxial arthrodesis, 187f
- Magnetic resonance imaging (MRI), 150f–152f, 150–162, 152t–153t, 154f–161f

advantages of, 150

artifacts on, 155

contrast in, 152, 152t

diagnostic errors from, 159, 162

gadolinium for, 156

history of, 150f, 150–151

image appearance in, 152f, 152–153, 152t–153t

- Magnetic resonance imaging (MRI)
(Continued)
- image enhancement in, 156
 - image interpretation in, 153–155
 - in differential diagnosis of back pain, 98
 - indications for, 153t, 154f–161f, 158–163
 - patient preparation for, 156–158
 - physical principles of, 151–152, 151f–152f
 - safety precautions in, 156–158
 - short-tau inversion recovery, 153
 - signal intensity in, 153t, 153–155, 154f, 156f
- Malformation(s). *See also under specific type, e.g., Scoliosis.*
- and growth and development, 5
 - Malgaigne pelvic fracture(s), 2264, 2265f
 - Malignant hyperthermia, 118
 - Mallet toe, 1027, 103f
 - Malnutrition, with severe cerebral palsy and scoliosis, 1224
 - Mandibulofacial dysostosis, anesthesia in, 125t
 - Mangled Extremity Severity Score (MESS), 1791, 1791t
 - Manipulation
 - for talipes equinovarus, 929, 933f–934f
 - of malreduced supracondylar humeral fractures, 2162–2163, 2164f
 - passive, of foot and ankle, in myelomeningocele, 1263 - Manske classification, of ulnar deficiency, 412f, 413
 - Marfan's syndrome, 1585–1591, 1586f–1588f, 1589t, 1591f
 - anesthesia in, 123t
 - clinical features of, 1586f–1587f, 1586–1588
 - diagnosis of, 1588, 1589t
 - differential diagnosis of, 1588, 1590
 - etiology of, 1585–1586
 - genetic basis of, 1585
 - historical aspects of, 1585
 - incidence of, 1585
 - infantile, 1591
 - kyphosis in, 347–349, 348f
 - prognosis in, 1591
 - scoliosis in, 302–308, 306f–307f
 - back pain in, 304–305
 - curve patterns in, 304, 306f
 - natural history of, 305
 - treatment of
 - nonoperative, 305–306
 - operative, 306, 307f, 308 - treatment of, 1590–1591, 1591f
 - vs. homocystinuria, 1598t

Maroteaux-Lamy syndrome, 1573t, 1580–1581 and kyphosis, 347

 - anesthesia in, 123t

Massage therapy, for rheumatoid arthritis, 1820

Master knot of Henry, mobility of, in talipes equinovarus, 927

McArdle's syndrome, 1491–1492

McFarland operation, for varus fifth toe, 1024, 1030p–1031p

McGregor's line, in Klippel-Feil syndrome, 176–177, 182f

McKay tendon transfer(s), for correction of dorsal bunion, 1372–1373, 1379f, 1386f–1387f

McKusick cartilage-hair hypoplasia, 1549

MCL. *See* Medial collateral ligament (MCL).

MCP joint(s). *See* Metacarpophalangeal (MCP) joint(s).

McRae's line, in Klippel-Feil syndrome, 176–177, 182f

MDP (methylene diphosphonate), 166–167

Meary's angle, 908, 910f

 - in pes cavus, 988, 989f

Mechanical axis

 - in femur, 1103, 1106f
 - deviation of, 1112f
 - vs. anatomic axis, in lower extremities, 2322f

MED. *See* Multiple epiphyseal dysplasia (MED).

Medial collateral ligament (MCL), injuries to

 - diagnosis of, 2363–2364
 - mechanism of, 2362
 - treatment of, 2365

Medial patellofemoral ligament, anatomy of, 2351, 2351f

Median nerve

 - dissection of, in amputation through arm, 2018p–2019p
 - injury to
 - from elbow dislocation, 2193, 2193f
 - from supracondylar humeral fractures, 2148f
 - protection of, in Steindler flexorplasty, 1414p–1415p

Melatonin, and spinal growth, 218

Melorheostosis, 1559f–1560f, 1559–1561

 - clinical course of, 1560–1561
 - clinical features of, 1560
 - differential diagnosis of, 1560
 - etiology of, 1559
 - orthopaedic management of, 1561
 - pathology of, 1559–1560
 - radiographic features of, 1559f–1560f, 1560

Menarche, and progression of curve in scoliosis, 215

Menelaus method, for prediction of remaining femoral/tibial growth, 1051, 1055f

Meningeal sac, mobilization of, in lumbar kyphectomy, 1282p–1285p

Meninges, pathology of, in myelomeningocele, 1252

Meningocele, 1249

Meniscus

 - anatomy of, 2366, 2366f
 - discoid lateral. *See* Discoid lateral meniscus.
 - injuries to, 2366–2367, 2366f–2367f

Mental retardation

 - in homocystinuria, 1597
 - in myotonic dystrophy, 1487

Merchant technique, for radiography of patellar dislocation, 796f–797f

Merosin, deficiency of, in congenital muscular dystrophy, 1481

Meryon's sign, in Duchenne muscular dystrophy, 1469

MESS (Mangled Extremity Severity Score), 1791, 1791t

Metabolic disorder(s)

 - of bone, 1685–1740. *See also under specific disorder, e.g., Osteogenesis imperfecta.*
 - pathophysiology of, 1685, 1686f

of muscle, 1489–1491

 - with bone dysplasia, classification of, 1509t–1510t

Metacarpal bone(s), fractures of, 2249–2250

Metacarpal joint, supracondylar osteotomy of, for radial polydactyly, 430, 431f

Metacarpal joint(s)

 - abduction/adduction contractures of, in cerebral palsy, 1223f–1224f
 - adduction contractures of, in cerebral palsy, 1223f–1224f
 - shortening of
 - in Hurler's syndrome, 1575, 1576f
 - in pseudohypoparathyroidism, 1703f, 1704

Metacarpophalangeal (MCP) joint(s)

 - flexion/extension contractures of, in cerebral palsy, 2192f
 - fusion of, 976f
 - reconstruction of, for rheumatoid arthritis, 474

"Metaphyseal blanch" sign, in slipped capital femoral epiphysis, 719, 719f

Metaphyseal chondrodysplasia, 1547–1549, 1548f–1549f

 - Jansen type, 1547
 - Schmid type, 1547–1548, 1548f–1549f

Metaphyseal cyst(s), in Legg-Calvé-Perthes disease, 677–678, 678f

Metaphyseal dysplasia, genu valgum from, 863

Metaphyseal fracture(s), of distal forearm, treatment of, 2235–2237, 2237f

Metaphyseal line(s), dense transverse, 132, 132f

Metaphyseal osteotomy, opening wedge, for angular deformities of leg, 1113–1114, 1115f

Metaphyseal-diaphyseal angle, in infantile tibia vara, 843, 843f

Metaphysis

 - cellular anatomy of, and acute hematogenous osteomyelitis, 1842, 1843f–1844f
 - cupping of, in rickets
 - nutritional, 1687f–1688f
 - vitamin D-resistant, 8f - femoral, changes in, in Legg-Calvé-Perthes disease, 675–678, 678f–679f

Metaphysitis, syphilitic, 1869, 1869f

Metastatic disease. *See also under specific site, e.g., Lung(s); under specific tumor type, e.g., Osteosarcoma.*

 - in bone, vs. osteopoikilosis, 1557

Metatarsal base

 - transfer of peroneus longus tendon to, for anterior tibial muscle paralysis, in poliomyelitis, 1365, 1365f, 1366p–1371p
 - transfer of tibialis anterior tendon to, in poliomyelitis, 1364

Metatarsal head

 - avascular necrosis of, and Freiberg's infarction, 901
 - first, osteochondritis dissecans of, and hallux rigidus, 1021f
 - transfer of toe extensor to, for pes cavus, 991, 994, 995f, 996p–997p

Metatarsal joint

 - duplication of, with polydactyly of toes, 1022, 1022f
 - fractures of, 2429–2430, 2430f
 - fifth, 2430–2431, 2430f–2433f
 - of foot, ossification of, 892f
 - tendon transfers to, for correction of dorsal bunion, 1372–1373, 1379f, 1386f–1387f

Metatarsal osteotomy

 - for hallux valgus, 1016, 1017f, 1018
 - for pes cavovarus, in Charcot-Marie-Tooth disease, 1436, 1439f
 - for pes cavus, 994–995, 998f–999f

Metatarsocuneiform joint, shape of, and hallux valgus, 1013, 1013f–1014f

Metatarsophalangeal (MTP) joint

 - arthrodesis of, for hallux valgus, in cerebral palsy, 1156, 1164f
 - first, fusion of, for hallux valgus, 1018f, 1018–1019
 - for free bone graft, 511f

Metatarsus adductus, 901–903, 902t, 902f–906f

 - and developmental dysplasia of hip, 517, 517f

- Metatarsus adductus (*Continued*)
 and intoeing, 65–66, 66f
 clinical features of, 902t, 902–903, 903f
 etiology of, 902
 incidence of, 902
 pathology of, 902
 radiographic features of, 902
 severity of, 901, 902f
 terminology used in, 901–902
 treatment of, 903, 904f–906f
- Metatropic dwarfism, 1673, 1674f–1677f, 1677
- Methylene diphosphonate (MDP), 166–167
- Methylprednisolone, for spinal cord injury, 2106–2107
- Meyerding grade, in spondylolisthesis, 356, 358f
- Meyers and McKeever classification, of tibial spine fractures, 2354, 2355f
- MIC2* gene, in Ewing's sarcoma, 2033
- Micromelia, rhizomelic, in achondroplasia, 1510
- Microsurgery, 503–504, 505f–511f, 512
 for free joint transfer, in radial clubhand, 511, 511f
 for free motorized muscle transfer, 504, 507f
 for free tissue transfer, 504, 505f–506f
 for free toe transfer, 504, 508, 508f
 for free vascularized bone transfer, 508, 509f–511f, 512
 for replantation, 503–504
 revascularization for, 504
- Midcarpus, closing wedge osteotomy of, in arthrogryposis, 1655
- Midfoot
 ischemic necrosis of, from correction of talipes equinovarus, with residual deformity, 953, 957f
 osteotomy of, for pes cavus, 999, 1003
 varus deformity of, in talipes equinovarus, 926
- Midfoot break, equinus deformity and, in cerebral palsy, 1136, 1266f
- Midpalmar space, infections of, 478
- Midtarsal dome osteotomy, for pes cavus, 999, 1003
- Midthigh amputation, for osteosarcoma, 1982p–1991p
- Milch classification, of humeral fractures, lateral condylar, 2177, 2179f
- Milwaukee brace
 for idiopathic scoliosis
 adolescent, 229–230, 230f
 infantile, 277
 for postirradiation kyphosis, 342, 344f
 for postural kyphosis, 324f
 for Scheuermann's kyphosis, 328, 329f
- Mirror hand, 423–426, 425f
 treatment of, 432
- Mitchell osteotomy, for hallux valgus, 1015–1016, 1016f
- Mitten cast(s), 384
- Möbius syndrome, 1756f
 anesthesia in, 123t
- "Moccasin deformity," in Proteus syndrome, 1668, 1669f
- Moiré topography, in scoliosis, 226, 227f
- Moist heat, during acute phase of poliomyelitis, 1323–1324
- Monoplegia, definition of, 44
- Monostotic fibrous dysplasia, 1909
- Monteggia fracture(s), 2218–2225, 2220f–2224f
 anatomic considerations in, 2218–2219
 classification of, 2219
 complications of, 2222–2225, 2224f
- Monteggia fracture(s) (*Continued*)
 diagnosis of, 2219
 mechanism of, 2219
 radiographic features of, 2219, 2220f
 treatment of, 2220–2222, 2221f–2223f
- Morel-Lavallee lesion, 2256, 2256f
- Moro reflex, 46–49, 51f, 1126
 abnormalities of, 48–49
- Morphine sulfate, intravenous, 113, 113t
- Morquio syndrome, 1573t, 1577–1580, 1578f–1580f
 clinical features of, 1577–1579
 diagnosis of, 1577–1580, 1578f–1580f
 genetic features of, 1577–1580, 1578f–1580f
 orthopaedic management of, 1579–1580, 1579f–1580f
 pathology of, 1577
- Morquio-Ullrich syndrome, anesthesia in, 123t
- Mose classification, of Legg-Calvé-Perthes disease, 687
- Moseley straight line graph(s), for prediction of remaining femoral/tibial growth, 1051–1053, 1056p–1057p, 1059f
- Motor and sensory neuropathy, hereditary, 1435–1456. *See also under specific type, e.g., Charcot-Marie-Tooth disease.*
 classification of, 1436t
- Motor function
 brain disorders and, 1121, 1122t
 fine, assessment of, in cerebral palsy, 1126
 in poliomyelitis, during convalescent phase, 1324–1325, 1325f
- Motor skill(s), developmental milestones in, 4f
 fine, 20t, 20–21
 gross, 19–20, 20t
- MRI. *See Magnetic resonance imaging (MRI).*
- MTP joint. *See Metatarsophalangeal (MTP) joint.*
- Mucin clot(s), analysis of, 1813, 1814t
- Mucopolysaccharidosis, 1572–1581, 1573t, 1574f–1580f
 and kyphosis, 346–347, 347f
 anesthesia in, 122t–124t
 classification of, 1573t
 diagnosis of, 1572, 1574
 radiographic features of, 1574
 type I, 1573t. *See also Hurler's syndrome.*
 type II, 1573t, 1576–1577
 type III, 1573t, 1577
 type I-S, 1573t
 type IV, 1573t. *See also Morquio syndrome.*
 type V, 1573t, 1580
 type VI, 1573t, 1580–1581
 and kyphosis, 347
 anesthesia in, 123t
- Multidisciplinary collaboration
 in treatment of myelomeningocele, 1260–1261
 in treatment of rheumatoid arthritis, 467
- Multiple enchondromatosis, 1927–1929, 1928f–1929f
- Multiple epiphyseal dysplasia (MED), 1531–1533, 1532f–1534f
 genetic features of, 1531
 orthopaedic considerations in, 1533f–1534f, 1534
 pathology of, 1531–1532
 radiographic features of, 1532f–1533f, 1533–1534
vs. Legg-Calvé-Perthes disease, 689–690, 694f
- Multiple gestation, and incidence of cerebral palsy, 1123
- Multiple injury(ies), 2072–2074, 2073f, 2073t–2074t
 assessment of, 2073–2074, 2073t–2074t
 management of, ABCs of, 2072
 response of children to, 2072
- Multiple pterygium syndrome, *vs. arthrogryposis*, 1652
- Muscle
 biopsy of
 in Becker muscular dystrophy, 1478
 in dermatomyositis/polymyositis, 1493
 in Duchenne muscular dystrophy, 1469
 calcification of, in myositis ossificans, 1497, 1497f
 fibrosis of, idiopathic, 1499
 hemorrhage in, from hemophilia, 1882f, 1882–1883
 treatment of, 1886
 limb lengthening and, 1087, 1087f
 metabolic disorders of, 1489–1491
 pressure in, and compartment syndrome, 2074
- Muscle contraction(s)
 during gait, 79t, 79–81, 80f
 prolonged, in myotonic dystrophy, 1487, 1488f
- Muscle fiber(s), loss of, in muscular dystrophy, 1465, 1466f
- Muscle imbalance, and pes cavus, 984
- Muscle spasms
 in poliomyelitis, during acute phase, 1323
 with joint inflammation, 1811
- Muscle strain, back pain from, 98
- Muscle strength
 assessment of, 44–46, 45t–48t
 grading of, 45t
 Lovett and Martin's system of, 45t
 Texas Scottish Rite Hospital's system of, 45, 46t
 kinetic, 44
 manual tests of, 59–61
 static, 44
- Muscle stretch reflex(es), 56
- Muscle transfer, free motorized, microsurgery for, 504, 507f
- Muscle weakness
 assessment of, 44–46
 clinical features of, 44
 in dermatomyositis/polymyositis, 1494
 in muscular dystrophy
 Duchenne, 1467
 Emery-Dreifuss, 1479
 facioscapulohumeral, 1480, 1480f
 in myasthenia gravis, 1500
 in poliomyelitis, 1322
 during acute phase, 1323
 during convalescent phase, assessment of, 1324
- Muscular atrophy, infantile, 1313t
- Muscular dystrophy, 1463–1484
 Becker, 780f, 1477–1478
 clinical features of, 1464t
 biochemical features of, 1465–1466
 classification of, 1463, 1465t
 clinical features of, 1464t
 congenital, 1481, 1484
 definition of, 1463
 Duchenne, 1467–1477, 1468f–1477f, 1472p–1475p
 anesthesia in, 121t
 clinical features of, 1464t, 1467, 1468f
 diagnosis of, 1469
 etiology of, 1463

- Muscular dystrophy (*Continued*)

medical problems with, 1469

physical examination in, 1468–1469, 1469f

treatment of, 1469–1477, 1470t, 1471f, 1472p–1475p, 1476f–1477f

gene therapy in, 1477

in upper extremities, 1477

lower limb surgery in, 1469–1470, 1470t, 1471f, 1472p–1475p

physical therapy in, 1469

spinal surgery in, 1470–1471, 1476f–1477f

vs. polymyositis, 1467t

electromyography in, 1467

Emery-Dreifuss, 1478–1479

clinical features of, 1464t

etiology of, 1463, 1465

facioscapulohumeral, 1480–1481, 1482p–1483p

clinical features of, 1464t, 1480f, 1480–1481

infantile, 1481

laboratory features of, 1481

treatment of, 808p–809p, 1481

historical aspects of, 1463, 1464f

limb-girdle, 1479–1480

clinical features of, 1464t

limping from, 88–89

myotonic

anesthesia in, 123t

etiology of, 1465

nerve conduction velocity in, 1467

pathology of, 1465, 1466f

progressive, differential diagnosis of, 1313t

scapuloperoneal, 1481

toe-walking in, 1135

Musculoskeletal system, injuries to, with pelvic fractures, 2256

Musculoskeletal tumor(s)

benign, 1901–1948. *See also under specific type, e.g., Bone cyst(s).*

biopsy of, 1899t, 1899–1900

classification of, 1899t, 1900

diagnosis of, 1897–1899, 1898t

imaging studies in, 1898–1899

physical examination in, 1897

staging of, 1899, 1899t

treatment of, 1899t, 1900

Mustard iliopsoas transfer, Sharrard

modification of, for gluteus medius paralysis, 1336, 1338p–1347p

Mustard procedure, 1273–1274, 1274f–1275f

Myasthenia congenita, anesthesia in, 123t

Myasthenia gravis

clinical features of, 1502–1503

diagnosis of, 1503

differential diagnosis of, 1313t

etiology of, 1502

incidence of, 1502

treatment of, 1503

Myasthenic syndrome(s), congenital, 1502

Mycobacterial infection(s), of hand, 479

Mycotic osteomyelitis, 1869–1870, 1870f

Myelography, with computed tomography, in lumbar disk herniation, 372f, 372–373

Myelomeningocele

ambulation in, 1253, 1255

level of lesion and, 1256–1257

clinical features of, scope of, 1249

closure of sac in, 1257

embryology of, 1250, 1251f

etiology of, 1251

Myelomeningocele (*Continued*)

foot and ankle in, 1262–1269, 1265f, 1266p–1267p, 1268f, 1270p–1271p

calcaneus deformity of, 1264–1265, 1265f, 1266p–1267p

equinus deformity of, 1263

passive manipulation of, 1263

rotational deformities of, 1269, 1272f

talipes equinovarus of, 1263–1264

valgus deformity of, 1265, 1268f, 1269, 1270p–1271p

vertical talus of, 1265

general health problems in, 1255–1259, 1258f

cognitive function and, 1255

diastematomyelia and, 1258

hydrocephalus and, 1257

hydromyelia and, 1258

of upper extremity, 1255

precocious puberty and, 1255

psychosocial, 1255–1256

tethered spinal cord and, 1258, 1258f

universal, 1255

urologic, 1258–1259

with lower lumbar level lesions, 1256–1257

with sacral level lesions, 1257

with thoracic level lesions, 1256

with upper lumbar level lesions, 1256

genetic factors in, 1251–1252

hip in, 1272–1277, 1274f–1277f

abduction/external rotation contractures of, 1272–1273

flexion deformity of, 1273

subluxation/dislocation of, 1273–1277, 1274f–1277f

external oblique muscle transfer for, 1274, 1276f

iliopsoas tendon transfer for, 1273–1274, 1274f–1275f

natural history of, 1273

pelvic osteotomy for, 1274, 1277f

varus osteotomy with tendon transfer for, 1274

incidence of, 1249–1250

knee in, 1269, 1272

long-term outcome in, 1253–1255

management of

fetal surgery in, 1257

multidisciplinary, 1260–1261

orthopaedic, 1259–1262. *See also under specific body part, e.g., Myelomeningocele, hip in.*

fractures and, 1262, 1263f

goals of, 1259

infection risk and, 1261

latex risk and, 1261

neonatal examination in, 1259–1260

periodic assessment in, 1260–1261

pressure sores and, 1261–1262, 1262f

orthoses in, 1290–1294, 1291f–1294f

ankle-foot, 1292, 1293f

for lower lumbar/sacral level lesions, 1292

for thoracic/upper lumbar level lesions, 1291–1292, 1292f

hip-knee-ankle-foot, 1293–1294, 1294f

knee-ankle-foot, 1292–1293, 1293f–1294f

principles of, 1290–1291

reciprocating gait, 1294–1295, 1295f

parapodiums in, 1295

principles of, 1253

Myelomeningocele (*Continued*)

role of neurosurgeon in, 1257

skin care in, 1295–1296

standing frames in, 1291, 1292f, 1295

ventriculoperitoneal shunting in, 1257

vocational rehabilitation in, 1296

wheelchairs in, 1295

natural history of, 1253

outcome in, long-term, 1253–1255

pathology of, 1252f, 1252–1253, 1254f

spine in, 1278f–1279f, 1282p–1291p

general management of, 1281, 1289, 1289f

hemimyelodysplasia of, 1291f

hyperlordosis of, 1289

kyphosis of, 1277–1280, 1278f–1279f

and pressure sores, 1279, 1279f

lumbar kyphectomy for, 1280, 1282p–1287p

partial vertebrectomy for, 1279, 1279f

patterns of, 1277, 1278f

radiographic features of, 1278f–1279f

scoliosis of, 1278f, 1280–1281

spinal fusion for, 1281, 1288–1289

scope of deformity in, 1277, 1278f

Myoelectric prostheses, 1800–1801

Myoglobinuria, paroxysmal, idiopathic, 1492

Myoneural level, neuromuscular disorders at, and motor function, 1121, 1122t

Myophosphorylase, deficiency of, 1491–1492

Myositis, suppurative, 1496

Myositis ossificans, 1496–1498, 1497f

anesthesia in, 123t

classification of, 1497

clinical features of, 1497

differential diagnosis of, 1498

from elbow dislocation, 2193

from humeral fractures, supracondylar, 2168, 2175f

from supracondylar humeral fractures, 2168, 2175f

imaging studies in, 1497f, 1497–1498

in hemophilia, 1885

pathology of, 1497

treatment of, 1498

vs. solitary osteochondroma, 1920

Myositis ossificans progressiva. *See Fibrodysplasia ossificans progressiva.*

Myostatic contracture(s), 45

Myotonia congenita, 1489, 1490t

anesthesia in, 123t

Myotonia, Thomsen's, 1489, 1490t

Myotonic dystrophy, 1487, 1488f, 1490t

Myotonic muscular dystrophy

anesthesia in, 123t

etiology of, 1465

Navicular

accessory, 895–899, 897f–898f

classification of, 895–896

clinical features of, 896

complications of, 897–899

imaging studies of, 896, 897f–899f

pathology of, 896

treatment of, 896

results of, 897

bipartite, 896, 898f

dorsal subluxation of, after correction of talipes equinovarus, 942

mobility of, in talipes equinovarus, 927

- Navicular (Continued)**
 subluxation of, in vertical talus, 959, 961f, 965f
- Neck**
 extension of, in Moro's reflex, 50f
 heterotopic ossification of, in fibrodysplasia ossificans progressiva, 1634, 1635f–1636f
- Neck righting reflex**, 54–55
- Neck-shaft angle**
 femoral, in developmental coxa vara, 767–768, 767f–768f, 770, 770f
 of femur, in hip subluxation, with cerebral palsy, 1187, 1189f
- Neolimbus**, development of, 520, 521f
- Neonatal history**, in orthopaedic history, 24
- Neonate(s)**
 flexion contractures in, physiologic, 3, 5f, 36, 38f
 hip instability in, and natural history of developmental dysplasia, 522–523
 joint range of motion in, 28–29
 new periosteal bone formation in, 131, 131f
 osteosclerosis in, 131, 131f
 screening examination of, 64
- Neoplastic disease**
 back pain from, 104–105, 104f–105f
 limping from, 93, 93f
- Nephrocalcinosis**, from phosphorus supplementation, for vitamin D-resistant rickets, 1694
- Nephropathy**, nonhereditary multicentric osteolysis with, 1571
- Nerve(s)**
 fatty infiltration of, in macrodactyly, 451f, 452
 injury to, from limb lengthening, 1088
 palsy of, in hemophilia, 1883–1884
- Nerve conduction velocity**
 in Charcot-Marie-Tooth disease, 1436, 1437f
 in muscular dystrophy, 1467
- Nerve root(s), peripheral**, pathology of, in myelomeningocele, 1252
- Nerve sheath tumor(s)**, of upper extremities, 503
- Nerve-stripping**, for macrodactyly, 457, 457f
- Neural arch(es)**, and vertebral bodies, synchondrosis between, 130f
- Neural axis abnormality(ies)**
 in congenital spinal anomalies, 282
 in idiopathic scoliosis
 infantile, 277
 juvenile, 280
- Neural canal**, rotatory translation of, 298f
- Neural level**, neuromuscular disorders at, and motor function, 1121, 1122t
- Neural tube defect(s)**. *See Myelomeningocele.*
- Neurapraxia**
 in brachial plexus palsy, 480
 in hemophilia, treatment of, 1889
 limb lengthening and, 1090
- Neuritis**, interstitial, hypertrophic, 1445, 1447
- Neuroectodermal tumor(s)**, and postirradiation kyphosis, 341f
- Neurofibroma**, of upper extremities, 502
- Neurofibromatosis**, 1624–1631, 1624f–1631f, 1625t
 and congenital tibial pseudarthrosis, 867–868, 1628, 1630f
 and kyphosis, 300–301, 302f
 and scoliosis, 299–302, 299f–305f
 dystrophic, 299–301, 302f
 treatment of, 300–302, 303f–305f
 nondystrophic, 299, 300f–301f
- Neurofibromatosis (Continued)**
 anesthesia in, 125t
 clinical features of, 196, 198, 198f
 diagnosis of, computed tomography in, 145f
 historical perspective on, 1624, 1624f
 macrodactyly with. *See Von Recklinghausen's disease.*
 radiographic features of, 199
 type 1, 1624–1629, 1625t, 1626f–1631f
 anesthesia in, 125t
 clinical features of, 455f, 455–456, 1625–1628, 1626f–1628f
 cognitive deficits in, 1628–1629
 diagnosis of, 1625, 1625t
 genetic basis of, 1625
 malignant degeneration in, 1628
 multisystem effects of, 1625
 skeletal anomalies in, 1628, 1629f–1631f
 type 2, 1629, 1631, 1631t
- Neurofibrosarcoma**, malignant, 105
- Neurogenic torticollis**, 191–193, 193f
- Neurologic assessment**, 46–56, 49t, 50f–55f
 in Larsen's syndrome, 1603–1604
 in scoliosis, 223–224, 223f–224f
 of back pain, 97
 preoperative, 239
 manual muscle testing in, 59–61
 of back pain, 97
 in scoliosis, 97
 of developmental reflexes, 46–56, 49t, 50f–55f. *See also Reflex(es).*
- Neurologic disorder(s)**
 and kyphosis, in neurofibromatosis, 302 and scoliosis, 218
 from instrumentation for scoliosis, 258
 from Klippel-Feil syndrome, 176
 in lumbar disk herniation, 371
 in spinal dysraphism, 1302
 limping from, 88–89
- Neuromuscular disorder(s)**, 1121–1247. *See also under specific disorder(s), e.g., Cerebral palsy.*
 levels affected by, 1121, 1122t
- Neuromuscular unit(s)**, guiding, in poliomyelitis, 1322
- Neuropathic arthropathy**, 1825–1826, 1826f–1827f
- Neurotmesis**, in brachial plexus palsy, 480
- Neurulation defect(s)**, 1250
- Nevus (nevi)**, in Klippel-Trénaunay syndrome, 1671, 1671f
- Newington brace**, for Legg-Calvé-Perthes disease, 696, 697f
- Niemann-Pick disease**, 1583
 anesthesia in, 123t
- Nievergelt-Pearlman syndrome**, with tarsal coalition, 976, 976f
- Noack's syndrome**, anesthesia in, 123t
- Nonsteroidal anti-inflammatory drug(s) (NSAIDs)**
 for rheumatoid arthritis, 1820
 postoperative, 112–113
- Noonan's syndrome**, anesthesia in, 124t
- NPO requirement(s)**, before anesthesia, 110, 110t
- NSAID(s)**. *See Nonsteroidal anti-inflammatory drug(s) (NSAIDs).*
- Nuclear imaging**, 165–168, 166f–167f
 advantages of, 167
 principles of, 165–166
 radiopharmaceuticals for, 165–167
- Nucleated cell count**, in synovial fluid, 1853
- Nursemaid's elbow**, 2206f, 2206–2207
- Nutritional rickets**. *See Rickets, nutritional.*
- Ober's fasciotomy**, for iliotibial band contractures, in poliomyelitis, 1331, 1337f
- Ober's test**, 40, 41f
 in Duchenne muscular dystrophy, 1469
- Obesity**, and adolescent tibia vara, 850–851, 852f
- Obturator nerve**
 dissection of, in hip disarticulation, 1972p–1973p
 protection of, during hip adduction contracture release, 1180p–1181p
- Occipital-atlantal anomaly(ies)**, 206, 209
- Occipital-atlantal assimilation**, in Klippel-Feil syndrome, 184f–185f, 185
- Occipital-atlantal dislocation**, 2091–2092, 2091f–2093f
- Occipital-atlantal instability**
 in Down syndrome, 1618, 1619f
 nontraumatic, 209
- Occipitocervical arthrodesis**
 Dewar's technique of, 185, 186f, 187
 for Klippel-Feil syndrome, 185–187, 186f
- Occiput-C5 fusion**, for chondrodysplasia punctata, 197f
- Occupational therapy**, for rheumatoid arthritis, 1820
- Ocular torticollis**, 194–195
- Oculoauriculovertebral dysplasia**, anesthesia in, 121t
- Odontoid**
 absence/hypoplasia of
 in metatropic dwarfism, 1673, 1676f
 in Morquio syndrome, 1578f, 1579, 1580f and C2 body, synchondrosis between, 130–131, 132f
 anomalies of, 209
 dysplasia of, in Down syndrome, 1618
 fractures of, 2092–2096
- Ogden classification**, of proximal tibiofibular joint dislocation, 2359, 2360f
- Olecranon**
 fractures of, 2200, 2201f–2203f
 in osteogenesis imperfecta, 1731, 1731f
 osteotomy of, for T-condylar fractures, 2202, 2204f
- Oligohydramnios**, and developmental dysplasia of hip, 516, 516f
- Ollier's disease**, 503, 1927–1929, 1928f–1929f
 anesthesia in, 124t
- Omer classification**, of radioulnar synostosis, 415
- Omohyoid muscle**, dissection of, in forequarter amputation, 2004p–2005p
- "On-top-plasty,"** for congenital band syndrome, 465f
- Onycho-osteodysplasia**, hereditary, 1599f–1601f, 1599–1602
- "Open-book" pelvic fracture(s)**, 2254f
- Opening wedge osteotomy**
 for angular deformities of leg, 1107, 1110f
 for cubitus varus, from supracondylar humeral fractures, 2168, 2172f–2174f
 for talipes equinovarus, with residual deformity, 948, 950, 950f
 for tibial phyeal arrest, 2405, 2406f
 metaphyseal, for angular deformities of leg, 1113–1114, 1115f

Open-up wedge osteotomy, for genu recurvatum, in poliomyelitis, 1359, 1360f
Ophthalmomarthritis, hereditary progressive, 1593
vs. Marfan's syndrome, 1590
Ophthalmologic disorder(s), in Hurler syndrome, 1575
Opioid(s), 113–114, 113t–114t
Opponens splint, long, for thumb, 396, 396f
Optic glioma, in neurofibromatosis, 1625, 1628
Oral reflex(es), 55–56
Orfit, for splints, 389, 389f
Orofaciodigital syndrome, anesthesia in, 124t
Orthopaedic examination, 25–56, 63–73. *See also* Examination.
focused, 65–70, 65f–70f
screening, 63–65, 64f
Orthopaedic history, 23–24
birth history in, 23–24
chief complaint in, 23, 71
family history in, 23
of growth and development, 24
Orthopaedic shoes, for pes planovalgus, 912
Orthoroentgenography, in leg length inequality, 1043, 1045f
Orthoses
below-knee, plantar flexion-assist, 1326, 1326f
for cerebral palsy, 1131–1132, 1132f–1133f
for cervical kyphosis, 199
for hemophilic arthropathy, 1887
for infantile tibia vara, 845–847, 847f
for leg length inequality, 1058, 1061f
for Legg-Calvé-Perthes disease, 694–696, 695f–698f
for lumbar disk herniation, 373
for myelomeningocele, 1291–1295,
1292f–1295f. *See also*
Myelomeningocele, management of, orthoses in.
for pes planovalgus, 912
for poliomyelitis, 1326f, 1326–1327
for scoliosis, idiopathic, 229–232, 230f–233f
thoracolumbar-sacral, for Scheuermann's kyphosis, 328
UCBL
for pes planovalgus, 914, 914f
for tarsal coalition, 980
Ortolani's test, 63–64, 64f
in developmental dysplasia of hip, 525, 525f
Os acromionale, with scapular fractures, 2123
Os odontoideum, 206, 207f
Os trigonum, 891–894, 895f
Osgood-Schlatter disease, 812–813, 814f
and tibial tuberosity fractures, 2343
Ossification
accessory centers of, 894f
in ankle, 2394–2395, 2399f
after limb lengthening, 1086f, 1086–1087
enchondral, failure of, and achondroplasia, 1510
heterotopic, in fibrodysplasia ossificans progressiva, 1634, 1635f–1636f
in carpal bone, in Larsen's syndrome, 1604, 1606f
secondary centers of
in elbow, 2139, 2139f
in lower extremity epiphysis, age at
appearance of, 8f
in upper extremity epiphysis, age at
appearance of, 6f
Ossification center(s), in cervical spine, 2087, 2088f

Ossifying fibroma. *See* Campanacci's disease.
Osteoarthritis
after hip dislocation, 2282
in multiple epiphyseal dysplasia, 1532f, 1534
in spondyloepiphyseal dysplasia
congenita, 1529
tarda, 1530, 1532f
of hip, femoral anteversion and, 784
scoliosis and, in adults, 216
Osteoblastoma, 1936f–1937f, 1936–1938
back pain in, 104
Osteochondral fracture(s), in patellar dislocation, 2352–2353
Osteochondritis, in femoral head, after treatment of Legg-Calvé-Perthes disease, 701, 704f
Osteochondritis dissecans, 789–793, 790f–792f, 792t
classification of, 790, 792t
clinical features of, 790
diagnosis of, 790–791, 790f–791f
etiology of, 789
hereditary, 789
limping from, 91
natural history of, 790
of capitellum, 2207–2208, 2208f
of first metatarsal head, and hallux rigidus, 1021f
of medial femoral condyle, 791f
pathology of, 790
treatment of, 791–793, 792f, 792t
arthroscopic, 791, 792f, 792t
nonoperative, 791
spongialization in, 793
Osteochondrodysplasia, classification of, 1507t–1508t
Osteochondroma
in hereditary multiple exostosis, 1922f, 1922–1923
excision of, 1924, 1925f
of spinal canal, computed tomography in, 148f
of tibial metaphysis, angular deformity from, 1104f
solitary, 1917–1920, 1919f, 1921t
clinical features of, 1919
differential diagnosis of, 1920
etiology of, 1917
pathology of, 1917–1919
radiographic features of, 1919f, 1919–1920
sarcomatous change in, 1920, 1921t
treatment of, 1920
treatment of, below-knee amputation in, 1996p–1999p
Osteochondromatosis, of upper extremities, 503
Osteochondrosis, 899–901, 899f–901f
of elbow, 2207–2208, 2208f
Osteoclasis
closed, for long bone deformities, in
osteogenesis imperfecta
with percutaneous intramedullary fixation, 1732
without internal fixation, 1731–1732
percutaneous, for malreduced supracondylar humeral fractures, 2162–2163, 2164f
Osteocyte(s), abundance of, in osteogenesis imperfecta, 1722, 1723f
Osteodystrophy, renal. *See* Renal osteodystrophy.
Osteofibrous dysplasia, of tibia and fibula, 1914–1917, 1917f–1919f. *See also* Campanacci's disease.

Osteogenesis imperfecta, 1717–1740
anesthesia in, 124t, 1733
basilar impression in, 1739–1740
classification of, 1718–1720, 1720t–1721t, 1722f
clinical features of, 1723–1725, 1725f
differential diagnosis of, 1727–1729
hyperplastic callus in, 1726–1727, 1728f
incidence of, 1722
laboratory studies in, 1727
long bone deformity in, 1729f–1737f, 1731–1738
closed osteoclasis with percutaneous intramedullary fixation for, 1732
closed osteoclasis without internal fixation for, 1731–1732
intramedullary (Bailey-Dubow) rod for, 1733–1738, 1734f–1737f
open osteotomy with intramedullary fixation for, 1732f, 1732–1733
long bone fractures in, 1730–1731, 1731f
pathology of, 1722–1723, 1723f–1724f
pathophysiology of, 1718, 1719f
prenatal diagnosis of, 1730
prognosis in, 1729–1730
radiographic features of, 1725–1726, 1726f–1728f
rehabilitation in, 1730
spinal deformity in, 1738–1739, 1739f
treatment of, 1729
tumors in, 1726–1727
Osteoid osteoma, 104, 104f, 1933f, 1933–1936, 1935f
clinical features of, 1934
differential diagnosis of, 1934
limping from, 93, 93f
natural history of, 1934
pathology of, 1933, 1933f
radiographic features of, 1934, 1935f
treatment of, 1934–1936
Osteoid tissue, uncalcified
in nutritional rickets, 1686, 1686f
in renal osteodystrophy, 1696f
Osteolysis, 1570–1572, 1571f–1572f
differential diagnosis of, 1571
Gorham's, 1570–1571, 1571f
idiopathic, 1572f
multicentric
hereditary
with dominant transmission, 1570, 1572f
with recessive inheritance, 1570
nonhereditary, with nephropathy, 1571
treatment of, 1571–1572
Osteoma, osteoid. *See* Osteoid osteoma.
Osteomyelitis
acute hematogenous, 1841–1859
clinical features of, 1845–1846
complications of, 1859, 1860f
differential diagnosis of, 1850–1852
epidemiology of, 1841
etiology of, 1841
imaging studies in, 1846–1848, 1847f–1852f
computed tomography in, 1847, 1850f
magnetic resonance, 1848, 1851f
radiography in, 1846, 1847f–1849f
radionuclide scintigraphy in, 1847
ultrasonography, 1848, 1852f
in infants, 1846
laboratory studies in, 1846
pathogens in, 1844t, 1844–1845

- Osteomyelitis (*Continued*)

pathophysiology of, 1841–1843, 1843f–1844f

progression to chronic form, 1849f

treatment of, 1852–1853, 1858–1859

 antibiotics in, 1858–1859

 aspiration in, 1852–1853

 surgical, 1858

brucellar, 1869

chronic, 1859–1860

 recurrent multifocal, 1864

definition of, 1841

diagnostic criteria for, 1842t

from congenital insensitivity to pain, 1448

from femoral shaft fractures, 2319

fungal, 1869–1870, 1870f

historical aspects of, 1841

in Gaucher's disease, 1645

in sickle cell disease, 1863–1864, 1893

limping from, 87

of hand, 479

subacute, 1860–1862, 1861f–1862f

tuberculous, 1865–1868, 1866f–1867f

 clinical features of, 1865, 1866f

 orthopaedic management of, 1867–1868

 radiographic features of, 1865, 1866f

 treatment of, 1866–1867, 1867f

vertebral

 and torticollis, 193–194

 back pain in, 103

 vs. diskitis, 87–88

viral, 1870–1871

vs. acute transient synovitis of hip, 1824

vs. hemarthrosis, 1880

vs. scurvy, 1707

vs. suppurative myositis, 1496

Osteonavicular ligament, plantar, dissection of, in correction of vertical talus, 970p–971p

Osteonecrosis, of proximal femur, from osteomyelitis, 1859, 1860f

Osteopathia striata, 1556, 1556f

Osteoperiosteal flap, for spinal fusion, in Scheuermann's kyphosis, 333

Osteopetrosis, 1550–1554, 1551f–1553f

 benign, 1550

 classification of, 1550

 complications of, 1552–1553

 differential diagnosis of, 1552, 1564t

 genetic features of, 1550

 laboratory features of, 1552

 malignant, 1550

 orthopaedic management of, 1553–1554

 pathology of, 1550, 1551f

 prenatal diagnosis of, 1552

 radiographic features of, 1551–1552, 1551f–1552f

 treatment of, 1553

 with renal tubular acidosis, 1551

Osteoplastic laminectomy, 339

Osteopoikilosis, 1557, 1557f

Osteoporosis, juvenile, idiopathic, 1715–1717, 1716f, 1716t

 back pain in, 102

 vs. osteogenesis imperfecta, 1728

Osteosarcoma, 1955–2040

 back pain from, 105

 biopsy of, 1960–1961

 clinical features of, 1957, 1958f

 common anatomic sites of, 1955

 diagnosis of

 computed tomography in, 147f

 magnetic resonance imaging in, 156f

 Pain

 assessment of, 112, 112f, 112t

 back. *See Back pain.*

 congenital insensitivity to, 1448, 1449t, 1450f–1452f

 pathologic fractures in, 1826f

 in Legg-Calvé-Perthes disease, 670

 Pain management. *See also Analgesia.*

 acute, 112f, 112–114, 112t–114t

 after scoliosis surgery, 241

 Paley classification, of short femur, 1775f

 Palmar dog-ear, from macrodactyly surgery, 457, 458f

 Palmar grasp reflex, 46, 50f

 Palmar incision, for syndactyly, 437–438, 438f–439f

Osteosarcoma (*Continued*)

 differential diagnosis of, 1959

 imaging studies in, 1957, 1958f–1962f, 1959 in osteogenesis imperfecta, 1726–1727

 juxtacortical, 1955

 laboratory studies in, 1959

 metastatic, 2029–2030

 of femur, 1958, 1958f–1959f, 1961p–1962p

 pathology of, 1955–1957, 1956f

 staging of, 1957, 1959–1960

 telangiectatic, 1955

 treatment of, 1962–1963, 1963f, 1964p–2026p, 2027f–2031f

 above-knee amputation in, ischial-bearing, 1982p–1991p

 amputation through arm in, 2018p–2021p

 Banks and Coleman hemipelvectomy in, 1964p–1971p

 chemotherapy in, 1962–1963

 choice of procedure in, 1963, 1963f

 elbow disarticulation in, 2022p–2025p

 forequarter amputation with posterior approach (Littlewood technique) in, 2000p–2013p

 hip disarticulation in, 1972p–1981p

 knee disarticulation in, 1992p–1995p

 limb salvage in, 1963, 2026–2029, 2028f–2030f

 rotationplasty in, 2026, 2027f

 shoulder disarticulation in, 2014p–2017p

Osteosclerosis

 in pyknodysostosis, 1566f, 1566–1567

 of long bones, in melorheostosis, 1559f, 1560

 of newborn, 131, 131f

Osteosynthesis, for intertrochanteric double osteotomy, 612p–613p

Osteotomy. *See under specific type, e.g., Pelvic osteotomy.*

Otoform, for total contact molding inserts, 390, 391f

Otopalatodigital syndrome, 1666, 1667f

Outlet view, of pelvic fractures, 2257, 2258f

Outtoeing

 from cerebral palsy, 1128f

 from tibial torsion, 866

Oval-8 ring splint, 393, 394f

Ovary(ies), intra-abdominal, 137f

Overuse injury(ies)

 limping from, 92

 of distal radial physis, 2238f, 2241

Oxford bone age score, modified, in slipped capital femoral epiphysis, and prediction of contralateral slip, 747

Oxygen consumption, during gait analysis, 1129–1130

Pain

 assessment of, 112, 112f, 112t

 back. *See Back pain.*

 congenital insensitivity to, 1448, 1449t, 1450f–1452f

 pathologic fractures in, 1826f

 in Legg-Calvé-Perthes disease, 670

Pain management. *See also Analgesia.*

 acute, 112f, 112–114, 112t–114t

 after scoliosis surgery, 241

Paley classification, of short femur, 1775f

Palmar dog-ear, from macrodactyly surgery, 457, 458f

Palmar grasp reflex, 46, 50f

Palmar incision, for syndactyly, 437–438, 438f–439f

Pamidronate, for osteogenesis imperfecta, 1729

Pancytopenia syndrome, Fanconi's, 1752

Panner's disease, 2207

Pannus, in rheumatoid arthritis, 1816, 1816f

Pantalar arthrodesis, with ankle fusion, in poliomyelitis, 1388–1389, 1394–1395

Papillomavirus infection(s), of hand, 479

Parachute reaction, 54, 55f

Paralysis

 definition of, 44

 in poliomyelitis, development of, 1323

 periodic, 1489–1491

 familial, anesthesia in, 121t

Paralytic disorder(s), genu recurvatum in, 825

Paramyotonia congenita, differential diagnosis of, 1490t

Paraparesis, spastic, hereditary, 1243–1244

Paraplegia

 definition of, 44

 in tuberculous spondylitis, 1834–1835

 Pott's, 1829

Parapodium, 1505

Paraxial limb deficiency(ies), 1747f, 1747–1748

Parent(s)

 communication with, during physical examination, 72–73

education of

 about hemarthrosis, 1886

 about rheumatoid arthritis, 468, 468f

grieving by, and timing of surgical procedures on hand, 381

Paresis, definition of, 44

Paronychia, 476f, 476–477

Paroxysmal myoglobinuria, idiopathic, 1492

Paroxysmal torticollis of infancy, 193

Pars interarticularis

 defects of, in spondylolisthesis, 354, 356f–357f

 repair of, in spondylolysis, 359, 359f–360f

stress fracture of

 and isthmic spondylolisthesis, 352, 353f

 nuclear imaging in, 166f

Patella

 arterial supply to, 2336, 2336f

 bipartite, 811

dislocation of, 2351f, 2351–2353, 2353f

 anatomic considerations in, 2351, 2351f

 chronic, 810

 classification of, 2351

 complications of, 2353, 2353f

 congenital, 811–812, 813f

 diagnosis of, 2352

 habitual, 810

 in idiopathic fibrosis of quadriceps muscle, 1499

 mechanism of, 2351–2352

 recurrent, 793–799, 794f–799f, 800p–809p, 2353

 clinical features of, 794

 diagnosis of, 794–797, 795f

 computed tomography in, 797

 radiography in, 794–797, 795f–799f

 etiology of, 793–794, 794f

 incidence of, 793

 treatment of, 797–799, 800p–809p

 Dewar-Galeazzi procedure in, 797, 800p–805p

 quadricepsplasty in, 806p–809p, 808

 realignment procedures in, 798–799

 results of, 799

 semitendinosis in, 797, 800p–805p

Volume 1, pgs. 1–710; Volume 2, pgs. 711–1502; Volume 3, pgs. 1503–2438.

- Patella (Continued)**
- treatment of, 2352–2353
 - excessive lateral pressure syndrome of, 810
 - fractures of, 2334–2341, 2336f–2341f
 - anatomic considerations in, 2336, 2336f
 - classification of, 2336–2337
 - complications of, 2341
 - diagnosis of, 2337–2338, 2337f–2338f
 - mechanism of, 2337
 - sleeve, 2337–2341, 2339f
 - treatment of, 2338–2341, 2339f–2341f
 - lateral tethering of, and recurrent dislocation, 793–794
 - position of, measurement of, 798f
 - recurrent subluxation of, 799, 810
 - clinical features of, 799
 - diagnosis of, 799
 - treatment of, 799, 810
 - Patella alta, diagnosis of, radiography in, 796–797, 798f
 - Patellar pain syndrome, 810
 - Patellofemoral angle, measurement of, in patellar dislocation, 796, 798f
 - Patellofemoral joint, instability of, 793–799, 794f–799f, 800p–809p
 - and recurrent dislocation, 793–799, 794f–799f, 800p–809p. *See also Patella, dislocation of, recurrent.*
 - in Down syndrome, 1623
 - in nail-patella syndrome, 1600, 1600f
 - Patellofemoral ligament, medial, anatomy of, 2351, 2351f
 - Pathologic fracture(s)**
 - femoral, 2302, 2303f
 - in congenital insensitivity to pain, 1826f
 - in eosinophilic granuloma, 1941f
 - in fibrous dysplasia, 1913, 1916f
 - in Gaucher's disease, 1645
 - in hemophilia, 1884
 - in osteopetrosis, 1552
 - with bone cyst, 1903
 - Patient education, about rheumatoid arthritis, 468, 468f
 - Patient-controlled analgesia (PCA), 113–114, 114t
 - after scoliosis surgery, 241
 - Pattern-bottom brace, for Legg-Calvé-Perthes disease, 658f
 - Pauciarticular juvenile arthritis, limping from, 88
 - Pauciarticular rheumatoid arthritis, 1816f, 1816–1817, 1817t
 - Pauwels' angle, in hip fractures, 2296, 2299f
 - Pauwels' intertrochanteric Y-osteotomy, for developmental coxa vara, 773, 774f, 775, 776p–779p
 - Pavlik disease, 537
 - Pavlik harness
 - effectiveness of, ultrasonography for demonstration of, 530, 530f–531f
 - for developmental dysplasia of hip complications of, 537
 - in neonate, 534–536, 535t, 535f–537f
 - application of, 535, 535f
 - inadequate flexion with, 535, 536f
 - indications for, 534–535, 535t
 - monitoring of, 535, 536f–537f
 - in young child, 536–537, 538f
 - results of, 536–537
 - for femoral shaft fractures, in infant 0 to 6 months, 2304, 2305f
 - for hip/knee dislocation, in Larsen's syndrome, 1607, 1608f
 - PCA. *See Patient-controlled analgesia (PCA).*
 - PCL. *See Posterior cruciate ligament (PCL).*
 - PCR (polymerase chain reaction), in osteomyelitis, 1844
 - Peak height velocity (PHV), and curve progression in scoliosis, 215f, 215–216
 - Pectenous muscle, dissection of, in open reduction of hip, medial approach to, 552p–555p
 - Pectoralis major muscle
 - clavicular head of, and shoulder motion, 1395
 - dissection of, in forequarter amputation, 2012p–2013p
 - Pectoralis major muscle transfer
 - bipolar, for elbow flexion, in arthrolysis, 1654–1655, 1660f
 - in poliomyelitis, 1418, 1420p–1423p
 - unipolar, for elbow flexion, in arthrolysis, 1655
 - Pectoralis minor muscle, dissection of, in forequarter amputation, 2012p–2013p
 - Pectoralis minor muscle transfer, in poliomyelitis, 1418
 - Pectoralis minor tendon transfer, for subscapularis muscle paralysis, in poliomyelitis, 1407, 1408f
 - Pectoroaxillary incision, for forequarter amputation, 2000p–2001p
 - Pectus excavatum, in Marfan's syndrome, 1586f, 1587
 - Pediatric trauma score, 2071, 2072t
 - Pedicle hook(s), for Texas Scottish Rite Hospital instrumentation, 246p–247p, 252p–253p
 - Pedicle screw(s), for spinal fusion, in idiopathic scoliosis, 237f
 - Pelvic obliquity
 - and gait, 77–78, 78f
 - in cerebral palsy, 1225, 1225f–1226f
 - in poliomyelitis, 1395
 - Pelvic osteotomy
 - Chiari, for hip arthrodesis, for proximal focal femoral deficiency, 1770, 1773f–1774f
 - for acetabular repositioning, 596–632, 618t, 621f, 622p–629p, 630f–633f, 634p–641p
 - Ganz, 633, 642, 643f
 - Pemberton, 596–597, 618, 622p–630p
 - Steel, 632, 642f
 - Tonnis, 632–633
 - for hip subluxation/dislocation
 - in cerebral palsy, 1196–1199, 1197f–1198f, 1200p–1201p, 1202f, 1204p–1207p, 1208f
 - in myelomeningocele, 1274, 1277f
 - Pelvic tilt, and gait, 77, 77t, 78f
 - Pelvis
 - anatomy of, 2251–2252, 2251f–2253f
 - deformity of, in poliomyelitis, 1331
 - fractures of, 2251–2271
 - anatomic considerations in, 2251–2252, 2251f–2253f
 - associated injuries in, 2255–2256
 - classification of, 2252, 2253f–2255f, 2254–2255
 - diagnosis of, 2256f, 2256–2257
 - computed tomography in, 2258
 - mechanism of, 2255
 - radiographic features of, 2257–2258, 2257f–2259f
 - Pelvis (Continued)
 - treatment of, 2258–2264, 2259f–2265f, 2264–2271, 2266f–2272f
 - avulsion type, 2259, 2259f–2260f, 2260
 - external fixation in, 2264–2266, 2266f
 - iliac wing, 2261, 2261f
 - internal fixation of sacroiliac joint in, 2267–2268, 2268f–2269f
 - open reduction/internal fixation of symphysis pubis in, 2266–2267, 2267f
 - pelvic ring disruption, 2263–2264, 2263f–2265f
 - straddle, 2263, 2263f
 - rotation of, and gait, 77t
 - Pemberton osteotomy, 596–597, 618, 620f, 622p–630p
 - bone grafts for, 628p–629p
 - complications of, 596, 618
 - contraindications to, 596
 - dissection in, 622p–625p
 - indications for, 596
 - positioning for, 622p–623p
 - postoperative care in, 628p
 - protection of sciatic nerve in, 624p–625p
 - results of, 620f
 - Percutaneous epiphysiodesis, for leg length inequality, 1062, 1072f
 - Perdriolle torsionometer, 225, 226f
 - Perichondral ring complex, and slipped capital femoral epiphysis, 715
 - Periosteal bone, formation of, in Caffey's disease, 1563f, 1564
 - Periosteal elevator(s), Cobb, for Texas Scottish Rite Hospital instrumentation, 250p–251p
 - Periosteal sleeve, in stabilization of supracondylar humeral fractures, 2145, 2148f
 - Periosteal stripping, for limb lengthening, in leg length inequality, 1078
 - Periosteum, osteosarcoma in, 1955, 1956f
 - Peripheral nerve(s)
 - injury to, from supracondylar humeral fractures, 2165
 - limb lengthening and, 1087
 - Peripheral nerve root(s), pathology of, in myelomeningocele, 1252
 - Peripheral neuropathy, 1435–1456. *See also under specific type, e.g., Charcot-Marie-Tooth disease.*
 - Peripheral reflex(es), 56
 - Periventricular leukomalacia, in cerebral palsy, 1124
 - Peroneal muscle
 - defect of, and talipes equinovarus, 923
 - paralysis of, in poliomyelitis, 1362–1364, 1364f, 1365, 1372f, 1374p–1377p
 - Peroneal nerve
 - common, protection of, in epiphysiodesis, for leg length inequality, 1068p–1069p
 - dissection of, in below-knee amputation, 1996p–1997p
 - injury to, from distal epiphyseal femoral fractures, 2332
 - palsy of, 1455–1456
 - Peroneal spastic flatfoot. *See Tarsal coalition.*
 - Peroneal tendon, lengthening of, for pes valgus, in cerebral palsy, 1152
 - Peroneus longus tendon, transfer of to base of second metatarsal, for anterior tibial muscle paralysis, in poliomyelitis, 1365, 1365f, 1368p–1371p
 - to peroneus brevis tendon, for pes cavus, 991

- Peroneus muscle, contractures of, in vertical talus, 964f
- Personal-social skill(s), developmental milestones in, 4f
- Pes calcaneus, limitation of ankle motion in, bone blocks for, 1395
- Pes cavovarus, in Charcot-Marie-Tooth disease, 1436, 1438f–1442f, 1440
- Pes cavus, 984–1011, 984f–990f, 992p–993p, 994f–995f, 996p–997p, 998f, 1000p–1001p, 1002f–1003f, 1004p–1009p clinical features of, 985–988, 985f–989f diagnosis of block test in, 986, 988f neurologic examination in, 986–987 etiology of, 984f, 984–985 in Charcot-Marie-Tooth disease, 995f, 998f, 1002f radiographic features of, 988–989, 989f–990f treatment of, 989–1011, 992p–1009p, 994f–1011f Achilles tendon lengthening in, 991, 995 calcaneal osteotomy in, 995, 999, 1000p–1001p, 1002f–1003f dorsal wedge osteotomy in, 999, 1004p–1005p Dwyer closing wedge osteotomy in, 995, 999, 1000p–1001p, 1002f factors affecting, 989–990 Japas V tarsal osteotomy in, 999, 1006p–1009p metatarsal osteotomy in, 994–995, 998f–999f midfoot osteotomy in, 999, 1003 midtarsal dome osteotomy in, 999, 1003 nonoperative, 989 soft tissue, 990–994, 992p–997p, 994f–995f triple arthrodesis in, 1003, 1010–1011, 1010f–1011f
- Pes planovalgus, 67, 67f, 908–917, 908f–920f clinical features of, 909–912, 911f definition of, 908 diagnosis of neurologic assessment in, 912 radiography in, 908, 910f differential diagnosis of, 908–909 incidence of, 908 Meary's angle in, 908, 910f natural history of, 912 severity of, 908, 909f surgical management of, 914–917, 915f–920f treatment of, 912–914, 913f–914f vs. vertical talus, 961
- Pes varus, in cerebral palsy, 1150f–1151f, 1150–1153, 1154p–1157p, 1164f–1166f calcaneal osteotomy for, 1153 clinical features of, 1150, 1150f peroneal tendon lengthening for, 1152 radiographic features of, 1150–1151, 1150f–1151f subtalar arthrodesis for, 1152 triple arthrodesis for, 1165f
- Pes varus in poliomyelitis, triple arthrodesis for, wedge resection in, 1379, 1389f paralytic, in poliomyelitis, 1364f
- Petrie cast after Salter innominate osteotomy, 640p for Legg-Calvé-Perthes disease, 656, 659f, 695
- Phalanges. *See* Finger(s); Toe(s).
- Phantom limb pain, 1796
- Phemister technique, of epiphysiodesis, for leg length inequality, 1058–1063, 1062f, 1064p–1071p, 1072f
- Philipsson's reflex, 51, 53f
- Phosphor plate(s), for computed radiography, 134
- Phosphorus, for vitamin D-resistant rickets, 1693–1694
- PHV (peak height velocity), and curve progression in scoliosis, 215f, 215–216
- Physcal arrest. *See* Growth arrest.
- Physcal bar, resection of, 2071, 2071f
- Physcal plate, abnormal growth of, in solitary osteochondroma, 1917
- Physical examination. *See* Examination.
- Physical therapy. *See also* Rehabilitation. for cerebral palsy, 1130–1131 in upper extremity, 1211 for Duchenne muscular dystrophy, 1469 for hemophilic arthropathy, 1887 for poliomyelitis after tendon transfer, 1328–1329 during chronic phase, 1325–1326 during convalescent phase, 1324 for recurrent patellar subluxation, 799 for rheumatoid arthritis, 468–469, 1820 for scoliosis, idiopathic, 232
- Physiology, in renal osteodystrophy, 1699, 1700f
- Physis (physes). *See also under individual bone.* anatomy of, 2062–2063, 2063f blood supply to, 2063, 2063f distal, and growth of upper extremity long bones, 10f distraction of, for limb lengthening, 1082–1083 femoral, changes in, in Legg-Calvé-Perthes disease, 678–679, 680f injuries to, 2062–2072, 2064f–2066f, 2068 and Harris growth arrest lines, 2063–2064, 2064f classification of, 2064–2065, 2064f–2065f complications of, 2066–2067, 2066f–2070f, 2070 treatment of, 2065 of ankle, premature closure of, after fractures, 2403–2405, 2404f, 2406f of long bones, yearly growth of, 2070t proximal, and growth of upper extremity long bones, 10f transplantation of, for limb lengthening, 1078 zones of, 1047, 1048f
- Phytanic acid, serum, in Refsum's disease, 1447–1448
- Pierre Robin syndrome, anesthesia in, 124t
- Pigmented nevi, in Proteus syndrome, 1666, 1668
- Pigmented villonodular synovitis, 1944–1946, 1946f–1947f
- Pin site(s), infection of, limb lengthening and, 1090
- PIP joint. *See* Proximal interphalangeal (PIP) joint.
- Pituitary dwarfism, 1711–1712
- PLACE scale, 112t
- Placing reaction, 49, 52f
- Plagiocephaly, in idiopathic scoliosis, infantile, 276, 276f
- Plantar closing wedge osteotomy, for dorsal bunion, 944, 945f
- Plantar flexion, muscles producing, 1361
- Plantar flexion deformity, of forefoot, in pes cavus, 985, 985f
- Plantar flexion-assist below-knee orthosis, 1326, 1326f
- Plantar grasp reflex, 46, 50f
- Plantar osteonavicular ligament, dissection of, in correction of vertical talus, 970–971p
- Plantar release, for pes cavus, 990–991, 992p–993p, 994f–995f
- Plastic deformation, 2059, 2060f
- Plastizote shoe insert(s), for pes planovalgus, 914, 914f
- Platelet(s), calmodulin in, and scoliosis, 218
- Polytyspondyly
- in pseudoachondroplasia, 1525, 1525f
 - in spondyloepiphyseal dysplasia congenita, 1528f
 - tarda, 1530, 1530f
- Plexiform neurofibroma(s), 1625, 1627f–1628f
- Plica syndrome, 810–811
- Plumb line, in diagnosis of scoliosis, 222, 222f
- PMMA (polymethylmethacrylate) bead(s), for wound care, 2383
- PNET(s) (primitive neuroectodermal tumors), 2030f–2037f, 2030–2038, 2032t. *See also* Ewing's sarcoma.
- Poisoning, lead, metaphyseal bands in, 132f
- Poland's syndrome, 1752, 1756f
- synbrachydactyly in, 1783, 1784f
- Poliomyelitis, 1321–1427
- elbow in, 1408–1419, 1409f, 1410p–1426p latissimus dorsi muscle transfer in, 1419, 1426f–1427f
 - pectoralis major muscle transfer in, 1418, 1420p–1423p
 - pectoralis minor muscle transfer in, 1418
 - Steindler flexorplasty of, 1408–1409, 1409f, 1410p–1417p
 - sternocephalic muscle transfer in, 1418–1419, 1424f
 - triceps brachii anterior transfer in, 1419, 1425f
 - triceps brachii paralysis in, 1419
- etiology of, 1321
- foot and ankle in, 1361–1373, 1364f–1365f, 1366p–1385p, 1372f–1387f
- anterior tibial muscle paralysis in, 1365, 1365f, 1366p–1371p
- arthrodesis of, 1373–1395, 1388f–1394f, 1390p–1393p
- distraction-compression bone graft, 80p–85p, 1395
 - extra-articular subtalar, 1388, 1396p–1399p
 - pantalar, with ankle fusion, 1388–1389, 1394–1395
 - triple, 1373–1387, 1388f–1389f, 1390p–1393p
- calcaneus deformity of, operative procedure for, 1365–1373, 1373f, 1379f, 1380p–1385p, 1386f–1387f
- equinovalgus deformity of, operative procedure for, 1364, 1365f
- equinovarus deformity of, operative procedure for, 1365, 1372f, 1374p–1377p
- extensor digitorum longus paralysis in, 1363–1364
- extensor hallucis longus paralysis in, 1363–1364
- peroneal muscle paralysis in, 1362–1364, 1364f, 1365, 1372f, 1374p–1377p
- tendon transfers for, 1362t–1363t

- Poliomyelitis (*Continued*)

 toe extensor muscle paralysis in, 1365, 1372f, 1374p–1377p

 triceps surae muscle paralysis in, 1365–1373, 1373f, 1378f–1379f, 1380p–1385p, 1386f–1387f

 forearm in, 1424–1427

 hip in, 1329–1351

 gluteus maximus paralysis in, 1336–1337, 1348–1349, 1348f

 Barr erector spinae transfer for, 1336–1337, 1348, 1348f

 Hogshead fascia lata transfer to lumbodorsal fascia for, 1349, 1349f

 motor evaluation in, 1336

 gluteus medius paralysis in, 1336, 1338p–1347p

 iliotibial band contractures in, 1329–1331, 1330f, 1332p–1335p, 1336, 1337f

 anatomic considerations in, 1330, 1330f

 and external tibial torsion, 1330–1331

 and flexion/valgus deformity of knee, 1330

 and lumbar lordosis, 1331

 and pelvic deformity, 1331

 and pes varus, 1331

 flexion, abduction and external rotation, 1330

 treatment of, 1331, 1332p–1335p, 1337f

 paralytic dislocation of, 1349–1351, 1352p–1355p

 arthrodesis in, 1350–1351, 1356f

 varization osteotomy in, 1349–1350, 1352p–1355p

 knee in, 1351–1361, 1356f–1361f

 flail, 1360–1361, 1361f

 flexion contractures of, 1358, 1358f

 genu recurvatum in, 1358–1360, 1359f–1360f

 quadriceps femoris paralysis in, 1351, 1356f–1357f, 1356–1358

 management of, 1322–1329, 1325f–1326f

 during chronic phase, 1325–1329, 1326f

 orthoses in, 1326f, 1326–1327

 physical therapy in, 1325–1326

 during convalescent phase, 1324–1325, 1325f

 in acute phase, 1322–1324

 surgical, principles of, 1327–1329

 paralysis in, development of, 1323

 pathology of, 1322

 prediction of remaining growth in, 1049, 1049t

 prognosis in, 1322

 shoulder in, 1395, 1406t, 1406–1408, 1407f–1409f

 arthrodesis of, 1407–1408

 classification of muscles and, 1395, 1406–1408, 1407f–1409f

 tendon transfers for, 1406–1407, 1406t, 1406f–1408f

 trunk in, 1395

 vaccine-associated, 1321

 Polyarticular rheumatoid arthritis, 1817, 1817f–1818f

 Polydactyly, 421–432, 422f–433f, 423t

 central

 classification of, 423, 424f–425f

 clinical features of, 423

 complications of, 432

 treatment of, 431–432

 classification of, 422f, 422–423, 423t

 etiology of, 421
- Polydactyly (*Continued*)

 historical perspective on, 421

 imaging studies of, 426

 in chondroectodermal dysplasia, 1679, 1679f–1680f

 incidence of, 422

 of toes, 1021–1023, 1022f

 with hallux varus, 1019, 1019f

 pathology of, 421–422

 radial, 437–448

 classification of, 422f, 422–423, 423t

 complications of, 432, 433f

 conditions associated with, 423

 treatment of, 426, 426f–431f, 430–431

 in Wassel type I and I thumbs, 426, 426f–427f

 in Wassel type III thumbs, 426

 in Wassel type IV thumbs, 426, 428f–431f, 430–431

 in Wassel type V and VI thumbs, 431

 ulnar

 classification of, 423, 425f

 complications of, 432

 treatment of, 432, 432f

 Polymyxin, for splints, 388f, 389

 Polymerase chain reaction (PCR), in osteomyelitis, 1844

 Polymethylmethacrylate (PMMA) bead(s), for wound care, 2383

 Polymyositis

 differential diagnosis of, 1313t

 vs. Duchenne muscular dystrophy, 1467t

 Polyneuritis

 acute infective, 1313t

 idiopathic, anesthesia in, 122t

 Polyostotic fibrous dysplasia, 1909, 1915f

 Polypropylene ankle-foot orthoses, for myelomeningocele, 1500, 1501f

 Polyradiculoneuritis, acute, 1454–1455

 Polysyndactyly, preaxial, 1024, 1024f

 "Popcorn" calcification, of long bones, in osteogenesis imperfecta, 1725, 1726f

 Popliteal artery

 dissection of, in knee disarticulation, 1994p–1995p

 injury to

 from distal epiphyseal femoral fractures, 2332, 2333f

 from proximal tibial phyeal fractures, 2350–2351

 Popliteal cyst(s), 828f–831f, 829–831, 832p–833p

 clinical features of, 828f, 829

 diagnosis of, 831, 831f

 differential diagnosis of, 829

 histologic classification of, 830f, 831

 pathology of, 829, 829f–830f, 831

 treatment of, 831, 832p–833p

 Popliteal vein, dissection of, in knee disarticulation, 1994p–1995p

 Position

 and spasticity, 27

 during acute phase of poliomyelitis, 1323

 fetal. *See* Fetal position.

 for assessment of cervical spine fractures, 2091, 2091f

 for percutaneous in situ pinning of slipped capital femoral epiphysis, 722, 724p–725p

 neonatal, vertex presentation and, 38f

 postnatal, and developmental dysplasia of hip, 516–517, 517f

 vertical suspension, 49

 Positive support response, 51–52

 Postaxial polydactyly, 1021, 1022f

 Posterior cervical line, 132, 132f

 Posterior cruciate ligament (PCL)

 anatomy of, 2354, 2354f, 2361f, 2361–2362

 avulsion fractures of, 2356

 injuries to, 2361f–2365f, 2361–2366

 diagnosis of, 2362–2364, 2363f–2364f

 mechanism of, 2362

 treatment of, 2365

 Posterior-lateral corner test, 2363, 2364f

 Postlaminectomy kyphosis, treatment of, 204–206, 205f

 Postpolio syndrome, 1321

 Postural clubfoot, *vs.* talipes equinovarus, 927

 Postural kyphosis, 323, 324f, 325t

 Posture

 definition of, 323

 development of, 323

 in cerebral palsy, from hamstring spasticity, 1157, 1166f

 in Duchenne muscular dystrophy, 1467, 1468f

 in septic hip, 1845

 in spondylolisthesis, 354, 355f

 in tuberculous spondylitis, 1833, 1833f

 Pott's disease. *See* Spondylitis, tuberculous.

 Pott's paraplegia, 1831

 Powers ratio, 2092, 2092f

 in occipito-atlantal instability, 1618, 1619f

 Prader-Willi syndrome, anesthesia in, 124t

 Preaxial polydactyly, 1021, 1022f

 Preaxial polysyndactyly, 1024, 1024f

 Precocious puberty, in myelomeningocele, 1255

 "Predisposed child" theory, in Legg-Calvé-Perthes disease, 661–662, 664f

 Pregnancy, screening for, before radiography, 137

 Prehensile function, with bilateral upper limb deficiencies, 1796–1797, 1797f–1798f

 Prematurity, rickets of, 1690

 Prenatal history, in orthopaedic history, 23–24

 Presentation, vertex, and neonatal position, 38f

 Pressure epiphysis, 1045

 Pressure sore(s)

 from improper casting, 2079, 2080f

 in myelomeningocele, 1261–1262, 1262f

 in myelomeningocele from kyphosis, 1279, 1279f

 on foot, prevention of, 1262, 1262f

 Primitive neuroectodermal tumor(s) (PNETs), 2030f–2037f, 2030–2038, 2032t. *See also* Ewing's sarcoma.

 Pro-alpha polypeptide chain(s), in collagen metabolism, 1718, 1719f

 Professional demeanor, during physical examination, 71

 Profundus tendon, injury of, 399f

 Progressive diaphyseal dysplasia, 1555f, 1555–1556

 Progressive noninfectious anterior fusion, 336–337, 339f

 Pronation

 assessment of, 31, 33f

 definition of, 27t

 Pronation contracture(s)

 after Steindler flexorplasty, 1418

 of forearm, in cerebral palsy, 1210, 1211f

 Pronator teres muscle, lengthening of, 1218p–1219p

 Prostaglandin E, and hyperostosis, 1564

- Prosthesis (prostheses)
 equinus, 1802
 fitting of, immediate, 1798
 for above-knee amputation, 1804
 for below-knee amputation, 1804
 for fibular deficiency, 1802–1803
 for foot amputation, 1803
 for foot deficiencies, 1803
 for knee disarticulation, 1804
 for lower limb deficiencies, 1801–1802
 for proximal focal femoral deficiency, 1761, 1761f, 1794
 for short femur, 1773f
 for tibial deficiency, 1782f, 1803
 for upper limb deficiencies, 1796–1798
 lower limb, 1801
 gait problems with, 1801–1802
 myoelectric, for upper extremities, 1800–1801
 principles of, 1799–1800, 1800f
 replacement of, 1800, 1802
 slipper, for congenital foot absence, 1783f
 upper limb, 1800–1801
 vs. allograft, for limb salvage, after osteosarcoma treatment, 2030–2031, 2031f
- Prosthetic foot, solid ankle-cushioned, for leg length inequality, 1058, 1058f
- Protective extension of arms reflex, 54, 55f
- Protein, in synovial fluid, 1813, 1814t
- Protein C, deficiencies in, and Legg-Calvé-Perthes disease, 657
- Protein S, deficiencies in, and Legg-Calvé-Perthes disease, 657
- Proteus syndrome, 1666–1669, 1668f–1669f, 1669t
 clinical features of, 1666–1669, 1668f–1669f
 diagnosis of, 1669, 1669t
 differential diagnosis of, 1669
 orthopaedic management of, 1669
- Protrusio acetabuli, in Marfan's syndrome, 1590, 1591f
- Proximal femoral focal deficiency. *See* Femoral deficiency, proximal focal.
- Proximal interphalangeal (PIP) joint boutonniere deformity of, 476
 flexion/extension splints for, 393, 394f
 reconstruction of, for rheumatoid arthritis, 474, 476f
 resection/arthrodesis of, for hammer toe, 1034p–1035p
 tension spring for, 393, 393f
- Pseudarthrosis
 after *in situ* spinal fusion, for spondylolisthesis, 365
 congenital
 of fibula, 883–884, 884f–886f, 887
 of tibia. *See* Tibia, congenital
 pseudarthrosis of.
 femoral, stabilization of, for proximal focal femoral deficiency, 1763, 1770f
 of forearm, free vascularized bone transfer for, 508, 509f–511f, 512
- Pseudoachondroplasia, 1523–1526, 1524f–1526f and kyphosis, 346
 clinical features of, 1523–1525, 1524t, 1524f–1525f
 genetic features of, 1523
 lumbar lordosis in, 1525f
 orthopaedic features of, 1526
 pathology of, 1523
- Pseudoachondroplasia (*Continued*)
 radiographic features of, 1525–1526, 1525f–1526f
 vs. achondroplasia, 1524t
- Pseudoadduction, of hip, 1178
- Pseudo-aneurysm(s), of upper extremities, 497
- Pseudodislocation, of sternoclavicular joint, 2118, 2120
- Pseudo-Galeazzi sign, 1191
- Pseudohypoparathyroidism, 1703f, 1703–1704
- Pseudometatropic dysplasia. *See* Kniest's dysplasia.
- Pseudosubluxation, of cervical spine, 131–132, 132f, 206f, 2089–2090, 2090f
- Pseudotumor(s), hemophilic, 1884, 1885f
 treatment of, 1889
- Psoas muscle, abscess of, 1850f
- Psoas tendon
 dissection of
 in Pemberton osteotomy, 622p–623p
 in Salter innominate osteotomy, 636p–637p
 lengthening of, for adduction contractures of hip, 1182p–1183p
- Psoas tenotomy, for hip flexion contractures, 1173
- Psoriatic arthritis, 1822–1823, 1823f, 1823t
- Psychological stress, limb lengthening and, 1091
- Psychosocial issue(s). *See under specific disorder*, e.g., Myelomeningocele.
- Psychosomatic back pain, 96, 105–106
- Puberty
 growth spurt during, 1048, 1050f
 precocious, in myelomeningocele, 1255
 Tanner's stages of development during, 19, 19f
- Pubic bone, osteotomy of, in Banks and Coleman hemipelvectomy, 1966p–1967p
- Pulmonary embolism, after spinal cord injury, 2107–2108
- Pulmonary function, in Duchenne muscular dystrophy, 1469
- "Pulseless pink hand," in supracondylar humeral fractures, 2163, 2163f
- Pulvinar, fibrofatty, in developmental dysplasia of hip, 520, 521f
- Puncture wound(s)
 infection from, 1863
 of foot, 2434
- Purpura fulminans, 1792–1793, 1793f, 1796, 1796f
- Putti's sign, 481, 483f
- Pyknodysostosis, 1565–1567, 1565f–1567f, 1566t
 clinical course of, 65
 clinical features of, 1565f, 1566
 differential diagnosis of, 1564t, 1566t, 1567
 genetic features of, 1565–1566
 orthopaedic management of, 1566f–1567f, 1567
 pathology of, 1566
 radiographic features of, 1566f, 1566–1567
 vs. osteogenesis imperfecta, 1728
- Pyogenic granuloma, of upper extremities, 502
- Pyogenic tenosynovitis, of hand, 478, 478f
- Pyomyositis, 1496
- Pyramidal level, neuromuscular disorders at, and motor function, 1121, 1122t
- Pyrazinamide, for tuberculous arthritis, 1831
- Pyridoxidine, for homocystinuria, 1597
- Q angle, in recurrent patellar dislocation, 794, 794f
- Quadriceps muscle
 fibrosis of, progressive, 1499
 misalignment of, and recurrent patellar dislocation, 793
 paralysis of, in poliomyelitis, 1351, 1356–1358, 1356f–1358f
 strength of, and ambulation in myelomeningocele, 1253, 1255
 stretching of, for congenital dislocation of knee, 823
 weakness of, after femoral shortening, 1076–1077
- Quadricepsplasty, for recurrent patellar dislocation, 798, 806p–809p
 dissection in, 806p, 809p
 incision in, 806p, 807p
 postoperative care after, 808p
- Quadriplegia
 definition of, 44
 in cerebral palsy, 1124, 1125f
- Race
 and chondrolysis, as complication of slipped capital femoral epiphysis, 712
 and developmental dysplasia of hip, 514, 514t
 and incidence of slipped capital femoral epiphysis, 712
 and interpretation of radiographs, 128
 and myelomeningocele, 1249–1250
 "Rachitic rosary," 1689
- Racquet incision
 for hip disarticulation, 1972p–1973p
 for radial polydactyly, 426, 429f
- Radial clubhand
 free joint transfer in, 511, 511f
 surgical treatment of, milestones in, 382
- Radial collateral ligament, dissection of, for radial polydactyly, 426, 430, 430f
- Radial deficiency, 412
- Radial head
 dislocation of
 anterior, from brachial plexus palsy, 481
 in achondroplasia, 1511, 1512f
 in Larsen's syndrome, 1611
 in nail-patella syndrome, 1601f
 with ulnar fractures. *See* Monteggia fracture(s).
- traumatic subluxation of, 2206f, 2206–2207
- Radial nerve
 dissection of
 in amputation through arm, 2020p–2021p
 in disarticulation of elbow, 2022p–2023p
 injury to, after humeral fractures, 2134
 palsy of, after Monteggia fractures, 2224–2225
- Radial polydactyly. *See* Polydactyly, radial.
- Radiation, exposure to, during computed tomography, 145
- Radiation therapy
 for aneurysmal bone cyst, 1909
 for Ewing's sarcoma, 2035, 2036f–2037f, 2038
 for soft tissue sarcoma, 2045, 2045f
 kyphosis after, 339–346, 343f–346f. *See also* Kyphosis, postirradiation.
- scoliosis from, 312
- Radicular neuropathy, sensory, and insensitivity to pain, 1449t, 1453

- Radioallergosorbent (RAST) test, for latex allergy, 117
- Radiocapitellar joint, dislocation of, in Larsen's syndrome, 1602, 1604f
- Radiofrequency coagulation, of osteoid osteoma, 1936
- Radiography, 127–137, 128f–136f. *See also under specific disorder.*
- computed, 134–135
 - digital, 135
 - during intertrochanteric varus osteotomy, and internal fixation with blade plate, for open hip reduction, 572p–573p
 - equipment for, 133–134, 134f
 - in differential diagnosis of back pain, 97–98
 - in idiopathic scoliosis, 224–226, 224f–227f
 - in slipped capital femoral epiphysis, 719–720, 719f–720f
 - interpretation of
 - avoidance of tunnel vision in, 128–129
 - gender and, 127–128
 - organ system examination and, 129
 - patient age and, 127
 - previous studies and, 129–130
 - race and, 128
 - signs and symptoms in, 128, 128f
 - soft tissue examination and, 129
 - viewing conditions and, 127
 - normal variants in, 130–132, 130f–133f
 - patient identification for, 135
 - patient preparation for, 135
 - patient protection during, 135–137, 136f–137f
 - pregnancy screening before, 137
 - technique of, 132–137, 134f–137f
 - child-friendly, 132–133
 - value of, 127
- Radiolucent table(s), for open in situ fixation of slipped capital femoral epiphysis, 723, 728f, 728–729
- Radionuclide scintigraphy, in osteomyelitis, 1847
- Radiopharmaceutical(s), for nuclear imaging, 165–167
- Radius
- synostosis
 - clinical features of, 415
 - complications of, 416
 - conditions associated with, 415
 - etiology of, 414–415
 - historical perspective on, 414–415
 - imaging studies of, 415
 - treatment of, 415–416
 - deficiency of, 1787, 1788f
 - fractures of
 - and phyeal arrest, 2068f
 - buckle, 2060f–2061f
 - distal phyeal, treatment of, 2237–2238
 - with ulnar fractures, 2225–2233, 2226f–2234f
 - anatomic considerations in, 2225
 - complications of, 2231–2233, 2234f
 - diagnosis of, 2225
 - malunion of, 2227, 2229, 2230f, 2231–2233, 2234f
 - mechanism of, 2225
 - radiographic features of, 2225, 2226f, 2227
 - treatment of, 2227–2231, 2228f–2232f
 - closed, 2227–2230, 2228f–2230f
 - open, 2230–2231, 2231f–2232f
 - anatomic (Continued)
 - head and neck fractures of, 2194f–2199f, 2194–2200
 - anatomic considerations in, 2194
 - classification of, 2194–2195, 2195f
 - complications of, 2197, 2200
 - diagnosis of, 2194
 - mechanism of, 2194, 2194f–2195f
 - radiographic features of, 2194, 2195f
 - remodeling after, 2197, 2199f
 - treatment of, 2195–2197, 2196f–2199f
 - closed reduction in, 2195–2196, 2196f–2197f
 - intramedullary reduction in, 2197, 2199f
 - open reduction in, 2197
 - percutaneous reduction in, 2196–2197, 2198f
 - physis of, overuse injuries of, 2238f, 2241
 - Rang classification, of physeal injuries, 2065, 2065f
 - Range of motion, 27t, 27–44, 28f–45f, 34t, 38t
 - examination of, 27t, 27–29
 - of ankle, 43–44, 43f–45f
 - of cervical spine, 32–35, 34t, 35f
 - of elbow, 31, 33f
 - of foot, 43–44, 43f–45f
 - of forearm, 31, 33f
 - of hip, 36–42, 38t, 38f–43f
 - age and, 37–38, 38t
 - of knee, 43, 43f
 - of shoulder, 28f–32f, 29–31
 - of thoracolumbar spine, 35–36, 36f–38f
 - Rash
 - in dermatomyositis, 1493f, 1493–1494
 - in rheumatoid arthritis, 1817, 1818f
 - RAST (radioallergosorbent) test, for latex allergy, 117
 - Ray(s), first, dorsiflexion of, for talipes equinovarus, 931, 931f
 - "Razorback" deformity, in scoliosis, 220, 239
 - Reciprocating gait orthoses (RGOs), for myelomeningocele patient, 1294–1295, 1295f
 - Rectus femoris muscle
 - dissection of
 - in open reduction of hip, using anterolateral approach, 562p–563p
 - in Pemberton osteotomy, 622p–623p
 - in Salter innominate osteotomy, 636p–637p
 - spasticity of, after hamstring lengthening, 1159, 1166–1167, 1172f
 - transfer of, for spasticity, 1167–1169, 1174p–1177p
 - Rectus grab, 1166–1167
 - Referring physician(s), communication with, 73
 - Reflex(es), 46–56, 49t, 50f–55f
 - abdominal, 56
 - assessment of, in scoliosis, 223, 223f
 - appearance/disappearance of, normal times for, 49t, 50f
 - assessment of, in cerebral palsy, 1126
 - Babinski's, 56
 - body righting, 55
 - cremasteric, 56
 - crossed extension, 51, 53f
 - deep tendon, 56
 - in cerebral palsy, 1126
 - extensor thrust, 52
 - foot grasp, 46, 50f
 - Galant's, 52, 54f
 - hand grasp, 46, 50f
 - Hoffmann's, 56
 - Reflex(es) (Continued)
 - Landau, 54
 - leg-straightening, 51–52
 - Moro's, 46–49, 51f
 - abnormalities of, 48–49
 - muscle stretch, 56
 - neck righting, 54–55
 - oral, 55–56
 - palmar grasp, 46, 50f
 - parachute, 54, 55f
 - peripheral, 56
 - Philipsson's, 51, 53f
 - placing reaction, 49, 52f
 - plantar grasp, 46, 50f
 - protective extension of arms, 54, 55f
 - rooting, 55–56
 - rotation, 52–53
 - startle, 49, 1126
 - stepping, 49, 51, 53f
 - sucking, 55–56
 - tonic neck, 53–54, 54f–55f
 - trunk incursion, 52, 54f
 - vertical suspension positioning, 49
 - walking, 49, 51, 53f
 - withdrawal, 51
 - Reflex sympathetic dystrophy, 2241
 - Refsum's disease, 1447–1448
 - Regional analgesia, 114–115
 - Regional anesthesia, 115–116
 - Rehabilitation. *See also Physical therapy.*
 - after distraction-compression ankle arthrodesis, in poliomyelitis, 1404p
 - after dorsal wedge osteotomy, for pes cavus, 1004p
 - after epiphysiodesis, for leg length inequality, 1066p, 1070p
 - after excision of discoid lateral meniscus, 820
 - after fractional lengthening of finger and wrist flexors, 1220p
 - after Grice procedure, 1156p
 - after hamstring lengthening, 1170p
 - for iliotibial band contractures, in poliomyelitis, 1334p, 1336, 1337f
 - after hip adduction contracture release, 1182p–1183p
 - after iliopsoas muscle transfer, for gluteus maximus paralysis, 1346p
 - after intertrochanteric oblique osteotomy of proximal femur, 1354p
 - after Japas V tarsal osteotomy, 1008p
 - after latissimus dorsi/teres major muscle transfer to rotator cuff, 490p
 - after lumbar kyphectomy, using Dunn-McCarthy technique, 1286p
 - after McFarland operation, 1030p
 - after pectoralis major transfer, 1422p
 - after proximal hamstring release, for hip subluxation/dislocation, in cerebral palsy, 1194p–1195p
 - after quadricepsplasty, for recurrent patellar dislocation, 808
 - after semitendinosus, for recurrent patellar dislocation, 804p
 - after subtalar extra-articular arthrodesis, 1398p
 - after synovectomy, for hemophilic arthropathy, 1888
 - after tibialis anterior tendon transfer to calcaneus, 1266p–1267p
 - after tibialis posterior tendon transfer to os calcis, for calcaneus deformity, 1384p
 - after use of foot frame, 953
 - after Zancolli procedure, 500p

Rehabilitation (*Continued*)
 in osteogenesis imperfecta, 1730
 vocational, for myelomeningocele patient, 1506

Reiter's syndrome, 1822

Relaxation time (T1), longitudinal, in magnetic resonance imaging, 152, 152f

Relaxation time (T2), transverse, in magnetic resonance imaging, 152, 152f

Renal failure, chronic, and slipped capital femoral epiphysis, 716

Renal osteodystrophy, 1696–1700, 1696f–1700f
 clinical features of, 1697
 laboratory studies in, 1696
 medical treatment of, 1698–1699
 orthopaedic management of, 1699–1700, 1700f
 pathophysiology of, 1696, 1696f
 radiographic features of, 1697–1698, 1697f–1699f

Renal tubular acidosis, osteopetrosis with, 1551

Renshaw classification, of caudal regression syndrome, 1306, 1522f–1525f

Reossification, in Legg-Calvé-Perthes disease, 675, 675t, 677f

Repetition time (TR), in magnetic resonance imaging, 152, 152t

Replantation, microsurgery for, 503–504

Reproductive organ(s), shielding of, during radiography, 135–137, 136f–137f

Respiratory distress, in scoliosis, 221

Respiratory infection(s), upper, and anesthesia, 110

Respiratory muscle(s), paralysis of, in poliomyelitis, 1323

Retardation, mental. *See* Mental retardation.

Retinacular artery(ies), occlusion of, in Legg-Calvé-Perthes disease, 668, 670f

Retinacular cyst(s), of upper extremities, 496–497

Retractive fibrotic tissue, and talipes equinovarus, 923

Revascularization
 in Legg-Calvé-Perthes disease, classification of, 681–682, 683f–684f
 microsurgery for, 504

Reverse transcriptase-polymerase chain reaction (RT-PCR), in Ewing's sarcoma, 2033

RGO(s) (reciprocating gait orthoses), for myelomeningocele, 1294–1295, 1295f

Rh blood group incompatibility, and cerebral palsy, 1123

Rhabdomyosarcoma, 2040–2042, 2041t

Rheumatoid arthritis, 467–476, 468f–476f, 1813–1821, 1815f–1819f, 1817t and torticollis, 194 back pain in, 103–104 classification of, 1813 clinical features of, 1816f–1817f, 1816–1819, 1817t in pauciarticular type, 1816f, 1816–1817, 1817t in polyarticular type, 1817, 1817f–1818f in systemic type, 1817–1819, 1818f definition of, 1813 etiology of, 1813, 1815 incidence of, 467, 1813 laboratory studies in, 1819 of hand, 1813–1817, 1817t treatment of, 473–474, 474f–475f pathology of, 1815–1816, 1815f–1816f patient/parent education in, 468, 468f

Rheumatoid arthritis (*Continued*)
 radiographic features of, 1819, 1819f treatment of, 1819–1821 intra-articular corticosteroid injections in, 469 multidisciplinary collaboration in, 467 physical therapy in, 468–469 role of hand surgeon in, 468 splinting in, 468–469 total joint arthroplasty in, 1821 Rheumatoid factor, 1817 Rhizomelic chondrodysplasia punctata, 1544f–1546f, 1545–1546 Rhizomelic micromelia, in achondroplasia, 1510 Rhizotomy
 for cerebral palsy, 1231–1232, 1232f–1233f lumbar hyperlordosis after, 309, 1232, 1232f Rhomboid muscle(s), dissection of, in forequarter amputation, 2004p–2005p Rib(s)
 deformities of, in scoliosis, 220, 239 fusion of, in congenital scoliosis, 286f resection of, in anterior instrumentation, for thoracolumbar scoliosis, 264p–265p strut graft of, for thoracic kyphosis, 148f Rib graft(s)
 for postirradiation kyphosis, 343f, 345 strut, for congenital kyphosis, 295, 295f–297f Rib-vertebral angle difference (RVAD), in idiopathic scoliosis infantile, 276, 276f juvenile, 280 Rickets, 129, 129f drug-induced, 1690 genu valgum from, 861, 863 hypophosphatemic tumor-related, 1695 nutritional, 1686–1689, 1686f–1689f clinical features of, 1688–1689 laboratory studies in, 1687f, 1687t, 1688 pathology of, 1686f, 1686–1687 radiographic features of, 1687f–1689f, 1689 treatment of, 1689 of prematurity, 1690 vitamin D-resistant, 1690–1695 clinical features of, 1691f, 1693 genetic basis of, 1693 laboratory studies in, 1690 medical treatment of, 1693–1694 orthopaedic management of, 1692f–1694f, 1694–1695 radiographic features of, 1691f–1692f, 1693 Rifampin, for tuberculous arthritis, 1831 Riley-Day syndrome, 1449t, 1453 anesthesia in, 124t Ring fixator, Ilizarov, for angular deformities of leg, 1116f, 1117 Ring splint(s), Oval-8, 393, 394f Risser sign, 215, 215f, 216 grades of, and indications for treatment, 228 Rocker-bottom foot, 959, 960f–961f. *See also* Talus, vertical Rockwood classification, of acromioclavicular joint injuries, 2115, 2117f Rod(s). *See also* Harrington rod(s). placement of, in anterior instrumentation, 274p–275p Rolyan Elastomer, for total contact molding inserts, 390, 390f Rooting reflex, 55–56

Rotation. *See also under specific joint, e.g., Hip.*
 definition of, 27t Rotation flap, for arthrogryposis, 444, 447f Rotation reflex, 52–53 Rotational maneuver, in Texas Scottish Rite Hospital instrumentation, 254p–255p Rotational osteotomy, of femur or tibia, for cerebral palsy, in knee, 1169–1173, 1178f Rotationplasty
 for femoral osteosarcoma, 2026, 2027f for proximal focal femoral deficiency, 1763, 1766f–1767f Rotator cuff
 lateral, lengthening of, for brachial plexus palsy, 108, 118f teres major/latissimus dorsi muscle transfer to, for brachial plexus palsy, 485, 486f, 488–491p Rotterdam straight line graph(s), for timing of epiphysiodesis, in leg length inequality, 1060f Round cell(s), in Ewing's sarcoma, 2032, 2032f Round-back deformity. *See* Kyphosis. Roux-Goldthwait procedure, for recurrent patellar dislocation, 797–798 RT-PCR (reverse transcriptase-polymerase chain reaction), in Ewing's sarcoma, 2033 Rubinstein-Taybi syndrome, 1665, 1665f “Rugger jersey” spine
 in osteopetrosis, 1551, 1552f in renal osteodystrophy, 1697 Rule of thirds, 2092, 2094f RVAD. *See* Rib-vertebral angle difference (RVAD). Ryanodine receptor, and malignant hyperthermia, 118

“Saber shin,” in congenital syphilis, 1869, 1870f SAC. *See* Space available for cord (SAC). SACH (solid ankle-cushioned heel) prosthetic foot, for leg length inequality, 1058, 1058f Sacral agenesis. *See* Caudal regression syndrome. Sacrofemoral angle, 1173, 1179f Sacroiliac joint
 injury to, in pelvic fractures, 2264, 2264f internal fixation of, 2267–2268, 2268f–2269f septic arthritis of, 1863 Sacroplasty, for reduction of spondylolisthesis, 368f, 368–369 Sacrum
 aneurysmal bone cyst of, 105f fractures of, 2261 Safety cast, 384, 387f after treatment of syndactyly, 442, 443f “Sagging rope” sign, in Legg-Calvé-Perthes disease, 678, 679f Sagittal balance, in Scheuermann's kyphosis, 326, 327f Saha trapezius transfer, for deltoid paralysis, 1406f, 1406t, 1406–1407 Salmonella osteoarticular infection(s), in sickle cell disease, 1864 Salter innominate osteotomy
 bone graft for, 638p–641p dissection in, 636p–637p for avascular necrosis, 596, 618–621, 621f, 630f–633f, 631, 634p–641p complications of, 621 incision for, 619 indications for, 619 Kalamchi modification of, 631, 633f

Salter innominate osteotomy (*Continued*)
 principles of, 618, 621f
 results of, 621, 631
 for concentric hip reduction, 618
 for Legg-Calvé-Perthes disease, 698–699, 701f–702f
 incision in, 634p–635p
 placement of Gigli saw for, 636p–639p
 postoperative care in, 640p
 Salter-Harris fracture classification, 2062f, 2062–2063
 of ankle fractures, 2396–2402, 2400f–2403f
 of distal femoral fractures, 2322–2323, 2322f–2324f
 of lateral condylar humeral fractures, 2177–2178
 of proximal tibial phyeal fractures, 2345–2346, 2345f–2346f
 Sanders classification, of calcaneal fractures, 2425
 Sandifer's syndrome, and torticollis, 194
 Sanfilippo syndrome, 1573t, 1577
 anesthesia in, 124t
 Saphenous nerve, dissection of, in midthigh amputation, 1984p–1985p
 Saphenous vein, ligation of, in open reduction of hip, medial approach to, 550p–551p
 Sarcoma
Ewing's. See Ewing's sarcoma.
osteogenic. See Osteosarcoma.
 soft tissue, 2040–2046, 2041t, 2043f–2045f
 nonrhabdomyosarcoma, 2042, 2043f
 in older children, 2042–2046, 2044f–2045f
 Sartorius muscle, dissection of, in Pemberton osteotomy, 622p–623p
 Saturn phenomenon, and prognosis in Legg-Calvé-Perthes disease, 682f, 687, 689
 Sausage dactylitis, in psoriatic arthritis, 1822, 1823f
 Sauve-Kapandji technique, for brachial plexus palsy, with elbow deformities, 487
 Scanner(s), for computed tomography, 138–139, 138f–141f
 Scanography, in leg length inequality, 1043, 1046f–1047f
 Scaphoid, fractures of, 2246–2248, 2247f–2249f
 Scapula
 anatomy of, 2121, 2121f
 dissection of muscles from, in forequarter amputation, 2002p–2007p
 fractures of, 2121–2123, 2121f–2123f
 anatomic considerations in, 2121, 2121f–2122f
 complications of, 2122
 diagnosis of, 2122, 2123f
 mechanism of, 2121
 scapulothoracic dissociation with, 2122–2123
 treatment of, 2122
 with os acromiale, 2123
 winging of, scapulocostal stabilization for, 1481, 1482p–1483p
 Scapulocostal stabilization, for facioscapulohumeral muscular dystrophy, 1481, 1482p–1483p
 Scapuloperoneal dystrophy, 1481
 Scapulothoracic dissociation, with scapular fractures, 2122–2123
 Scapulothoracic joint, range of motion of, 28f, 29–30, 30f

Scar(s)
 hypertrophic, of hand, 382, 383f
 total contact molding inserts for, 390f–391f, 390–392
 SCFE. *See Slipped capital femoral epiphysis (SCFE).*
 Scham's sign, in slipped capital femoral epiphysis, 719, 720f
 Schanz screw(s), placement of, in reduction of hip dislocation, 2280, 2281f
 Scheie's syndrome, 1573t, 1580
 anesthesia in, 124t
 Scheuermann's kyphosis, 101f, 101–102, 324–336, 326f–338f, 328t
 clinical features of, 325
 diagnosis of, 324–325, 328f
 etiology of, 325
 incidence of, 323–324
 location of, 323–324, 326f
 lumbar, 102, 102f
 nonoperative treatment of, 328–334, 329f–330f
 bracing in, 328, 329f
 casts in, 329, 330f
 operative treatment of, 329–333, 331f–332f
 approach to, 330–331
 thoracotomy, 333
 compression rods for, 331, 335
 excessive correction in, 336, 337f
 goals of, 330, 331f
 Harrington compression rods in, 331, 335
 hook-rod systems in, 334–335, 335f
 indications for, 329
 laminar resection in, 336
 Luque instrumentation in, 334
 maintenance of correction in, 336, 338f
 preoperative evaluation in, 331
 spinal fusion in
 choice of level for, 331–333, 332f
 level of, 331–333, 332f
 video-assisted, 333–334
 with anterior release, 333
 thoracoscopic, 333–334
 with instrumentation, 327f, 331, 333, 335
 with osteoperiosteal flap, 333
 with posterior instrumentation, 334–336, 335f–338f
 with thoracoscopic anterior release, 333–334
 with thoracoscopic diskectomy, 334
 pathogenesis of, 324, 327f
 radiographic features of, 324f, 326–328, 328t
 spondylolysis in, 326
 treatment of, 325–326
 vs. postural kyphosis, 323, 324f
 Schmid metaphyseal chondrodyplasia, 1547–1548, 1548f–1549f
 Schmidt and Weiner classification, of calcaneal fractures, 2424, 2424f
 Schmorl's node(s), in Scheuermann's kyphosis, 324, 328f
 School(s), scoliosis screening in, 216–217
 Schwannoma
 malignant, 2045f
 of upper extremities, 503
 Sciatic nerve
 dissection of
 in hip disarticulation, 1976p–1977p
 in midthigh amputation, 1986p–1987p
 injury to
 from hip dislocation, 2281
 from Salter innominate osteotomy, 621

Sciatic nerve (*Continued*)
 palsy of, 1455–1456
 protection of
 during proximal hamstring release, for hip subluxation/dislocation, 1190p–1191p
 in Pemberton osteotomy, 624p–625p
 Sciatica, with lumbar disk herniation, treatment of, 373
 Scintigraphy. *See Bone scan(s).*
 Scissoring, in cerebral palsy, 1127, 1127f
 from adduction contractures of hip, 1173, 1179f
 Cllerae, blue, in osteogenesis imperfecta, 1724–1725
 Scleroderma, anesthesia in, 124t
 Sclziometer, 217f, 217–218
 Scoliosis, 213–312
 after cervical spine fractures, 2098–2099
 after laminectomy, 308–312, 310f–311f
 after spinal cord injury, 2107–2108, 2109f
 classification of, 213, 214t
 computed tomography in, 227–228
 congenital, 283–291, 283f–292f
 cervicothoracic, 285, 286f
 classification of, 283, 283f
 magnetic resonance imaging in, 287, 288f
 natural history of, 285, 286f, 287
 progression of curve in, 287, 288f–289f
 radiographic evaluation of, 287, 288f–289f
 treatment of
 nonoperative, 287
 operative, 287–291, 289f–292f
 with heart disease, 308, 309f
 curve in
 cerebral palsy and, 1225, 1225f–1226f
 magnitude of
 and indications for treatment, 228, 228t, 233–234
 measurement of, 225, 225f–226f
 pattern of, and preoperative planning, 234t, 234–235, 234f–235f
 terminology related to, 214t
 definition of, 213, 214f
 from radiation therapy, 312
 from thoracotomy, 308
 genetic factors in, 218–219, 219f
 hysterical, 312
 idiopathic, 213–261
 adolescent, 213–261, 217f, 219f–227f, 230f–231f, 233f–243f, 244p–257p, 258f–261f
 and assessment of growth potential, 239–240, 239f–240f
 back pain in, 102
 connective tissue disorders and, 218
 curve in
 and fusion level, 235–236, 236f–238f
 stable vertebra and, 234–235, 235f
 diagnosis of
 peak ages for, 213
 plumb line in, 222, 222f
 genetic factors in, 218–219, 219f
 imaging studies in, 224–228, 225f–227f
 computed tomography, 227–228
 magnetic resonance, 227
 plain radiography, 224–226, 224f–227f
 surface, 226, 227f
 infantile, 276f–279f, 276–280
 abnormalities associated with, 277
 clinical features of, 276, 276f
 natural history of, 276
 progression of curve in, 276–277, 276f–277f

Scoliosis (*Continued*)
 resolution of curve in, 277
 treatment of
 nonoperative, 277
 operative, 277–280, 278f–279f
 juvenile, 280, 281f
 mortality rate in, 216
 natural history of, 214–216, 215f, 216t
 after skeletal maturity, 216
 before skeletal maturity, 214–216, 215f, 216t
 curve magnitude and, 216, 216t
 curve progression and, 214
 gender and, 215
 peak height velocity and, 215f, 215–216
 remaining growth and, 215f, 215–216
 neurologic disorders and, 218
 pathophysiology of, 219–220, 220f
 patient evaluation in, 221–224, 221f–224f
 neurologic examination and, 223–224, 223f–224f
 physical examination and, 221–222, 221f–223f
 presenting complaint and, 221
 prevalence of, 213–214
 screening for, 216–218, 217f
 in school programs, 216–217
 methods of, 217f, 217–218
 treatment of
 and postoperative pain management, 241
 anterior instrumentation in, 259–261, 259f–261f, 262p–275p
 Dwyer, 259, 259f
 for thoracic deformity, 261, 261f
 for thoracolumbar/lumbar curves, 262p–275p
 solid rod, 260–261, 260f–261f
 Zielke derotation spondylodesis, 259–260, 260f
 bone grafting in, 240–241
 determination of transfusion requirements for, 240
 nonsurgical, 228–232, 230f–233f. *See also* Brace(s), for idiopathic scoliosis.
 biofeedback in, 232
 electrical stimulation in, 232
 physical therapy in, 232
 observation as, 228
 posterior instrumentation in, 241–243, 241f–243f, 244p–257p, 258f
 complications of, 258–259
 Cotrel-Dubousset, 242–243, 243f
 Harrington, 241f, 241–242
 Isola, 243, 258f
 Luque double L-rod segmented, 242, 242f, 244p–245p
 new methods of, 259
 Texas Scottish Rite Hospital, 243, 243f, 246p–257p
 Wisconsin segmental spinal, 242, 242f, 244p–245p
 principles of, 228, 228t
 spinal cord monitoring during, 241
 surgical, 232–241, 234f–240f, 235t
 goals of, 232
 indications for, 233–234
 instrumentation systems for, 232–233
 preoperative planning for, 233–234, 234t, 234f–240f

Scoliosis (*Continued*)
 video-assisted thorascopic surgery in, 240
 in arthrogryposis, 1660
 in Beals' syndrome, 1595f
 in cerebral palsy, 1225–1229, 1229f–1234f
 curve in, 1225, 1225f–1226f
 incidence of, 1225
 nonoperative treatment of, 1225
 pelvic obliquity and, 1225, 1225f–1226f
 progression of, 107
 spinal fusion for, 1225–1231, 1228f–1230f
 anterior, 1227, 1230
 complications of, 1230f, 1230–1231
 Dunn-McCarthy technique of, 1227, 1229f
 Galveston technique of, 1227, 1230f
 Luque technique of, 1226–1227
 in Charcot-Marie-Tooth disease, 1441, 1445f–1446f
 in cleidocranial dysostosis, 1569f, 1570
 in diastrophic dysplasia, 1536, 1537f
 progression of, 1540f, 1541
 in Duchenne muscular dystrophy, surgical treatment of, 1470–1471, 1476f–1477f
 in fibrodysplasia ossificans progressiva, 1634, 1637f
 in Larsen's syndrome, treatment of, 1611–1614, 1613f–1614f
 in Marfan's syndrome, 1590. *See also* Marfan's syndrome, scoliosis in.
 in muscular dystrophy, Emery-Dreifuss, 1479
 in myelomeningocele, 1278f, 1280–1281
 spinal fusion for, 1281, 1288–1289
 in neurofibromatosis, 299–302, 299f–305f, 1628, 1629f. *See also* Neurofibromatosis, and scoliosis.
 in osteogenesis imperfecta, 1738–1739, 1739f
 in spinal muscular atrophy, 1314, 1316f–1318f, 1317
 in spondyloepiphyseal dysplasia congenita, 1529
 in spondylolisthesis, 97f
 magnetic resonance imaging in, 227
 Marfan's syndrome and, 302–308, 306f–307f
 operative management of, blood loss during, 116
 physical examination in, 69, 70f
 screening for, 64
 with hip subluxation/dislocation, in cerebral palsy, 1187, 1188f
 “Scotty dog” sign
 in spondylolisthesis, 354, 356f
 in spondylolysis, 100, 100f
 Screening examination, 63–65, 64f
 Screw(s)
 placement of, in anterior instrumentation, for idiopathic scoliosis, 272p–273p
 transphyseal, in epiphyseal stapling, for leg length inequality, 1063, 1074f
 Scurvy, 1705–1707, 1706f
 Season, and incidence of slipped capital femoral epiphysis, 712
 Seat belt injury(ies), classification of, 2103, 2107f
 Secondary sexual characteristics, development of, 19, 19f
 SED. *See* Spondyloepiphyseal dysplasia (SED).
 Sedation
 by nonanesthesiologist, 111, 111t
 for computed tomography, 142–143, 145
 Segmental spinal dysgenesis, 297
 Seizure(s), medications for, and rickets, 1690

Semimembranous tendon, lengthening of, for iliotibial band contractures, 1334p–1335p
 Semitendinosus muscle
 dissection of, in hamstring lengthening, for iliotibial band contractures, 1334p–1335p
 transfer of, for quadriceps femoris paralysis, 1351, 1356–1358, 1357f
 Semitenodesis
 dissection for, 800p–803p
 for recurrent patellar dislocation, 797, 800p–805p
 dissection in, 800p–803p
 postoperative care after, 804p–805p
 Sensibility, in hand, and reconstructive surgery, 404f, 404–405
 Sensory impairment, in upper extremity, cerebral palsy and, 1209
 Sensory neuropathy, congenital, 1449t, 1453
 Septic arthritis. *See Arthritis, septic.*
 Sequestrum, after limb lengthening, 1090–1091
 Serpentine foot, for pes planovalgus, 921f, 921–922
 Serratus anterior muscle, dissection of, in forequarter amputation, 2006p–2007p
 Serratus anterior tendon transfer, for subscapularis muscle paralysis, in poliomyelitis, 1407, 1407f
 Severed extremity(ies), replantation of, 503–504
 Severin classification, in developmental dysplasia of hip, 532, 533t
 Sex. *See Gender.*
 Sexual characteristics, secondary, development of, 19, 19f
 Sexual precocity, in fibrous dysplasia, 1911
 Shapiro classification, of osteogenesis imperfecta, 1721, 1721t
 Sharrard procedure, 1273–1274, 1275f
 Shelf acetabular augmentation, 645f–649f, 646
 for subluxation/dislocation of hip, in cerebral palsy, 1198, 1200p–1201p
 Shelf arthroplasty, for Legg-Calvé-Perthes disease, 700–701
 Shh (sonic hedgehog gene), in limb development, 1745, 1747
 “Ship's funnel” thigh, in proximal focal femoral deficiency, 1761f
 Shoe insert(s), for pes planovalgus, 914, 914f
 Shoe lift(s), for leg length inequality, 1058, 1061f
 Shoes, orthopaedic, for pes planovalgus, 912
 Short stature
 classification of, 1505
 in achondroplasia, 1517, 1519, 1519f
 in cleidocranial dysostosis, 1569
 in hypochondroplasia, 1521
 Short-limb gait, 84
 Short-tau inversion recovery (STIR) magnetic resonance imaging, 153
 Shoulder
 abduction contracture of, with external rotation, 108, 118f–119f
 abduction of, 29f, 31, 32f
 adduction contracture of, with internal rotation, 106–108, 108f, 110p–117p
 adduction of, 29f, 31, 32f
 anatomy of, 2128–2129, 2129f
 arthrodesis of, in poliomyelitis, 1407–1408
 deformities of, from brachial plexus palsy, 481, 482f–483f, 483t
 management of, 484–486, 488p–495p, 496f–497f
 disarticulation of, 2014p–2017p

- Shoulder (*Continued*)

elevation of, 29, 30f, 31, 32f

extension of, 29f, 29–30

flexion of, 29, 29f

in poliomyelitis, 1395, 1406t, 1406–1408, 1407f–1409f. *See also* Poliomyelitis, shoulder in.

instability of, after glenohumeral joint dislocation, 2131

muscles of

depressor group, 1406–1407, 1407f–1409f

innervation of, 47t

prime movers, 1395

steering group, 1395, 1406

range of motion of, 29–31, 29f–32f

rotation of, 30–31, 31f–32f

after lateral rotation osteotomy of humerus, 494p–495p

synovial chondromatosis of, 1946f

traumatic dislocation of. *See* Glenohumeral joint, traumatic dislocation of.

Shunting, ventriculoperitoneal, in myelomeningocele, 1257

Sickle cell disease, 1892f–1893f, 1892–1894

avascular necrosis in, 1894

bone changes in, 1894

bone infarction in, 1892–1893

hand-foot syndrome in, 1892f–1893f, 1893

osteomyelitis in, 1863–1864, 1893

pathophysiology of, 1892

tourniquets and, 1894

vertebral involvement in, 1893f, 1893–1894

Signal intensity, in magnetic resonance imaging, 153t, 153–155, 154f, 156f

Silicone sheet(s), for total contact molding inserts, 390, 391f

Silipos sleeve, for securing total contact molding inserts, 392, 392f

Sillence classification, of osteogenesis imperfecta, 1718–1719, 1719t

Silver-Russell syndrome, 1039, 1041f

Silver's syndrome, anesthesia in, 124t

Silverskiöld test, 1135, 1135f

Single-photon emission computed tomography (SPECT), 167

in differential diagnosis of back pain, 98

in spondylolysis, 358

Sitting balance, in spinal muscular atrophy, 1316f–1317f

Skeletal anomaly(ies)

in homocysturia, 1597

in Marfan's syndrome, 1587, 1587f

in neurofibromatosis, 1628, 1629f–1631f

Skeletal dysplasia. *See* Bone dysplasia.

Skeletal growth, normal, and prediction of leg length inequality, 1045–1048, 1047f–1048f

Skeletal muscle, periodic paralysis of, 1489–1491

“Skeletal standstill,” in Legg-Calvé-Perthes disease, 661–662, 664f

Skeletal traction, for femoral shaft fractures, 2307f, 2307–2308

Skeleton

immature, 2059–2072, 2060f–2072f, 2070t

buckle fractures of, 2059–2060, 2060f–2061f

greenstick fractures of, 2060, 2061f

physeal injuries to, 2062–2072, 2063f–2072f, 2070t. *See also* Physis (physes).

plastic deformation of, 2059, 2060f

remodeling/overgrowth in, 2060–2062, 2062f

Skeleton (*Continued*)

normal radiographic variants of, 130–132, 130f–133f

Skewfoot, 921f, 921–922

Skin

care of, in myelomeningocele, 1295–1296

hyperlaxity of, in Ehlers-Danlos syndrome, 1639, 1639f

in myelomeningocele, 1252, 1252f

lesions of

in neurofibromatosis, 1625, 1626f–1628f

in spinal dysraphism, 1302

of hand, protection of, dressings for, 386, 386f–387f

Skin distraction, in measurement of range of motion, of thoracolumbar spine, 35

Skin graft(s)

for coverage after amputation, 1796, 1796f

for cutaneous syndactyly, 436, 436f

attachment of, 440, 441f

harvest of, 440, 440f

patterns of, 439, 440f

Skin traction, for femoral shaft fractures, 2305, 2307f, 2307–2308

Skip metastases, from osteosarcoma, 1955–1956

Skull

in osteopetrosis, 1552

wormian bones of, in osteogenesis imperfecta, 1725–1726, 1727f

“Skyline view,” in recurrent patellar dislocation, 794, 796, 796f

Sleep apnea, in achondroplasia, 1514

Sliding calcaneal osteotomy, for pes planovalgus, 916, 919f–920f

Sling(s), Snyder, for Legg-Calvé-Perthes disease, 659f

“Slings and springs,” for traction, in Legg-Calvé-Perthes disease, 692, 695f

Slip ring technology, in computed tomography, 139, 140f–141f

Slipped capital femoral epiphysis (SCFE), 711–759

acute, 712–713, 713f

acute-on-chronic, 713f, 714

age at onset of, 712

bilateral, incidence of, 712

chronic, 713, 713f

classification of, 712–714, 713f–714f, 745

clinical features of, 718f, 718–719

complications of, 747–753, 749f–755f, 756p–757p

avascular necrosis as, 744–745, 750–753, 750f–755f, 756p–757p. *See also* Avascular necrosis (AVN).

chondrolysis as, 747–750, 749f. *See also* Chondrolysis.

diagnosis of, radiography in, 129, 129f

epidemiology of, 711–712

etiology of, 715–716

endocrine, 715–716

mechanical, 715

historical perspective on, 711

imaging studies of, 719–721, 719f–721f

in hypothyroidism, 1713, 1714f

in renal osteodystrophy, 1697f–1700f, 1699

incidence of, 711–712

limping from, 90–91, 801f

magnetic resonance imaging in, 721

pathology of, 716–718, 717f

pathophysiology of, 711

stable, vs. unstable, 714

Slipped capital femoral epiphysis (SCFE) (*Continued*)

treatment of, 722–747, 723f, 724p–743p

bone graft epiphysiodesis in, 731, 732p–735p, 736

in anterior slips, 745

in situ pinning in, 722–723, 724p–727p, 728–731, 730f

open, using radiolucent table, 723, 728f, 728–729

percutaneous, using fracture table, 722–723, 723f, 724p–727p

in unstable cases, 744–745

in valgus slips, 745

initial, 722

long-term outcome after, 753, 755, 758–759

primary osteotomy in, 736–739, 737f, 739f, 740p–741p

base-of-neck, 737f, 737–738, 742p–743p

Dunn procedure for, 736–737, 737f, 740p–741p

goals of, 736

Imhauser, 736, 737f, 738

intertrochanteric, 738–739, 739f

Kramer and Barmada, 737f, 737–738, 742p–743p

locations for, 736, 737f

Southwick, 738–739, 739f

prophylactic, of contralateral hip, 745, 747, 748f

remodeling after, 758

residual deformity after, 745, 746f

spica cast in, 739, 744

untreated, long-term outcome in, 753

with screw impingement on acetabulum, 146f

Slipped vertebral apophysis, 99, 374–375

Slipper prosthesis, for congenital foot absence, 1783f

SMA. *See* Spinal muscular atrophy (SMA).

Smith-Lemli-Optiz syndrome, anesthesia in, 124t

Smith-Peterson approach, to bone graft epiphysiodesis, for slipped capital femoral epiphysis, 731p, 732p–733p

Snapping iliotibial band syndrome, 787

Snapping knee, congenital, 825, 826f

Snyder sling, for Legg-Calvé-Perthes disease, 659f

Social skill(s), developmental milestones for, 4f, 21, 21t

Socioeconomic status, and Legg-Calvé-Perthes disease, 665

Sofield procedure, for long bone deformities, in osteogenesis imperfecta, 1732f, 1732–1733

Soft tissue

bleeding in, from hemophilia, 1881–1883, 1883f

calcification of, in hypervitaminosis D, 1704, 1705f

examination of, and radiographic diagnosis, 129

hypertrophy of, in Klippel-Trénaunay syndrome, 1671–1672

injuries to, from child abuse, 2079

magnetic resonance imaging of, 159

sarcoma of, 2040–2046, 2041t, 2043f–2045f

nonrhabdomyosarcoma, 2042, 2043f

in older children, 2042–2046, 2044f–2045f

“starburst” reaction in, in osteosarcoma, 2027f

Soft tissue tumor(s), malignant, of upper extremities, 502–503
 Softcast, 386
 Solid ankle-cushioned heel (SACH) prosthetic foot, for leg length inequality, 1058, 1058f
 Somatomedin C, and Legg-Calvé-Perthes disease, 662
 Somatosensory evoked potential(s) (SSEPs), during scoliosis surgery, 241
 Sonic hedgehog gene (*Shh*), in limb development, 1745, 1747
 Sotos' syndrome, anesthesia in, 125t
 Southwick measurement, of femoral head-neck angle, 714, 714f
 Southwick osteotomy, for slipped capital femoral epiphysis, 738–739, 739f
 Space available for cord (SAC), measurement of
 in atlantoaxial rotatory displacement, 188f
 in Klippel-Feil syndrome, 177–178, 181f–182f
 Spastic diplegia
 and hallux valgus, 1013f
 gait in, 81f
 Spastic gait, 84
 Spastic hemiplegia, with equinovarus deformity, 1142, 1142f
 Spastic thumb, 446
 Spasticity
 assessment of, 27
 in cerebral palsy, 1123
 medical treatment of, 1132–1134
 Spatial resolution, in computed tomography, 140, 142f
 Special Olympics, qualification for, Down syndrome and, 1618
 SPECT. See Single-photon emission computed tomography (SPECT).
 Speech, developmental milestones in, 21
 Sphenoid bone, synchondrosis of, 130, 130f
 Spica cast(s)
 after intertrochanteric varus osteotomy, with internal blade plate fixation, 576p
 for femoral fractures, 2308, 2308f–2309f
 for femoral shaft fractures
 ambulation in, 2314, 2315f
 in child 7 months to 5 years, 2304, 2306f
 for hip fractures, 2291–2301
 for slipped capital femoral epiphysis, 739, 744
 Spina bifida. *See* Myelomeningocele.
 Spina bifida occulta, 1249, 1305, 1306f
 of cervical spine, in diastrophic dysplasia, 1536, 1537f
 Spinal canal
 deformity of, in scoliosis, 220, 220f
 osteochondroma of, computed tomography in, 148f
 Spinal cord
 anterior horn cells of, in poliomyelitis, 1322
 blood flow to, induced hypotension and, 116
 decompression of, 2106
 in spondylolisthesis, 362–363, 364f
 disorders of, 1249–1319. *See also under specific disorder, e.g., Myelomeningocele.*
 differential diagnosis of, magnetic resonance imaging in, 158–159, 161f
 injuries to, 2087–2110. *See also Cervical spine, fractures of; Lumbar spine, injuries to; Thoracic spine, injuries to.*
 monitoring of, during scoliosis surgery, 241

Spinal cord (*Continued*)
 neurogenic abnormalities of, and pes cavus, 984, 984f, 987–988
 pathology of, in myelomeningocele, 1252
 tethered, 1303, 1304f
 back pain in, 102
 in myelomeningocele, 1258, 1258f
 magnetic resonance imaging in, 161f
 tumors of, back pain from, 105
 Spinal deformity(ies), assessment of, 68–70, 70f
 Spinal dysgenesis, segmental, 1509
 Spinal dysraphism. *See also under specific type, e.g., Myelomeningocele.*
 cutaneous lesions in, 1302
 musculoskeletal disorders in, 1302
 neurologic disorders in, 1302
 with other congenital spinal anomalies, 281–282
 Spinal fusion
 cervical
 for atlantoaxial rotatory displacement, 189
 for cervical kyphosis, 199, 201f, 202
 for Klippel-Feil syndrome, 178f–179f, 183, 185–187, 186f–187f
 for neurofibromatosis, 198f
 for postlaminectomy kyphosis, 204, 205f, 206
 for cervical kyphosis, in diastrophic dwarfism, 1539f, 1541, 1541f
 for cervical spine instability, in Down syndrome, 1619–1621, 1620f
 for kyphosis
 congenital, 294–295, 295f
 in myelomeningocele, Luque technique of, 1286p–1287p, 1288f
 postirradiation, 345–346, 346f
 thoracolumbar, in achondroplasia, 1515f, 1515–1516
 for lumbar spine injuries, 2105, 2106f
 for scoliosis
 congenital, 287–288
 Galveston technique of, 1227, 1230f
 in cerebral palsy, 1225–1231, 1228f–1230f
 Luque technique of, 1226
 in Larsen's syndrome, 1612, 1613f
 in Marfan's syndrome, 306, 307f, 308
 in myelomeningocele, 1281, 1288
 juvenile idiopathic, 280, 281f
 level of, 235–236, 236f–238f
 nondystrophic, 299, 300f
 with lumbar curve, 236, 237f
 with neurofibromatosis, 301, 304f–305f
 for spinal muscular atrophy, 1314, 1317, 1317f–1318f
 for spondylolisthesis, 362, 364–365
 high-grade, 363
 in situ, 363–369, 369f–370f
 for stenosis, in achondroplasia, 1515
 for thoracic spine injuries, 2105–2106, 2109f
 Luque instrumentation for
 in cerebral palsy, 1224–1225
 in myelomeningocele, 1286p–1287p, 1288f
 in osteogenesis imperfecta, 1738–1739, 1739f
 in spinal muscular atrophy, 1318f
 occiput-C5, for chondrodysplasia punctata, 197f
 posterior, for scoliosis, in Duchenne muscular dystrophy, 1471, 1476f–1477f
 prophylactic, after radiation therapy, 339–345
 with anterior diskectomy, for infantile idiopathic scoliosis, 277–278, 278f–279f

Spinal fusion (*Continued*)
 with instrumentation, for Scheuermann's kyphosis, 327f, 333
 Spinal hook(s), for spinal fusion, in idiopathic scoliosis, 237f–238f
 Spinal laminar line of Swischuk, 2089, 2089f
 "Spinal locomotor generator," 76
 Spinal muscular atrophy (SMA), 1311–1319, 1312t–1313t, 1312f–1318f
 classification of, 1311–1312, 1312f, 1312t
 diagnosis of, 1312
 differential diagnosis of, 1312, 1313t
 genetic studies of, 1311
 hip subluxation/dislocation in, 1314, 1314f–1315f
 scoliosis in, 1314, 1316f–1318f, 1317
 treatment of, 1312–1317, 1314f–1318f
 spinal fusion in, 1314, 1317f–1318f
 Spinal stenosis, in achondroplasia, 1516f, 1516–1517
 Spinal tumor(s), malignant, back pain from, 105
 Spine
 aneurysmal bone cyst in, 1906, 1907f
 cervical. *See Cervical spine.*
 congenital anomalies of, 280–298, 282f–298f
 abnormalities associated with, 281–283, 282f
 and dislocation, 297–298
 and kyphosis, 291–297, 292f–297f. *See also Kyphosis.*
 and scoliosis, 283–291, 286f–292f. *See also Scoliosis, congenital.*
 and segmental spinal dysgenesis, 297
 and vertebral displacement, 297, 298f
 etiology of, 280–281
 deformities of, in osteogenesis imperfecta, 1738–1739, 1739f
 dislocation of, congenital, 297–298
 fractures of, 99
 growth of, melatonin and, 218
 in cerebral palsy, 1225–1230, 1224f–1230f.
 See also Scoliosis, in cerebral palsy.
 interpedicular distance in, achondroplasia and, 1513, 1513f
 lumbar. *See Lumbar spine.*
 osteotomy of, for congenital scoliosis, 291, 292f
 thoracic. *See Thoracic spine.*
 torsion of, in scoliosis, 219–220, 220f
 tuberculosis of, 1831–1835, 1832f–1835f. *See also Spondylitis, tuberculous.*
 Spin-echo relaxation time (T2), in magnetic resonance imaging, 152, 152f
 Spin-lattice relaxation time (T1), in magnetic resonance imaging, 152, 152f
 Spinocerebellar ataxia, 1246–1248, 1247f
 Spinomuscular level, neuromuscular disorders at, and motor function, 1121, 1122t
 Spinous process(es), absence of, in scoliosis, 221, 221f
 Splenectomy, for Gaucher's disease, 1645
 Splint(s)
 for developmental dysplasia of hip, in young child ages 1–6 months, 537–538, 538f
 for elbow, 396–398, 398f
 for fingers, 392–393, 392f–394f
 for forearm, 396, 397f
 for hallux valgus, 1014, 1015f
 for hand, 387–398, 388f–398f. *See also Hand, splints for.*
 for metatarsal fractures, 2429, 2430f
 for osteogenesis imperfecta, 1730

- Splint(s) (Continued)**
- for poliomyelitis, 1327
 - for rheumatoid arthritis, 468–469
 - for supracondylar humeral fractures, 2147, 2149, 2150f
 - for wrist, 394, 395f–397f, 396
 - with tenosynovitis, 470
 - improper application of, 2079, 2080f
 - Split hand symbrachydactyly**, 1783, 1784f
 - Spondylitis**
 - ankylosing, 1821–1822, 1822f
 - back pain in, 103
 - infectious, 103
 - and torticollis, 193–194
 - tuberculous, 1831–1835, 1832f–1835f, 1865, 1866f
 - clinical features of, 1833, 1833f
 - paraplegia in, 1834–1835
 - pathophysiology of, 1831–1833, 1832f–1833f
 - radiographic features of, 1833–1834, 1834f–1835f
 - treatment of, 1834
 - Spondyloarthropathy, 1821–1823, 1822f–1823f, 1823t
 - Spondyloepiphyseal dysplasia (SED)**, 1527–1530, 1527f–1530f
 - congenital, 1527–1529, 1527f–1529f
 - clinical features of, 1527, 1527f
 - genetic features of, 1527
 - kyphosis in, 1508f
 - orthopaedic features of, 1527–1529, 1528f–1529f
 - radiographic features of, 1527–1528, 1528f
 - genu valgum from, 861, 865f
 - kyphosis in, 346–347
 - tarda, 1529–1530, 1530f
 - Spondylolisthesis, 351–369, 352f–368f
 - back pain in, 99–100
 - classification of, 351–354, 352f–353f
 - clinical features of, 354, 355f
 - computed tomography in, 354, 357f
 - diagnosis of, computed tomography in, 149f
 - dysplastic, 351–352, 352f
 - in Larsen's syndrome, 1613, 1614f
 - in scoliosis, 97f
 - from cerebral palsy, 1228
 - isthmic, 352, 353f
 - Meyerding grade in, 356, 358f
 - natural history of, 356
 - radiographic findings in, 354, 356, 356f–358f
 - range of impairment from, 351
 - treatment of, 358–369, 359f–370f
 - high-grade, 363
 - instrumented reduction in, 365–368, 368f
 - low-grade, 360–363, 362f–363f
 - age at onset and, 361
 - indications for, 361
 - positioning for, 362
 - spinal cord decompression in, 362–363, 364f
 - Wiltse muscle-splitting technique of, 362, 362f
 - with cauda equina encroachment, 361
 - prophylactic, 357
 - sacroplasty in, 368f, 368–369
 - spinal fusion in, 362, 364f–365f
 - in situ, 364–369, 366f–370f
 - with spondylolysis, 358–360, 359f–360f
 - Spondylolysis
 - back pain in, 99–100, 100f
 - diagnosis of, computed tomography in, 149f
 - in osteopetrosis, 1553f, 1553–1554
 - Spondylolisthesis (Continued)**
 - in pyknodysostosis, 1566f–1567f, 1567
 - in Scheuermann's kyphosis, 326
 - in scoliosis, from cerebral palsy, 1228
 - treatment of, 358–360, 359f–360f
 - Spondylomegaly**, of lumbar vertebrae, in Proteus syndrome, 1668f
 - Spondyloptosis**, reduction of, 366f, 368–369
 - Spongialization**, 793
 - Sporotrichosis**, 1870
 - Sports**. *See* Athletic activity(ies).
 - Sports injury(ies)**, Down syndrome and, 1618
 - Sprengel's deformity**, in Klippel-Feil syndrome, 174, 180f
 - SSEP(s) (somatosensory evoked potentials)**, during scoliosis surgery, 241
 - S-shaped foot**, for pes planovalgus, 921f, 921–922
 - Staheli shelf procedure**, for acetabular augmentation, 646
 - Staheli test**, 1173, 1179f
 - Stance phase**, 75, 76f
 - muscle activity during, 80–81
 - Standing frame(s)**, 1291, 1292f, 1295
 - Staphylococcus aureus infection(s)**, and osteomyelitis, 1845
 - "Starburst" soft tissue reaction, in osteosarcoma, 2027f
 - Starfish hand**, in achondroplasia, 1511, 1511f
 - Startle reflex**, 49, 1126
 - Static muscle strength**, 44
 - Stature**
 - in Marfan's syndrome, 1586–1587
 - prediction of, Anderson-Green-Messner charts for, 1048–1051, 1049f, 1050f–1055f, 1052t
 - short**
 - classification of, 1506
 - in achondroplasia, 1517, 1519, 1519f
 - in cleidocranial dysostosis, 1569
 - in hypochondroplasia, 1521
 - Steel hip arthrodesis**, for proximal focal femoral deficiency, 1768, 1771f
 - Steel osteotomy**, for acetabular repositioning, 632, 642f
 - Steel's rule of thirds**, 2092, 2094f
 - Steinberg sign**, in Marfan's syndrome, 1587, 1587f
 - Steindler flexorplasty**, 1408–1409, 1409f, 1410p–1417p
 - for elbow flexion, in arthrogryposis, 1655
 - pronation contractures after, 1418
 - Steinmann pin(s)**
 - insertion of
 - for tibial osteotomy, 1111, 1111f
 - in closing wedge valgus osteotomy, 614p–617p
 - in dorsal wedge osteotomy, for pes cavus, 1004p–1005p
 - in intertrochanteric double osteotomy, 612p–613p
 - in intertrochanteric varus osteotomy, and internal fixation with blade plate, 564p–575p
 - in Japas V-osteotomy, for pes cavus, 1008p–1009p
 - placement of, in bone graft epiphysiodesis, for slipped capital femoral epiphysis, 731, 732p–733p
 - Step length**, 75
 - Step time**, 75
 - Stepping reflex**, 49, 51, 53f
 - "Steri-strip stent" dressing, for hand, 386f
 - Sternoclavicular joint**
 - medial, anatomy of, 2115, 2116f
 - pseudodislocation of, 2118, 2120
 - Sternocleidomastoid muscle**
 - contractures of, in congenital torticollis, 173f–175f
 - release of, for congenital muscular torticollis, 172–173, 176f–177f
 - Sternocleidomastoid muscle transfer**, in poliomyelitis, 1418–1419, 1424f
 - Sternocleidomastoid tendon transfer**, for supraspinatus muscle paralysis, in poliomyelitis, 1407, 1409f
 - Steroid(s)**. *See* Corticosteroid(s).
 - Stevens-Johnson syndrome**, anesthesia in, 125t
 - Stewart-Milford classification**, of hip fracture-dislocations, 2273, 2274f
 - Stickler's syndrome**, 1593
 - vs. Marfan's syndrome, 1590
 - Stimson technique**
 - for reduction of glenohumeral dislocation, 2130, 2131f
 - for reduction of hip dislocation, 2275, 2279f
 - STIR (short-tau inversion recovery) magnetic resonance imaging**, 153
 - "Stork's leg" appearance, in Charcot-Marie-Tooth disease, 1435, 1436f
 - Straddle fracture(s)**, 2263, 2263f
 - Straight-leg-raising test**, 97
 - Strayer procedure**, for equinus deformity, in cerebral palsy, 1136, 1138f
 - Streeter band syndrome**. *See* Congenital band syndrome.
 - Streptococcal infection(s)**, and osteomyelitis, 1845
 - Stress**, psychological, limb lengthening and, 1091
 - Stress fracture(s)**
 - at L5, and isthmic spondylolisthesis, 352, 353f
 - in pyknodysostosis, 1566f, 1567
 - of hip, 2286, 2288f
 - treatment of, 2292–2293
 - of pars interarticularis
 - and isthmic spondylolisthesis, 352, 353f
 - nuclear imaging in, 166f
 - tibial, 2387–2389, 2388f
 - "Stress" Trendelenburg's test, 42, 43f
 - Stress-strain curve(s)**, for immature vs. mature bone, 2058t
 - Stretching exercise(s)**. *See* Exercise.
 - Stride length**, 75
 - Strut graft(s)**
 - for cervical kyphosis, in Larsen's syndrome, 202, 203f
 - for congenital kyphosis, 295, 295f–297f
 - for postirradiation kyphosis, 343f, 345
 - Stulberg classification**, of Legg-Calvé-Perthes disease, 687, 691f, 691t
 - and long-term prognosis, 702
 - Sturge-Weber syndrome**, anesthesia in, 125t
 - Subaxial instability**, 207, 209f, 210
 - Subchondral fracture(s)**
 - and severity of Legg-Calvé-Perthes disease, 656, 659f
 - in Legg-Calvé-Perthes disease, 699f
 - Subclavian artery**, dissection of, in forequarter amputation, 2010p–2011p
 - Subclavian artery supply disruption sequence hypothesis**, 1754–1755, 1754f–1756f, 1755t
 - Subclavian vein**, dissection of, in forequarter amputation, 2010p–2011p

- Subclavian vessel(s), laceration of, from clavicular injuries, 2121
- Subclavius muscle, dissection of, in forequarter amputation, 2008p–2009p
- Subcortical nuclei, and neonatal reflexes, 46
- Subcutaneous instrumentation, with spinal fusion, for infantile idiopathic scoliosis, 279–280
- Sublaminar hook, in Texas Scottish Rite Hospital instrumentation, 256p–257p
- Sublaminar wire(s), for stabilization of cervical spine, in Down syndrome, 1620, 1620f
- Subscapularis muscle
- and shoulder motion, 1395
 - paralysis of, in poliomyelitis
 - latissimus dorsi tendon transfer for, 1407, 1409f
 - pectoralis minor tendon transfer for, 1407, 1408f
 - serratus anterior tendon transfer for, 1407, 1407f
 - teres major tendon transfer for, 1407, 1409f
- Subscapularis tendon, lengthening of, for brachial plexus palsy, 484–485
- Subtalar arthrodesis, extra-articular, of foot and ankle, for poliomyelitis, 1388, 1396p–1399p
- Subtalar coalition, limping from, 92f
- Subtalar joint
- arthrodesis of, for pes valgus, in cerebral palsy, 1150
 - arthroereisis of, for pes planovalgus, 915, 915f
 - bony coalition of, computed tomography in, 143f
 - coalition of, 978f
 - in pes cavus, 989, 989f
 - motion of, in tarsal coalition, 974, 975f
 - release of, for talipes equinovarus, 937, 938f–939f
 - valgus deformity of, in myelomeningocele, 1268–1269
- Subungual exostosis, 1032, 1033f
- Sucking reflex, 55
- “Sunburst” appearance, in osteosarcoma, 1957, 1958f–1959f
- Supination
- assessment of, 31, 33f
 - definition of, 27
- Supination contracture(s), of forearm, in poliomyelitis, 1424–1427
- Suppurative myositis, 1496
- Supracondylar osteotomy
- for femoral anteverision, 786
 - of metacarpal joint, for radial polydactyly, 430, 431f
- Supramalleolar osteotomy
- for talipes equinovarus, with persistent toe-in gait, 950–951, 952f
 - for tibial torsion, 867, 867f
- Supraspinatus muscle
- and shoulder motion, 1395
 - paralysis of
 - levator scapulae transfer for, 1407, 1408f
 - sternocleidomastoid transfer for, 1407, 1409f
- Sural nerve, biopsy of, in Charcot-Marie-Tooth disease, 1436
- Suspension traction, after hamstring lengthening, 1336, 1337f
- Suture(s), for hand surgery, 384–385, 385f
- Swayback, in developmental dysplasia of hip, 89, 90f
- Swing phase, 75, 76f
- muscle activity during, 80–81
- Sybrachydactyly
- classification of, 1783, 1784f
 - treatment of, 1783–1785, 1784f–1785f
 - vs. congenital band syndrome, 461, 463f
- Syme amputation, with knee arthrodesis, for femoral deficiency, 1762f, 1762–1763, 1764p–1767p
- Syme prosthesis, for proximal focal femoral deficiency, 1794
- Symmetric tonic neck reflex, 54, 55f
- Symphysis pubis, open reduction/internal fixation of, 2266–2267, 2267f
- Synchondrosis
- ischio pubic, 130f, 131
 - vs. fractures, 130f, 130–131
 - vs. infections/neoplasms, 130f, 131
- Syndactyly, 433–444, 434f–447f, 435t
- classification of, 434
 - conditions associated with, 435, 435t
 - cutaneous, 434
 - treatment of, 436–442, 436f–444f
 - dressing application in, 441–442, 443f
 - flap closure in, 439, 439f
 - graft attachment in, 440, 441f
 - graft harvest and donor site closure in, 440, 440f
 - incision for, 437–438, 437f–439f
 - preparation for, 437
 - principles of, 436, 436f
 - release of tourniquet in, 440
 - separation of digits in, 438–439, 439f–440f
 - skin grafts for, 436, 436f
 - patterns of, 439, 440f
 - tip reconstruction in, 441, 441f
- Etiology of, 433
- “fenestrated,” 1786, 1787f
 - historical perspective on, 433
 - imaging studies of, 435
 - incidence of, 434–435, 435f
 - of thumb-index web, 442, 444, 445f–447f
 - of toes, 1023–1024, 1023f–1024f
 - pathology of, 433–434, 434f
 - treatment of
 - complications of, 444
 - timing of, 436
 - with central polydactyly, 423, 423f
- Synostosis, from radial/ulnar fractures, 2233
- Synovectomy
- for hemophilic arthropathy, 1888–1889
 - for rheumatoid arthritis, 1820
 - of hand, 474
 - for tuberculous arthritis, 1831
 - of wrist, 471
- Synovial chondromatosis, 1944, 1946f
- Synovial cyst(s), of upper extremity, 496–497
- Synovial fluid
- analysis of, 1811, 1812f, 1813, 1814t
 - in tuberculous arthritis, 1831
 - aspiration of, in septic arthritis, 1852
 - properties of, 1811
 - viscosity of, 1813, 1814t
- Synovial popliteal cyst(s), 830f, 831
- Synovial sarcoma, 2044f
- Synoviorthesis, for hemophilic arthropathy, 1889
- Synovitis
- and Legg-Calvé-Perthes disease, 665
 - in dermatomyositis/polymyositis, 1494
- Synovitis (*Continued*)
- of finger, 472–473, 473f
 - of hip, acute transient, 1823–1825
 - of thumb, 472
 - of wrist, 469, 469f
 - treatment of, 470–472, 470f–472f
 - exercise in, 470
 - intra-articular corticosteroid injections in, 470, 470f–471f
 - pigmented villonodular, 1944–1946, 1946f–1947f
 - toxic, 1823–1825
 - vs. septic arthritis, 1850
 - transient, limping from, 86, 87f
- Synovium, pathology of, in rheumatoid arthritis, 1815–1816, 1815f–1816f
- Syphilis, 1869, 1869f–1870f
- Syringomyelia, 1449t, 1454, 1454f
- after spinal cord injury, 2108
 - back pain in, 102
- Syrinx, diagnosis of, magnetic resonance imaging in, 161f
- T1 (relaxation time), longitudinal, in magnetic resonance imaging, 152, 152f
- T2 (relaxation time), transverse, in magnetic resonance imaging, 152, 152f
- Tachdjian brace, for Legg-Calvé-Perthes disease, 696, 697f
- Talectomy, for equinovarus, in myelomeningocele, 1264
- Talipes calcaneovalgus, 903–904, 907, 907f
- Talipes equinovarus, 922–957, 924f–957f
- classification of, 927, 930f–931f
 - diagnosis of, 927, 927f–930f
 - differential diagnosis of, 927
 - embryology of, 923
 - etiology of, 923
 - in arthrogryposis, 1653–1654
 - in diastrophic dysplasia, 1535, 1535f
 - in Freeman-Sheldon syndrome, 1663f
 - in myelomeningocele, 1263–1264
 - pathologic anatomy of, 923–927, 924f–926f
 - recurrence of, lateral column shortening for, 946, 948, 949f
 - residual hindfoot, 929, 935f
 - severity of, 930f–931f
 - treatment of
 - nonoperative, 927–934, 932f–934f
 - operative, 933–941, 936f–941f
 - Achilles tendon lengthening in, 931–932, 937, 938f
 - “cable cast” in, 937
 - calcaneal osteotomy in, 948, 950, 950f
 - calcaneocuboid release in, 935–936, 937f
 - complications of, 941–945, 943f–945f
 - current technique in, 937–941, 938f–941f
 - talonavicular pinning in, 935, 936f
 - tibiotalar release in, 936
 - timing of, 935
 - results of, 922
 - revision after, 945–946, 947f
 - with residual deformity, Dwyer technique for, 948, 950, 950f
 - with residual deformity, Ilizarov technique for, 951, 953, 954f–957f
- Talipes equinus
- and knee extension, 1356f
 - in arthrogryposis, 1649f
 - in myelomeningocele, 1263
 - Talocalcaneal angle, in vertical talus, 961, 965f

- Talocalcaneal bridge, fetal, and tarsal coalition, 967, 974f
- Talocalcaneal coalition, 978f
treatment of, 981, 981f
- Talocalcaneal ligament, release of, for talipes equinovarus, 939, 940f
- Talocalcaneocuboid coalition, 976f
- Talocalcaneonavicular joint, dorsolateral dislocation of, open reduction of, 966–967, 968p–973p
- Talonavicular joint
exposure of, in triple arthrodesis, 1390p–1391p
pinning of, for talipes equinovarus, 935, 936f
release of, for talipes equinovarus, 939, 939f
- Talonavicularocuneiform complex, medial imbrication of, for pes planovalgus, 916, 918f, 920f
- Talus
abnormalities of, in talipes equinovarus, 923–924, 924f–925f
anatomy of, 2416, 2417f
beaking of, in tarsal coalition, 980, 980f
blood supply to, 2416, 2418f
bony connection with navicular, and tarsal coalition, 967, 975f
fractures of, 2414–2420, 2417f–2420f
classification of, 2416, 2418f–2419f
complications of, 2420
diagnosis of, 2416–2417, 2419f
mechanism of, 2416
osteochondral, 2420–2423, 2421f–2422f
classification of, 2421, 2421f
diagnosis of, 2421–2422
mechanism of, 2420–2421
treatment of, 2422f, 2422–2423
treatment of, 2417–2420, 2420f
in lateral process fractures, 2419–2420
in talar body fractures, 2419
in talar neck fractures, 2417–2419, 2420f
vertical, 959–967, 960f–966f, 968p–973p
clinical features of, 959, 960f
conditions associated with, 959
congenital, 67–68
definition of, 959
differential diagnosis of, 961
etiology of, 959
in arthrogryposis, 1654
in myelomeningocele, 1265
pathologic anatomy of, 959–961, 961f–964f
radiographic features of, 961, 965f–966f
treatment of, 966–967, 968p–973p
complications of, 967
- Tanner system, for patient evaluation, in scoliosis, 224
- Tanner's stages of development, 19, 19f
- TAP (Tone and Positioning) splint, 396, 397f
- TAR (thrombocytopenia-absent radius) syndrome, anomalies in, 1752, 1753f
- Tarsal bone(s), ossification of, 892f
- Tarsal canal artery, anatomy of, 2416, 2418f
- Tarsal coalition, 68, 967, 974t, 974f–981f, 974–982, 976t
clinical features of, 974–976, 975f–977f, 976t
conditions associated with, 974, 976, 976f–977f
etiology of, 967, 974, 974f–975f
historical aspects of, 967
imaging studies of, 976, 978f–980f, 980
incidence of, 967, 974t
- Tarsal coalition (*Continued*)
limping from, 92, 92f
treatment of, 980–981, 981f
complications of, 982
results of, 981–982
- Tarsal navicular, osteochondrosis of, 899–900, 899f–900f
- Tarsal osteotomy, Japas-V, for pes cavus, 999, 1006p–1009p
- Tarsometatarsal joint
fractures of, 2425–2428, 2426f–2429f
anatomic considerations in, 2426
classification of, 2426–2427, 2427f
complications of, 2428
diagnosis of, 2427, 2428f
mechanism of, 2426, 2426f
treatment of, 2427–2428, 2429f
Heyman-Herndon release of, for metatarsus adductus, 903, 906f
- Tay-Sachs disease, anesthesia in, 125t
- T-condylar fracture(s), of humerus, 2202, 2204f
- TE (echo time), in magnetic resonance imaging, 152, 152t
- Technetium bone scan(s). *See* Bone scan(s).
- Teeth, abnormalities of, in osteogenesis imperfecta, 1725, 1725f
- Telangiectatic osteosarcoma, 1955
- Teleoroentgenography, in leg length inequality, 1043, 1045f
- Tendon sheath, giant cell tumors of, in upper extremities, 502
- Tendon transfer(s). *See also under specific tendon.*
for poliomyelitis, principles of, 1327–1328
for upper extremity weakness, in Charcot-Marie-Tooth disease, 1444–1445
in poliomyelitis, 1362t–1363t
in shoulder, for poliomyelitis, 1406t, 1406–1407, 1406f–1408f
- Tenosynovitis
of hand, pyogenic, 478, 478f
of wrist, 469–470
treatment of, 470–472, 470f–472f
exercise in, 470
intra-articular corticosteroid injections in, 470, 470f–471f
operative, 470–472, 472f
- Tenotomy, for congenital muscular torticollis, 172
- Tensile force, in developmental coxa vara, 769f
- Tension band wire(s), for patellar fractures, 2339, 2340f
- Tension spring, for proximal interphalangeal joint, 393, 393f
- Tensor fascia lata, dissection of, in Pemberton osteotomy, 624p–625p
- Tensor fascia lata muscle, dissection of, in trochanteric epiphysiodesis, 588p–589p
- Teratogen(s), and limb absence, 1752
- Teratologic hip dislocation, 513, 646, 650, 650f
- Teres major muscle, transfer to rotator cuff, for brachial plexus palsy, 485, 486f, 488p–491p
- Teres major tendon transfer, for subscapularis muscle paralysis, in poliomyelitis, 1407, 1409f
- Teres minor muscle, division of, for brachial plexus palsy, with shoulder deformities, 118f, 486
- Terminal limb deficiency(ies), 1747, 1747f–1748f
- Tetanus immunization, 2075
- Tethered spinal cord. *See* Spinal cord, tethered.
- Tetracycline labeling, of osteoid osteoma, 1934–1935
- Tetraplegia, definition of, 44
- Texas Scottish Rite Hospital for Children
history and physical exam form of, 58
muscle strength grading system of, 45, 46t
- Texas Scottish Rite Hospital (TSRH)
instrumentation, for idiopathic scoliosis, 243, 243f, 246p–257p
dissection in, 250p–251p
hooks and screws for, 246p–257p
incision in, 248p–250p
patient positioning in, 248p–249p
placement of pedicle and laminar hooks in, 252p–253p
- Thanatophoric dwarfism, 1521–1523, 1522f
- Thenar release, in arthrogryposis, 1660
- Thenar space, infections of, 478
- Thigh-foot axis, assessment of, 66, 66f
- Thigh-leg test, for leg length inequality, 1043–1044, 1044f
- Thomas test, 38, 39f, 1173, 1178f
- Thomsen's myotonia, 1489, 1490t
anesthesia in, 123t
- Thoracic cage, deformities of, in scoliosis, 220
- Thoracic dysplasia, asphyxiating, 1682f, 1683
- Thoracic kyphosis, in Scheuermann's kyphosis, 101, 101f
- Thoracic spine
anatomy of, 2101, 2101f
dextroscoliosis of, 147f
injuries to, 2101–2110, 2101f–2110f, 2106t
classification of, 2103, 2106t, 2107f
complications of, 2107–2108, 2109f
diagnosis of, 2102f, 2102–2103, 2104f
differential diagnosis of, 2103, 2105f
mechanism of, 2102
treatment of, 2105–2107, 2108f–2109f
without radiographic abnormality, 2110, 2110f
- kyphosis of, in neurofibromatosis, 145f
- lordosis of, with lumbar kyphosis, in
myelomeningocele, 1279f
- segmentation-fusion anomalies of, computed tomography in, 144f
- Thoracoacromial artery, dissection of, in shoulder disarticulation, 2014p–2015p
- Thoracolumbar curve(s), in idiopathic scoliosis
anterior instrumentation for, 262p–275p
diaphragm dissection in, 266p–267p
dissection in, 264p–271p
incision in, 262p–263p
positioning in, 262p–263p
protection of aorta and vena cava in, 268p–269p
rib resection in, 264p–265p
solid rod, 260–261, 260f–261f
spinal fusion for, 236, 238f
- Thoracolumbar spine
extension of, 36, 37f
flexion of, 36, 36f–37f
kyphosis of
in achondroplasia, 1516–1517, 1516f–1517f
in Hurler's syndrome, 1573, 1573f–1574f
lateral bending of, 36, 37f
range of motion of, 35–36, 36f–38f
rotation of, 36, 38f
- Thoracolumbosacral orthoses (TLSOs)
for idiopathic scoliosis, 229–231, 230f–231f, 233f
for Scheuermann's kyphosis, 328

Thoracoplasty, for rib deformities, in scoliosis, 239
 Thoracoscopic anterior release, with spinal fusion, for Scheuermann's kyphosis, 333–334
 Thoracotomy
 for spinal fusion, in Scheuermann's kyphosis, 333
 scoliosis from, 308
 Thorax, narrow, in thanatophoric dwarfism, 1522f, 1523
 "Thorn," in developmental dysplasia of hip, 532–533, 534f
 "Three-column spine," 2101, 2101f
 Three-point fixation, in casting, 2079, 2079f
 Thrombocytopenia, with absent radius, anesthesia in, 125t
 Thrombocytopenia-absent radius (TAR) syndrome, anomalies in, 1752, 1753f
 Thrombosis
 and Legg-Calvé-Perthes disease, 657
 in homocystinuria, 1597
 Thumb
 angled, 449
 anomalies of, in fibrodysplasia ossificans progressiva, 1634, 1634f
 clamped, 447
 rotation flap for, 444, 447f
 deformities of
 from rheumatoid arthritis, 474
 in radial deficiency, 1789
 duplication. *See Polydactyly, radial.*
 dysplasia of, 444–449
 historical perspective on, 444–445
 hypoplastic, 448–449
 macrodactyly of, 458, 460f
 mobility of, and hand function, 406f
 spastic, 446
 splints for, 394, 395f–396f, 396
 synovitis of, 472
 trigger thumb, 445
 triphalangeal, treatment of, 431, 431f
 Thumb sign, in Marfan's syndrome, 1587, 1587f
 Thumb-index web, syndactyly of, 442, 444, 445f–447f
 Thumb-in-palm deformity
 in cerebral palsy, surgical treatment of, 1213, 1222, 1222f–1224f
 in Freeman-Sheldon syndrome, 1663f
 Thurston-Holland sign, 2063, 2141, 2142f, 2168, 2176f, 2177, 2178f
 Thyroxine, for hypothyroidism, 1713
 Tibia
 anatomy of, 2371–2372, 2372f
 bowing of, 867
 congenital anterolateral. *See also Tibia,*
 congenital pseudarthrosis of.
 benign, 881–883, 882f
 congenital posteromedial, 883, 883f–884f
 from congenital band syndrome, 462f
 with talipes calcaneovalgus, 905
 congenital pseudarthrosis of, 867–881, 869f–880f
 classification of, 868
 clinical features of, 868
 etiology of, 867–868
 in neurofibromatosis, 867–868, 1628, 1630f
 natural history of, 868, 869f
 pathology of, 867–868

Tibia (*Continued*)
 treatment of, 868–881, 869f–880f
 bypass bone graft in, 869, 871f, 873
 electrical stimulation in, 878
 external fixation and distraction
 osteogenesis in, 878–881, 879–880
 fracture prevention in, 868–869, 870f–872f
 intramedullary fixation in, 872f–877f, 873–877
 principles of, 868
 vascularized fibular graft in, 877–878, 878f
 with late fracture, 881
 distal
 fractures of. *See Ankle, fractures of.*
 injuries to, classification of, 2392
 epiphysiodesis of, for valgus deformity, 2376, 2376f
 fractures of, 2371–2373, 2372f
 diaphyseal, 2376–2381, 2378f–2382f
 classification of, 2376
 diagnosis of, 2377, 2378f
 distal metaphyseal, 2387
 from congenital pseudarthrosis, 868–869, 870f–872f
 late, 881
 in toddlers, 91–92, 92f
 open, 2381–2387, 2382f–2387f
 classification of, 2381
 complications of, 2385–2387, 2386f–2387f
 delayed union of, 2386, 2387f
 external fixation of, 2383, 2384f–2385f
 prognosis in, 2383, 2385
 vascular injury in, 2383
 wound care in, 2382f, 2382–2383
 proximal metaphyseal, 2373–2376, 2373f–2376f
 complications of, 2373, 2374f
 greenstick, 2373, 2373f
 treatment of, 2375, 2375f–2376f
 stress, 2387–2389, 2388f
 with ipsilateral femoral fractures, 2389f, 2389–2390
 hemophilic pseudotumor of, 1885f
 intramedullary fixation of, Bailey-Dubow rod for, 1734f, 1734–1735
 lengthening of, for leg length inequality, 1081–1082
 ossification of, 893f, 2391
 osteofibrous dysplasia of, 1914–1917, 1917f–1919f. *See also Campanacci's disease.*
 osteosarcoma of, 1960f
 physeal arrest in, after ankle fractures, 2403–2405, 2404f, 2406f
 prediction of remaining growth in, 1048–1053, 1049t, 1050f–1055f, 1052t, 1056p–1057p
 for timing of epiphysiodesis, 1053–1058, 1056p–1057p, 1058f
 Menelaus method for, 1051, 1055f
 proximal, epiphysiodesis of, for leg length inequality, 1068p–1071p
 proximal physeal fractures of, 2344–2351, 2345f–2351f
 anatomic considerations in, 2344–2345
 classification of, 2345–2346, 2345f–2346f
 complications of, 2348–2351
 diagnosis of, 2347

Tibia (*Continued*)
 mechanism of, 2346–2347
 treatment of, 2347–2348, 2348f–2350f
 pseudarthrosis of, and amputation, 881
 rotational osteotomy of, for cerebral palsy, in knee, 1169–1173, 1178f
 sectioning of, in below-knee amputation, 1998p–1999p
 shortening of, for leg length inequality, 1077, 1077f–1078f
 spinal fractures of, 2353–2357, 2354f–2359f
 anatomic considerations in, 2354, 2354f
 classification of, 2354, 2355f
 complications of, 2355–2357
 diagnosis of, 2354–2355
 mechanism of, 2354
 treatment of, 2355–2356, 2356f–2359f
 arthroscopic, 2355–2356, 2358f
 casting in, 2355
 tuberosity fractures of, 2341–2344, 2342f, 2344f
 anatomic considerations in, 2341–2342
 classification of, 2342, 2342f
 complications of, 2343–2344
 diagnosis of, 2343
 mechanism of, 2342–2343
 treatment of, 2343, 2344f
 valgus deformity of, 1107, 1109f
 from proximal metaphyseal fractures, 2373, 2374f
 in hereditary multiple exostosis, 1923
 varus deformity of, with Blount's disease, 1104f, 1107, 1109f
 Tibia vara, 840–855, 840f–858f
 adolescent, 850–855, 852f–856f, 853t
 clinical features of, 851, 852f–853f
 definition of, 850
 etiology of, 850–851
 radiographic features of, 851, 852f–853f, 853t
 treatment of, 851–855, 854f, 856f
 external fixation in, 853, 855, 856f
 lateral epiphysiodesis in, 851–853, 854f
 osteotomy in, 851, 852f–853f
 classification of, 840
 definition of, 840
 from focal cartilaginous dysplasia, 855, 857f–858f, 858t
 infantile, 840–850, 843t, 843f–849f
 classification of, 844f, 844–845, 846f
 clinical features of, 841f, 841–842, 843f, 843t
 differential diagnosis of, 844
 etiology of, 840
 physiology of, 840–841
 radiographic features of, 842–844, 843t, 843f–844f
 treatment of, 845–850, 847f–849f
 orthoses in, 845–847, 847f
 osteotomy in, 846f–849f, 847–850
 Tibial artery(ies)
 anatomy of, 2416, 2418f
 anterior, injury to, from tibial osteotomy, 850
 Tibial deficiency, 1778–1784, 1779f–1784f
 anomalies in, 1780
 clinical features of, 1779–1780, 1783f–1784f
 Jones classification of, 1779, 1782f–1784f
 prostheses for, 1805
 treatment of, 1782–1784

- Tibial epiphysis
hypoplasia of, and genu valgum, 1680, 1680f
ossification of, in infantile tibia vara, 844, 844f
- Tibial fracture(s), proximal, genu valgum from, 861, 864f
- Tibial growth plate, preservation of, ankle arthrodesis with, 1395, 1400p–1405p
- Tibial metaphysis, osteochondroma of, angular deformity from, 1104f
- Tibial osteotomy
distal, for valgus ankle deformity, in myelomeningocele, 1265, 1268f
for adolescent tibia vara, 851, 852f–853f
for Ellis–van Creveld syndrome, 1681
for infantile tibia vara, 846f–849f, 847–850
for metaphyseal chondrodysplasia, 1548f–1549f
- Tibial plateau fracture(s), diagnosis of, computed tomography in, 143f
- Tibial tendon(s). *See Tibialis anterior tendon; Tibialis posterior tendon.*
- Tibial torsion, 863–867, 867f
and alignment of foot, 1378, 1388f
and intoeing, 66, 66f, 866
clinical features of, 866
developmental anatomy and, 864, 866
from hip contractures, in poliomyelitis, 1330–1331
in femoral anteversion, 783, 786f
lateral, 866
treatment of, 866–867, 867f
- Tibialis anterior muscle, paralysis of, in poliomyelitis, 1364–1365, 1365f, 1366p–1371p
- Tibialis anterior tendon, transfer of
for correction of vertical talus, 972p–973p
for revision of talipes equinovarus treatment, 945–946
lateral, to metatarsal base, in poliomyelitis, 1364
split, for equinovarus deformity, in cerebral palsy, 1145–1146, 1148p–1149p
to calcaneus, 1264–1265, 1266p–1267p
- Tibialis muscle, anterior, contraction of, during gait, 79–81, 80f
- Tibialis posterior muscle, unopposed activity with triceps surae muscle, in poliomyelitis, 1365, 1372f, 1374p–1377p
- Tibialis posterior tendon
anterior transfer of
for pes cavus, 991
through interosseous membrane, in Duchenne muscular dystrophy, 1470, 1472p–1475p
lengthening of, for equinovarus deformity, in cerebral palsy, 1143, 1143f
transfer of
split, for equinovarus deformity, in cerebral palsy, 1143–1145, 1144f–1147f
through interosseous membrane, for anterior tibial muscle paralysis, 1374p–1377p
to dorsum of foot, for pes cavovarus, 1327, 1440f
to os calcis, for calcaneus deformity, in poliomyelitis, 1380p–1385p
- Tibiofemoral angle, changes in, in physiologic genu varum, 839, 842f
- Tibiofemoral joint, congenital subluxation of, 825
- Tibiofibular diastasis, in tibial deficiency, 1779, 1782f
- Tibiofibular joint, proximal, dislocation of, 2357–2361, 2359f–2361f
anatomic considerations in, 2359, 2359f
classification of, 2359, 2360f
complications of, 2361
diagnosis of, 2359–2360
mechanism of, 2359–2360
treatment of, 2360, 2361f
- Tibionavicular ligament, release of, for talipes equinovarus, 926, 926f
- Tibiotalar joint, release of, for talipes equinovarus, 937, 938f
- Tile and Pennal classification, of pelvic fractures, 2252, 2254, 2254f
- Tillaux fracture(s), 2405–2410, 2407f–2410f
closed reduction of, 2405, 2407
mechanism of, 2405, 2407f
open reduction of, 2407, 2409f–2410f
prognosis in, 2407, 2410
radiographic features of, 2405, 2408f
- Tissue transfer, free, microsurgery for, 504, 505f–506f
- TLSO(s) (thoracolumbosacral orthoses)
for idiopathic scoliosis, 229–231, 230f–231f, 233f
for Scheuermann's kyphosis, 328
- Toddler(s). *See also under specific disorder.*
ambidexterity in, 21
limping in, differential diagnosis of, 85, 86f
- Toddler's fracture, 2377
of tibia, 91–92, 92f
- Toes(s)
clawing of, in pes cavus, 991, 995f, 998f
curly, 1032, 1032f
deformities of, 1012f–1029f, 1012–1036, 1030p–1035p. *See also under Hallux*
fifth, varus, 1024, 1027f–1029f,
1030p–1031p, 1032. *See also Varus fifth toe.*
fractures of, 2431–2432
glomus tumors of, 1032
great, malformations of, in fibrodysplasia ossificans progressiva, 1632, 1633f
hammer, 1027, 1034p–1035p
ischemic necrosis of, from correction of talipes equinovarus, with residual deformity, 953, 957f
mallet, 1027, 1033f
polydactyly of, 1021–1023, 1022f
with hallux varus, 1019, 1019f
syndactyly of, 1023–1024, 1023f–1024f
- Toe extensor(s)
paralysis of, in poliomyelitis, 1365, 1372f, 1374p–1377p
transfer to metatarsal heads, for pes cavus, 991, 994, 995f, 996p–997p
- Toe phalanges, transplantation of, to partial fingers, 1783, 1785f
- Toe transfer, free, microsurgery for, 504, 508, 508f
- Toe-ing-in
assessment of, 65f–66f, 65–67
persistent, with talipes equinovarus, supramalleolar osteotomy for, 950–951, 952f
- Toe-walking
conditions causing, 1135
in toddlers, 85, 86f
- Tone and Positioning (TAP) splint, 396, 397f
- Tone-reducing ankle-foot orthoses, for cerebral palsy, 1131
- Tonic neck reflex(es), 53–54, 54f–55f
- Tonnis osteotomy, for acetabular repositioning, 632–633
- TORCHES infection(s), and cerebral palsy, 1123
- Torode and Zieg classification, of pelvic fractures, 2254–2255, 2255f, 2259f, 2261f
- Toronto brace, for Legg–Calvé–Perthes disease, 695f, 695–696
- Torsionometer, Perdriolle, 225, 226f
- Torticollis
acquired, 187–195, 188f–193f
and atlantoaxial rotatory displacement, 187–191, 188f–193f
inflammatory/septic, 193–195
neurogenic, 191–193, 193f
ocular, 194–195
congenital, 171–187, 173f–187f
and Klippel–Feil syndrome, 173–187, 178f–187f. *See also Klippel–Feil syndrome.*
differential diagnosis of, 171–172, 172f
muscular, 171–173, 173f–177f
clinical features of, 171, 173f–174f
etiology of, 171
treatment of, 172–173, 175f–177f
paroxysmal, of infancy, 193
with developmental dysplasia of hip, 517, 517f, 519, 519f
- Torus fracture(s), 2059–2060, 2060f–2061f
- Total contact molding insert(s), 390f–391f, 390–392
- Total joint arthroplasty
for hemophilic arthropathy, 1889
for rheumatoid arthritis, 1821
of hip
for cerebral palsy, 1201
for osteopetrosis, 1553
- Totally implantable lengthening device(s), for limb lengthening, 1094–1095
- Tourniquet, for regional anesthesia, 115
- Tourniquet(s), and sickle cell disease, 1894
- Toxic synovitis, 1823–1825
vs. septic arthritis, 1850
- T-plate, for wrist fusion, 471, 472f
- TR (repetition time), in magnetic resonance imaging, 152, 152t
- Traction
before femoral osteotomy, 697
for acetabular fractures, 2268
for atlantoaxial rotatory displacement, 189
for developmental dysplasia of hip, in child ages 6 months to 2 years, 539–540, 539f–540f
for femoral shaft fractures, 2305, 2307f, 2307–2308
for hemophilic arthropathy, 1889
for Legg–Calvé–Perthes disease, 692–693, 695f
for reduction of hip dislocation, 2275, 2280
for reduction of spondylolisthesis, 366, 367f
for supracondylar humeral fractures, 2156, 2159f
suspension, after hamstring lengthening, 1336, 1337f
with femoral osteotomy, for limb lengthening, 1079, 1079f
- Traction epiphysis, 1045
- "Traction reduction," for developmental dysplasia of hip, 540
- Traction/countertraction technique, for reduction of glenohumeral dislocation, 2130, 2130f

- Transarticular screw fixation, C1–2, for atlantoaxial rotatory displacement, 191, 192f
- Transfusion(s)**
during scoliosis surgery, requirement for, preoperative assessment of, 240
for scoliosis surgery, using Texas Scottish Rite Hospital instrumentation, 246p minimization of, 116–117
- Transiliac lengthening, for limb lengthening, 1081, 1081f
- Transitional popliteal cyst(s), 830f, 831
- Transphyseal screw(s), in epiphyseal stapling, for leg length inequality, 1063, 1074f
- Transplantation
bone marrow
for Hurler's syndrome, 1575
for osteogenesis imperfecta, 1729
for osteopetrosis, 1553
physeal, for limb lengthening, 1078
umbilical cord blood, for osteopetrosis, 1553
- Trapezius muscle, dissection of, in forequarter amputation, 2002p–2003p
- Trapezius transfer, Saha, for deltoid paralysis, 1406f, 1406t, 1406–1407
- Trauma**
ABCs of, 2072–2073
and adolescent tibia vara, 850–851
and Legg-Calvé-Perthes disease, 662–663
and lumbar disk herniation, 371
and pes cavus, 985
multiple. *See* Multiple injury(ies).
- Traumatic amputation, 1790–1792, 1790f–1793f, 1791t. *See also* Amputation, traumatic.
- Treacher Collins syndrome, anesthesia in, 125t
- Trendelenburg gait, 78, 84, 84f
during ambulation with prosthesis, 1801
- in developmental dysplasia of hip, 526, 527f
in Duchenne muscular dystrophy, 1468f, 1469
in Legg-Calvé-Perthes disease, 672f
- Trendelenburg's test, 41–42, 42f–43f
in Legg-Calvé-Perthes disease, 672, 672f
- Trehphination, of unicameral bone cysts, 1905–1906
- Trethowan's sign, in slipped capital femoral epiphysis, 719, 719f
- Trevor's disease, 1946, 1947f–1948f
- Triamcinolone, intra-articular injection, for rheumatoid arthritis, of hand, 474, 474f–475f
- Triceps brachii muscle
anterior transfer of, in poliomyelitis, 1419, 1425f
paralysis of, in poliomyelitis, 1419
strength of, after Steindler flexorplasty, 1408, 1418
- Triceps insufficiency, muscle transfers for, after talipes equinovarus surgery, 946, 947f
- Triceps reflex, 56
- Triceps surae muscle
contractures of, in vertical talus, 964f
function of, 1361
paralysis of
 calcaneocavus foot deformity from, 1365, 1378f
 in poliomyelitis, 1365–1373, 1373f, 1378f–1379f, 1380p–1385p, 1386f–1387f
unopposed activity with tibialis posterior muscle, in poliomyelitis, 1365, 1372f, 1374p–1377p
- Triceps transfer, for elbow flexion, in arthrogryposis, 1655
- Tricepsplasty, elbow release with, for arthrogryposis, 1654, 1656p–1659p
- Trident hand, in achondroplasia, 1511, 1511f
- Trigger finger, in Hurler's syndrome, 1575, 1577f
- Trigger thumb, 445
- Triphalangeal thumb, treatment of, 431, 431f
- Triplane fracture(s), 2410–2414, 2410f–2417f
closed reduction of, 2411–2412
mechanism of, 2410–2411, 2410f–2411f
open reduction of, 2412, 2414, 2415f–2417f
prognosis in, 2414
radiographic features of, 2411, 2412f–2414f
- Triple arthrodesis. *See* Arthrodesis, triple.
- Triradiate cartilage
anatomy of, 2252f
injury to, in acetabular fractures, 2269–2270, 2270f
- Trisomy 21. *See* Down syndrome.
- Trisprial tomography, of scaphoid fractures, 2247
- Trochanter, greater
avulsion injury of, 2260f
position of, in avascular necrosis, 583
- Trochanteric advancement, lateral, for avascular necrosis, in developmental dysplasia of hip, 610p–611p
- Trochanteric epiphysiodesis
after femoral osteotomy, 697, 700f
for avascular necrosis, in developmental dysplasia of hip, 584, 588p–595p
destruction of growth plate in, 592p–593p
dissection in, 588p–591p
incision for, 588p–589p
postoperative care in, 594p–595p
removal of bone plug in, 590p–591p
- greater, for developmental coxa vara, 772–773
- Trochanteric osteotomy, in iliopsoas muscle transfer, 1340–1341p
- Trochanteric transfer, for avascular necrosis, in developmental dysplasia of hip, 584, 598p–611p
dissection in, 598p–603p
incision for, 598p–599p
osteotomy in, 602p–607p
position for, 598p
postoperative care in, 608p
with lateral wedge valgus osteotomy, 584–586, 614p–617p
- Trochlea, fractures of, 2205
- Truncation artifact(s), on magnetic resonance imaging, 155
- Trunk, in poliomyelitis, 1395
- Trunk incursion reflex, 52, 54f
- TSRH instrumentation. *See* Texas Scottish Rite Hospital (TSRH) instrumentation.
- Tuberculosis
of spine, 1831–1835, 1832f–1835f. *See also* Spondylitis, tuberculous.
osteoarticular infections from, 1865–1868, 1866f–1867f. *See also* Arthritis, tuberculous; Osteomyelitis, tuberculous.
- Tuberous sclerosis, anesthesia in, 125t
- Tubigrip, for securing total contact molding inserts, 390, 391f, 392
- TUBS, in shoulder instability, 2131
- Tumor(s). *See under* specific site, e.g., Bone tumors; *under* specific type, e.g., Osteosarcoma.
- Tumor-related hypophosphatemic rickets, 1695
- UCBL orthosis
for pes planovalgus, 914, 914f
for tarsal coalition, 980
- Ulcer(s)
decubitus. *See* Pressure sore(s).
in dermatomyositis, 1494f
- Ulna
deformities of, in hereditary multiple exostosis, 1922, 1923f
fractures of
 distal phyeal, treatment of, 2238
 with radial fractures, 2225–2233, 2226f–2234f
 anatomic considerations in, 2225
 complications of, 2231–2233, 2234f
 diagnosis of, 2225
 malunion of, 2227, 2229, 2230f, 2231–2233, 2234f
 mechanism of, 2225
 radiographic features of, 2225, 2226f, 2227
 treatment of, 2227–2231, 2228f–2232f
 closed, 2227–2230, 2228f–2230f
 open, 2230–2231, 2231f–2232f
 with radial head dislocation. *See* Monteggia fracture(s).
 malunion of, after Monteggia fracture, 2224
- Ulnar deficiency, 411–414, 412f, 414f, 1787–1790, 1789f–1790f
classification of, 412f, 412–413, 1788, 1789f
clinical features of, 412, 1788–1789, 1790f
etiology of, 411, 1787–1788
historical perspective on, 411
imaging studies in, 413
pathology of, 413
treatment of, 413–414, 414f, 1789–1790
- Ulnar dimelia, 423–426, 425f
- Ulnar nerve
compression neuropathy of, in Volkmann's contracture, 408, 410f
dissection of
 in amputation through arm, 2018p–2019p
 in disarticulation of elbow, 2022p–2023p
- injury to
 from elbow dislocation, 2193
 from pinning of humeral fractures, 2153, 2155, 2155f, 2160–2162, 2162f
- palsy of, congenital band syndrome and, 462f
- protection of
 during percutaneous pinning of supracondylar humeral fractures, 2153, 2155, 2155f, 2160–2162, 2162f
 in Steindler flexorplasty, 1410p–1413p
- Ulnar polydactyly. *See* Polydactyly, ulnar.
- Ulnar-femoral syndrome, 1754, 1754t
- Ulnohumeral joint, range of motion of, 31, 33
- Ultrasonography, 163–165, 165f
B-mode, 163
- Doppler, 163
- image generation in, 163
- in acute transient synovitis of hip, 1824
- in developmental dysplasia of hip, 527–530, 528f–531f, 529f
advantages of, 527
- anatomic landmarks on, 528f
- Graf classification in, 527–530, 529f, 529t
sensitivity of, 528–530, 530f
- in hemophilic arthropathy, 1883, 1883f
- in leg length inequality, 1045
- in Legg-Calvé-Perthes disease, 683
- in measurement of femoral anteverision, 784
- in slipped capital femoral epiphysis, 721

- Ultrasonography (Continued)**
- indications for, 164f–165f, 165
 - physical principles of, 163
- Umbilical cord blood transplantation**, for osteopetrosis, 1553
- Unipolar pectoralis major transfer**, for elbow flexion, in arthrogryposis, 1655
- Upper extremity(ies).** *See also* Arm; Forearm; Hand; Wrist.
- acute care of, 398–404, 399f–403f. *See also* Hand, acute care of.
 - amputation of, 1793
 - aneurysms of, 497
 - cerebral palsy in, 1211–1222, 1212f–1213f, 1214p–1221p, 1222f–1225f. *See also* Cerebral palsy, upper extremity in.
 - congenital anomalies of, 410–467. *See also under specific anomaly, e.g., Ulnar deficiency.*
 - classification of, 410t, 410–411
 - diagnosis of, 411, 411t
 - embryology of, 379–380
 - epidermoid inclusion cysts of, 497
 - epiphyses of
 - closure of growth plates in, age at, 7f
 - secondary ossification centers in, age at appearance of, 6f
 - foreign bodies in, 497
 - function of, in myelomeningocele, 1255
 - growth of, cerebral palsy and, 1209
 - injuries to, 2115–2250. *See also under specific site, e.g., Clavicle, injuries to.*
 - long bones of, growth of, physiologically, 10f
 - manual muscle tests of, 59–61
 - microsurgery on, 503–504, 505f–511f, 512. *See also* Microsurgery.
 - muscles of, innervation of, 47t
 - physical examination of, 380
 - prostheses for, 1800–1801
 - pseudoaneurysms of, 497, 1796–1798
 - regional anesthesia for, 115–116
 - surgical procedures on
 - goals of, 379
 - timing of, 380–382, 381t–384t, 383f
 - tumors of, 487, 496–497, 502–503
 - imaging studies of, 496
 - neoplastic, 502–503
 - non-neoplastic, 496–497, 502–503
 - vascular malformations of, 497, 502
 - weakness of, in Charcot-Marie-Tooth disease, 1441, 1444, 1447f
- Upper respiratory infection(s), and anesthesia, 110**
- Urinalysis**, of glycosaminoglycans, in mucopolysaccharidosis, 1572
- Urologic disorder(s)**, from myelomeningocele, 1258–1259
- Uveitis**, in rheumatoid arthritis, 1816
- V radiolucency**, in coxa vara, 766, 766f
- Vaccination**, for tetanus, 2075
- Vaccine-associated paralytic poliomyelitis (VAPP)**, 1321
- VACTERL association**, 1754
- Valgus deformity**, 25
- from traction, for developmental dysplasia of hip, 583, 583f
 - of foot. *See* Pes valgus.
- Valgus osteotomy**
- closing wedge, of proximal femur, with distal-lateral advancement of greater trochanter, 614p–617p
 - for angular deformities of leg, 1107, 1110f
 - for developmental coxa vara, 772–773, 774f–775f
 - for hip fractures, 2296, 2298f–2299f
 - for Legg-Calvé-Perthes disease, 699–700
 - of proximal femur, for nonreconstructable hip in cerebral palsy, 1201, 1208f
- Valgus overcorrection**, from treatment of talipes equinovarus, 942–943, 943f–944f
- van Ness rotationplasty**, for proximal focal femoral deficiency, 1763, 1768f–1769f
- VAPP (vaccine-associated paralytic poliomyelitis)**, 1321
- Varicostas**, in Klippel-Trénaunay syndrome, 1671
- Varization osteotomy**, for paralytic dislocation of hip, in poliomyelitis, 1349–1350, 1352p–1355p
- Varus deformity**, 25
- from treatment of developmental dysplasia of hip, 583, 585f–586f
 - of ankle, after fractures, 2405, 2406f
 - of tibia, with Blount's disease, 1104f, 1107, 1109f
- Varus femoral osteotomy**, for Legg-Calvé-Perthes disease, 696–698, 699f–700f
- Varus fifth toe**, 1024, 1027f–1029f, 1030p–1031p, 1032
- Varus features**, of 1024, 1027f
- Treatment of**
- Butler procedure in, 1024, 1028f–1029f
 - McFarland operation in, 1024, 1030p–1031p
- Varus osteotomy**
- closing wedge, for ankle valgus, in hereditary multiple exostosis, 1924, 1925f
 - with tendon transfer, for subluxation/dislocation of hip, in myelomeningocele, 1274
- Vascular fibrous tissue**, of bone marrow space, in Caffey's disease, 1562, 1562f
- Vascular injury(ies)**, 2078f, 2078–2079
- from elbow dislocation, 2193
 - from femoral shaft fractures, 2319–2320
 - from open tibial fractures, 2383
 - from supracondylar humeral fractures, 2163–2165
- Vascular malformation(s)**
- diagnosis of, magnetic resonance imaging in, 157f
 - of upper extremities, 497, 502
- Vascular system**
- anatomy of, and osteomyelitis, 1841–1842, 1843f
 - injuries to, with pelvic fractures, 2256
- Vascularized fibular graft**, for congenital anterolateral tibial bowing, 877–878, 878f
- Vastus lateralis muscle**, dissection of in intertrochanteric varus osteotomy, and internal fixation with blade plate, 572p–573p
- Vastus medialis obliquus (VMO) ligament**, tears of, 2351–2352
- VATER association**, 1752, 1754
- VATS (video-assisted thoracoscopic surgery)**
- for idiopathic scoliosis, 240
 - for Scheuermann's kyphosis, 333–334
- Vaulting**, during ambulation with prosthesis, 1802
- VDRO (femoral varus derotation osteotomy)**, for hip subluxation/dislocation, in cerebral palsy, 1189, 1194, 1194f
- Vena cava**, protection of, in anterior instrumentation, for idiopathic scoliosis, 268p–269p, 271p
- Venous malformation(s)**, diagnosis of, magnetic resonance imaging in, 157f
- Ventriculoperitoneal shunting**, in myelomeningocele, 1257
- Verbal skill(s)**, developmental milestones for, 21, 21t
- Verebelyi-Ogston procedure**, 1653–1654
- Verrucous hyperplasia**, in neurofibromatosis, 1625
- Vertebra(e)**
- abscesses of, in tuberculous spondylitis, 1832–1833, 1833f
 - block, and congenital scoliosis, 284, 284f, 288f
 - destruction of, in tuberculous spondylitis, 1831–1832, 1832f–1835f
 - displacement of, congenital, 297, 298f
 - formation of, defects of
 - and congenital kyphosis, 291, 293f, 294–295, 295f–298f
 - and congenital scoliosis, 283–284, 283f–284f
 - growth of, 239, 240f
 - in sickle cell disease, 1893f, 1893–1894
 - of cervical spine
 - anomalies of, in Klippel-Feil syndrome, 174–176
 - range of motion of, 34t
 - pathology of, in myelomeningocele, 1252
 - rotation of, in scoliosis, measurement of, 225–226, 226f–227f
- Scalloping of**, in neurofibromatosis, 301, 303f–304f
- Segmentation of**
- defects of, and congenital scoliosis, 283–285, 284f–286f
 - failure of, and congenital kyphosis, 291, 294f, 295, 297
- Vertebra plana**, in eosinophilic granuloma, 1939, 1940f
- Vertebral apophysis**, slipped, 99, 374–375
- Vertebral body(ies)**
- and neural arches, synchondrosis between, 130f
 - beaking of
 - in Hurler's syndrome, 1575, 1576f
 - in Morquio syndrome, 1578f, 1579
 - shape of, in idiopathic scoliosis, 219–220, 220f
 - trapezoid-shaped, in Scheuermann's kyphosis, 324, 327f
- Vertebral end plate(s)**, removal of, in anterior instrumentation, for idiopathic scoliosis, 270p–271p
- Vertebral osteomyelitis**
- and torticollis, 193–194
 - back pain in, 103
 - vs. diskitis, 87–88
- Vertebrectomy**, Dunn-McCarthy technique of, for kyphosis, in myelomeningocele, 1280, 1282p–1287p
- Vertex presentation**, and neonatal position, 38f

- Vertical suspension positioning reflex, 49
Vertical talus. *See Talus, vertical.*
Vickers' ligament
 in Madelung's deformity, 416–417, 417f
 release of, with dome osteotomy, for
 Madelung's deformity, 419–420, 420f
Video-assisted thoracoscopic surgery (VATS)
 for idiopathic scoliosis, 240
 for Scheuermann's kyphosis, 333–334
Villonodular synovitis, pigmented, 1944–1946,
 1946f–1947f
Viral osteomyelitis, 1870–1871
Visual analog scale(s), for assessment of pain,
 112, 112f
Vitallium staple(s), for epiphyseal stapling, for
 leg length inequality, 1074f
Vitamin B₆, for homocystinuria, 1597
Vitamin C, deficiency of, and scurvy,
 1705–1707, 1706f
Vitamin D
 and calcium metabolism, 1685, 1686f
 deficiency of
 and rickets, 1686. *See also Rickets.*
 radiography in, 129, 129f
 for renal osteodystrophy, 1698
 for rickets, 1689
Vocational rehabilitation, for
 myelomeningocele patient, 1296
Volar fasciotomy, of forearm, 2078f
Volkmann's ischemic contracture
 from Monteggia fractures, 2225
 in hemophilia, 1883, 1883f
 of forearm, 408, 409f–410f, 410
 from casting of supracondylar humeral
 fractures, 2156, 2165
 von Recklinghausen's disease, 451. *See also*
Neurofibromatosis, type 1.
Von Rosen splint, for developmental dysplasia
 of hip, 538f
Von Rosen view, in developmental dysplasia of
 hip, 531
 von Willebrand's disease, 1879
V-osteotomy, Japas tarsal, for pes cavus, 999,
 1006p–1009p
Vulpius procedure, for Achilles tendon
 lengthening, 1136, 1137f–1138f
- Waddell's triad, 2302
Wagner apparatus, for limb lengthening, 1079,
 1080f, 1091–1092
Wagner's intertrochanteric double osteotomy,
 for avascular necrosis, 612p–613p. *See also*
Intertrochanteric osteotomy.
Wake-up test, 241
Waldenström classification, of Legg-Calvé–
 Perthes disease, 673, 673t
 for radiographic staging, 22f–23f, 675t
Walking. *See also Ambulation.*
 energy for, 76–77, 77t
 velocity of, 76
Walking reflex, 49, 51, 53f
Walton classification, of muscular dystrophy,
 1463, 1465t
Warner's syndrome, 1782
Wart(s), of hand, 479
Watson-Jones approach, to bone graft
 epiphysiodesis, for slipped capital femoral
 epiphysis, 731, 732p–733p
- Watson-Jones classification, of tibial tuberosity
 fractures, Ogden modification of, 2342f
WBC (white blood cell) count, in acute
 hematogenous osteomyelitis, 1846
"Web creep," after treatment of syndactyly, 444
Weber forceps, for reduction of triplane
 fractures, 2414, 2415f–2416f
Wedge osteotomy
 for congenital scoliosis, 291, 292f
 Irwin, for genu recurvatum, in poliomyelitis,
 1359, 1360f
Weight, in relation to age, 10, 13f–18f
 for boys, 10, 13f, 15f, 17f
 for girls, 10, 14f, 16f, 18f
Welander's syndrome, anesthesia in, 125t
Werdnig-Hoffmann disease, 1311, 1312f
 anesthesia in, 125t
Wheelchair(s), for myelomeningocele patient,
 1295
"Wheeled stander," 1291, 1292f
Whip, during ambulation with prosthesis, 1803
"Whistling face" syndrome, 1662, 1663f
White and Stubbins technique, of
 epiphysiodesis, for leg length inequality,
 1061–1062, 1062f
White blood cell (WBC) count, in acute
 hematogenous osteomyelitis, 1846
Whitesides needle technique, of
 intercompartmental pressure
 measurement, 2076, 2077f
Whitlow, herpetic, 477, 477f
Wiesel-Rothman technique, for measurement
 of occipito-atlantal instability, 1618, 1619f
Wilberg's center-edge angle, in developmental
 dysplasia of hip, 531, 532f
Williams rod, for fixation of congenital
 anterolateral tibial bowing, 874, 874f–875f,
 877f
Wilmington brace, for scoliosis, idiopathic,
 230–231
Wilms' tumor, with Beckwith-Wiedemann
 syndrome, 1040
Wilson shelf procedure, 647f–648f
Wilson's sign, 790
Wiltse muscle-splitting technique, for
 spondylolisthesis, 362, 362f
Wiltse technique, for congenital fibular
 pseudarthrosis, 886f, 887
Winberger's sign, in scurvy, 1706, 1706f
Wisconsin instrumentation, segmental spinal,
 for idiopathic scoliosis, 242, 242f,
 244p–245p
Withdrawal reflex, 51
Wolff-Parkinson-White syndrome, anesthesia
 in, 125t
Wolf's law, 2059
Wong-Baker Faces Rating Scale, 112, 112f
Word-graphic rating scale(s), for pain
 assessment, 112f
Wormian bone(s), in osteogenesis imperfecta,
 1725–1726, 1727f
Wound(s)
 from animal bites, 478
 from lawnmower injuries, care of, 2434
 from open fractures, care of, 2073, 2382,
 2382f
 gunshot, 2075
 infection of, from instrumentation for
 scoliosis, 258, 1228
 on hand, closure of, 384–385, 385f
 puncture
 infection from, 1863
 of foot, 2434
- Woven bone**
 in Campanacci's disease, 1915, 1917f
 in fibrous dysplasia, 1911, 1913f
Wrap-around artifact(s), on magnetic
 resonance imaging, 155
Wrisberg ligament discoid lateral meniscus,
 814, 814f
Wrist
 deformities of, from brachial plexus palsy,
 481, 483, 484f
 flexor tendons of, fractional lengthening of,
 1216p–1219p
 fracture of, and physeal arrest, 2068f
 fracture-dislocations of, 2248–2249
 fusion of, for synovitis, 471, 472f
 in arthrogryposis, 1655
 in cerebral palsy, surgical procedures on,
 1211, 1212p–1219p
 in rheumatoid arthritis, 1817, 1817f
 splints for, 394, 395f–397f, 396
 synovitis of, 469, 469f
 treatment of, 470–472, 470f–472f
 exercise in, 470
 tenosynovitis of, 469–470
 treatment of, 470–472, 470f–472f
 intra-articular corticosteroid injections
 in, 470, 470f–471f
 operative, 470–472, 472f
 ulnar deficiency in, treatment of, 413
Wrist sign, in Marfan's syndrome, 1587, 1587f
Wry neck, 171–173, 173f–177f
Wynne-Davies criteria, for ligamentous laxity,
 515, 515f
- Xp21, and Duchenne muscular dystrophy, 1463
X-ray equipment, 133–134, 134f
- Y-ligament of Bigelow**, anatomy of, 2273, 2273f
Y-osteotomy, Pauwels' intertrochanteric, for
 developmental coxa vara, 773, 774f, 775,
 776p–779p
Young and Burgess classification, of pelvic
 fractures, 2252, 2253f
Yount's fasciotomy, for iliotibial band
 contractures, in poliomyelitis, 1331, 1337f
"Y-scapular" view, in proximal humeral
 physeal fractures, 2125, 2126f
- Zancolli procedure, for brachial plexus palsy,
 with elbow deformities, 487, 498p–501p
Z-foot, for pes planovalgus, 921f, 921–922
Ziehl-Neelsen stain, in tuberculous
 osteomyelitis, 1866, 1867f
Zielke derotation spondylodesis
 instrumentation, for idiopathic scoliosis,
 259–260, 260f
Zone of polarizing activity, and upper
 extremity development, 379
Zone of polarizing activity (ZPA), in limb
 development, 1745, 1746f, 1747
Zone of Ranvier, 2060
ZPA (zone of polarizing activity), in limb
 development, 1745, 1746f, 1747
Z-plasty, for congenital muscular torticollis,
 172–173, 176f–177f