

A novel approach to the use of genetically modified herbicide tolerant crops for environmental benefit

Alan M. Dewar¹, Mike J. May¹, Ian P. Woiwod², Lisa A. Haylock¹, Gillian T. Champion¹, Beulah H. Garner¹, Richard J. Sands¹, Aiming Qi¹ and John D. Pidgeon¹

¹*Broom's Barn Research Station, Higham, Bury St Edmunds, Suffolk IP28 6NP, UK*

²*Rothamsted Research, Harpenden, Herts. AL5 2JQ, UK*

The proposed introduction of genetically modified herbicide tolerant (GMHT) crops, with claims of improved weed control, has prompted fears about possible environmental impacts of their widespread adoption, particularly on arable weeds, insects and associated farmland birds. In response to this, we have developed a novel weed management system for GMHT sugar beet, based on band spraying, which exploits the flexibility offered by the broad spectrum partner herbicides. Here we show the results from two series of field experiments which, taken together, demonstrate that, by using this system, crops can be managed for enhanced weed and insect biomass without compromising yield, thus potentially offering food and shelter to farmland birds and other wildlife. These results could be applicable widely to other row crops, and suggest that creative use of GMHT technology could be a powerful tool for developing more sustainable farming systems in the future.

Keywords: genetically modified crops, biodiversity, sustainability, sugar beet, weeds

1. INTRODUCTION

Sugar beet is a poor competitor with weeds in arable fields because it is slow growing early in the season and has a low canopy in its first year of a biennial life cycle. Good weed control is therefore essential to produce economically viable yields (Jansen 1972), but is not easy to achieve with current selective herbicides and/or inter-row tillage. From the point of view of competition for resources, principally light, weed control need not be carried out until the 6-8 leaf growth stage of the crop (Scott *et al.* 1979), but the weaknesses of current conventional herbicides dictates that weed control commences pre-emergence or at the cotyledon stage of weeds (and crop). Thus very few weeds are present throughout the season in most crops. However, the few crops that are weedy do offer a food source for migrating seed eating birds in the autumn (Wilson *et al.* 1999; Watkinson *et al.* 2000). It is the potential loss of these weedy crops, amid general alarm over population decline of farmland bird species (Chamberlain *et al.* 2000), that has prompted concerns about genetically modified herbicide tolerant (GMHT) technology in the UK from English Nature (English Nature 1998, 2000) and some environmental scientists (Krebs *et al.* 1999; Hails 2000) and NGOs.

However, our novel approach to weed management in GMHT sugar beet exploits the much greater flexibility and efficacy of the broad-spectrum herbicides, glyphosate and glufosinate-ammonium, to which GM tolerances have been produced. We have developed a simple over-the-row band spraying technique to control in-row weeds first, whilst those between the rows can be controlled by a later overall spray application. This exploits both the temporal and spatial flexibility offered by the GMHT system, to allow weed control tailored precisely to avoidance of competition.

Two series of experiments have tested the hypothesis that in GMHT systems, weed management options could be found which would benefit weed and invertebrate populations of relevance to farmland birds without reducing crop yield.

2. METHODS

A first series of five experiments investigated the effect of weed management strategies on yield of glyphosate tolerant GM sugar beet (L#77 from Monsanto) in East Anglia in 1999 and 2000. The experiments were carried out on soil types typical of those on which sugar beet is grown in the UK. Between 12 and 22 different weed species were present on each experiment, many of them important components of farmland bird diets (Krebs *et al.* 1999). *Chenopodium album* was an important weed on four sites, *Fallopia convolvulus* and *Veronica persica* on three, *Sinapis arvensis*, *Persicaria maculosa*, volunteer cereals, *Cirsium arvense* on two and *Tripleurospermum maritimum*, *Persicaria lapathifolia* and *Alopecurus myosuroides* on one. Total population densities in untreated plots ranged from 29 to 75 m⁻².

The conventional herbicide programmes varied at each site depending on the weed species present. The number of active ingredients ranged from 3 to 8, the simplest including phenmedipham, metamitron, and ethofumesate (site 1, 1999), and the most complex including those three plus paraquat, diquat, desmedipham, lenacil and cycloxydim. Other active ingredients used on the other sites included clopyralid, and triflurosulfuron-methyl. The number of applications ranged from 2 to 4. In 1999, applications of some treatments, particularly in the conventional programmes, were delayed by adverse weather conditions. Treatments of glyphosate (at 1080 g a.i. sprayed ha⁻¹) were applied either overall at several timings between 207 and 864 day degrees above 3°C (°Cd) after sowing, or over the sugar beet rows only at similar but fewer timings up to 586°Cd. The overall treatments were followed by a second application between 698 and 1022°Cd and the band sprays by an overall application

between 401 and 811 °Cd, both depending on the timing of the first sprays. Glyphosate treatments were compared to untreated controls and programmes of current commercial herbicides applied pre-emergence (in two trials), but mostly post-emergence starting between 79 and 222 °Cd.

The biomass of weeds present in each treatment was assessed on six occasions throughout the season, the earliest at the time of the first glyphosate applications in late May (240 °Cd) and the latest in mid-August (1450 °Cd). Biomass was assessed using scores on a linear scale (0-10), where 0 = no living weeds and 10 = full biomass for the time of year and plant stage with no effect on plants. In band-sprayed plots the score was a mean of the sprayed area down the row and unsprayed area between the row. Where weed numbers were low, scores in untreated plots were sometimes less than 10. Sugar yield was assessed at harvest in late August/early September. This was earlier than commercial crops due to the constraints of the consent and the audit requirements of British Sugar.

The environmental impact of all the conventional herbicide programmes was assessed by the Millieumeetlat scoring system (CLM 2000), which evaluates toxicity, mobility and persistence of pesticides based on data from submissions for the registration of pesticides in Europe.

In a second series of four experiments using the same cultivar and husbandry, a subset of the treatments included in the yield experiments was set up in adjacent areas at three (1, 3 and 4) of those five sites, and one at a different site (6). Plot sizes were larger to allow the collection of invertebrates with reduced interference from neighbouring plots. For this study, we selected Carabidae, Staphylinidae and Aranae, sampled in pitfall traps, as indicators of arthropod abundance. Pitfall traps give a measure of abundance and activity and are widely used to study ground dwelling invertebrates (Baars 1979). Carabidae are regarded as useful environmental indicators

(Luff & Woiwod 1995), and both staphylinids (Powell *et al.* 1985) and spiders (Haughton *et al.* 1999) contain species that are known to respond to herbicide regimes. Traps were set for 7 days per sampling occasion (4) in 1999, and for 14 days per occasion (3) in 2000 to increase sample size. Samples have to be standardised for number of individuals for meaningful comparisons of diversity to be made. Here we used the log-series α diversity parameter for reasons given by Taylor *et al.* (1976).

3. RESULTS

(a) Weed management and yield effects

Cumulative weed biomass in untreated plots at four of the five sites was high, but low at site 3 in 2000, where weed numbers were lowest. All treatments at all sites significantly reduced weed numbers and biomass compared to the untreated plots, but control from the early overall glyphosate programmes was generally better than that from the conventional treatments, particularly at sites 2 and 5 (Figure 1). Weed biomass was greater than conventional following the later overall sprays of glyphosate. Biomass in band sprayed plots was much higher between rows than was evident from the scores.

Sugar yields from all trials were lower than would normally be expected from a commercial crop (range 4.9-6.1 t ha⁻¹ in plots treated with conventional herbicides) as a result of the imposed early harvest in late August or early September to comply with British Sugar GM audit requirements. Yield reductions in the untreated plots compared to those treated with conventional herbicides ranged from 24 to 88%. Glyphosate, first applied overall sometime between 240 and 320°Cd, gave the best yields in each trial (range 5.9-6.7 t ha⁻¹) - on average 9.7% greater than the conventional treatments, although the differences were only significant at sites 2 and 5 (Figure 1). In previous trials, yield reductions ranging from 5-15% in conventional treatments compared to the best glyphosate treatment (Brants & Harms 1998; Moll

1997; Wilson 1999; Wilson *et al.* 2002) have been attributed partially to the slight, but occasionally important, phytotoxic effects of conventional herbicides on the beet plants themselves (Wilson *et al.* 2002). In our experiments some of the yield improvement from the glyphosate treatments was probably also a result of better weed control throughout the season.

Delays in glyphosate treatment had a significant effect on final sugar yield (Figure 1), described by the following equation:

$$Y = Y_0 + \alpha / (1 + e^{\beta(X - X_0)}) \text{ (eqn1)}$$

where Y is the sugar yield, Y_0 the yield from the untreated plots, α the maximum reduction of sugar yield observed, β the rate of yield reduction due to delays in treatment. $Y_0 + \alpha$ combines to represent the maximum obtainable sugar yield when weeds are effectively controlled to the full, X_0 the thermal time at which the reduction of sugar yield is at half value of α , and X the thermal time from sowing. When α and Y_0 were allowed to vary from experiment to experiment, but the other parameters were fixed, the total variance accounted for (R^2) was 97.1% ($d.f.=34$).

The yields from the band spray treatments could be described by a simple linear relationship:

$$Y = Y_p - \gamma X \text{ (eqn2)}$$

where Y is sugar yield, Y_p the intercept indicating the potential yield at a given site in a given year, γ the slope measuring the reduction of sugar yield per unit of delay in thermal time from sowing, and X the thermal time delay from sowing. Comparison of regressions from all sites showed that each had a different Y_p but a common slope γ , which was not significantly different from 0, and accounted for 90.2% of the total variance ($d.f.=18$) in the observed sugar yields (Figure 1). Delays in overall sprays

following band treatments resulted in the same amount of sugar yield reduction per unit of thermal time in each trial.

These results suggest that weed control with overall glyphosate applications should commence around 275°Cd for optimum yield return and before 535°Cd if significant yield loss is to be avoided. This broadly agrees with previous work (Scott *et al.* 1979; Wilson *et al.* 2002; Schweizer & Dexter 1987). Results from the band-spray treatments suggest that, following a first spray applied at between 207 and 530°Cd after sowing, the second could be applied much later between 586°Cd and 725°Cd (average 656 °Cd) before significant reductions in yield compared to the conventional regime occur.

(b) *Environmental impact*

In our assessment, using the Millieumeetlat system of the direct environmental impact of the herbicide regimes, scores for the conventional herbicides ranged from 32 to 218 for water organisms, 11 to 960 for soil organisms, and 155 to 16540 for deeper water. The equivalent scores for glyphosate treatments were 0, 5-6 and 0 respectively, even though the latter used the maximum dose recommended on draft labels, and conventional treatments, especially in 1999, were less intensive compared to most commercial treatments used that season as a result of the later sowing. A score greater than 100 is considered unacceptable for an individual application in the Millieumeetlat system. All herbicides were within the acceptable limits for water organisms, but lenacil in experiments 2, 3 and 5 and clopyralid in experiment 4 were above this limit for deeper water, and lenacil (experiments 2 and 5) and paraquat plus diquat (experiments 4 and 5) were above this limit for soil organisms.

(c) *Effects on arthropods*

In our study sites the number of species of carabids, staphylinids and spiders was typical of arable fields (Kromp 1999). Carabids were numerically greater than the

other two groups at all sites, comprising at least 44% of the total collected (site 4), but as high as 84% at site 6, which was situated next to a beetle bank. Staphylinids comprised between 5% (sites 6 and 3) and 30% (site 4), while spiders comprised between 10% (site 6) and 31% (site 1) of the total. Site 3 was the most diverse (particularly in spiders), and caught the largest number of specimens over the sampling period (31911) even taking account of the longer period of collection (Table 1). Site 1 had moderate populations, while site 4 had the fewest individuals.

Among the carabids, the dominant species at all sites was *Pterostichus melanarius*, which comprised at least 70% of the carabids. The Aleocharinae were the dominant group of staphylinids at two of the four sites (6 and 3), constituting 79 and 50% of the populations there respectively. *Philonthus cognatus* was the most important staphylinid species at the other two sites making up 43% at site 1 and 53% at site 4. Spider communities were also dominated by a single species at three of the four sites. *Oedothorax apicatus* was the most common species at sites 1 (61%), 6 (49%), and 3 (69%), but *Erigone atra* was dominant at site 4 (36%). In all three groups the top five species made up at least 87% of the total at any site.

The impact of herbicide treatments on the relative abundance of the three groups depended on the density and diversity of weeds present and the timing and efficiency of their removal. At sites 3 and 6 there was no consistent effect of treatments on the numbers of carabids, staphylinids or spiders at any time during the growth of the crop, only an occasional transitory effect. This was almost certainly due to the low weed populations at these sites (*circa* 11-12 m² in untreated plots), which did not alter the structure of the habitat sufficiently to influence the populations of these arthropods. Indeed the maximum ground cover afforded by those weeds in late July, was only 23 and 16% respectively in the untreated plots at the two sites, compared to 35% at site 1 and 96% at site 4.

At these latter two sites, weed numbers in untreated plots were two to five times greater (27 and 61m⁻² respectively) and the weeds, especially at site 4, were much taller. There were strongly significant correlations at both sites between weed biomass (including dead and dying weeds) in late July and the cumulative numbers of staphylinid beetles collected in the pitfall traps during the sampling period June-August (Figure 2). The correlation was much weaker, although still significant, for carabids but non-significant for spiders.

Thus fewest carabids and staphylinids were found in plots treated with conventional herbicides or early overall applications of glyphosate, and most where weed control was delayed (both sites 1 and 4) or partial as in the band sprayed plots (site 4). There was no difference for any species of carabid or staphylinid, or their combined totals, on any sampling occasion or when considering cumulative totals, between the conventional treatments and the early overall glyphosate treatment. This suggests that the response of the beetles was to the removal of weeds, and not to the chemicals used.

These effects of herbicides were similar to those reported in cereal crops (Powell *et al.* 1985; Haughton *et al.* 1999; Moreby & Southway 1999). In one study carried out in a row crop, Purvis and Curry (1984) reported that carabids were rarely affected by weediness in sugar beet fields but staphylinid beetles were substantially increased, especially *P. cognatus*, while spiders were unaffected. Spiders are known to be affected by herbicide regimes but the response is more apparent in some families than others (Haughton *et al.* 1999).

As expected there were site differences in invertebrate diversity. The most diverse site was 3 and the least, 6, which was significantly less diverse. However there were no significant differences between any of the other sites (Table 1). Within any site there was no significant difference in the log series α index of biodiversity

between any treatment on any one sampling date, or when the cumulative catch over all sampling dates was considered, even at the two sites which showed significant effects of treatments on the number of carabid and staphylinid beetles. There was no difference between the conventional or early applied glyphosate herbicides. The lack of effects of treatments on biodiversity for individual sample dates is not surprising as any actual change in species complement would only be likely to occur over long periods for the relatively widespread and abundant farmland species being sampled, many of which have only one generation per year.

4. DISCUSSION

These experiments demonstrate that GMHT technology allows a flexible knowledge-based management approach to weed control in sugar beet, permitting higher weed populations early in the season than is possible in conventional systems. The models of yield effects described here could be used to determine weed control requirements quantitatively. In some low weed pressure situations, such as at site 3 (Figure 1), only one well-timed spray would be needed to achieve satisfactory commercial weed control.

Inputs could be tailored to weed pressure and environmental objectives, such as weed free fields for bird species such as the stonecurlew (*Burhinus oedicanus*) or to provide low growing vegetation for skylarks (*Alauda arvensis*), both species of current conservation concern in the UK. These and other scenarios, for example availability of weed seed late in the season, can be created by band spraying appropriately in GMHT sugar beet. Weeds can also help minimise insecticide use by reducing colonisation of beet by migratory insect pests, such as aphids, either offering alternative hosts, or providing olfactory and/or visual distraction (Dewar *et al.* 2000; Finch & Collier 2000). In addition, the avoidance of tractor hoeing in our GMHT

weed management systems also means that there is potential for improving the habitat for some species of ground nesting farmland birds.

Concerns have long been expressed about the effects of intensification of agriculture on the farmland environment, most recently by Donald *et al.* (2001). Some of the immediate environmental issues about GMHT are being addressed based on current agronomic practice in the Government's major Farm Scale Evaluations (FSE) project which examines one management option for GM crops (Firbank *et al.* 1999). Our work reported here examines a much wider range of weed management options, albeit on a small scale and on a single crop. Farmers could achieve higher yields with early overall applications of glyphosate than with conventional herbicides, but with the same low weediness as conventional herbicides; alternatively they could achieve equivalent yields to conventional herbicides with band sprays of glyphosate followed by late overall applications, but with the additional environmental benefits (insect food and habitat) from conservation of the weed flora for longer. This sets the FSE work in context as studying an environmental worst case option for GMHT crop production systems. In this paper we suggest that the way forward in row crops might be to use the technology to maximise environmental benefit and sustainability in a way that does not conflict with agronomic and financial benefits.

Acknowledgements

We thank Robert Hambridge, Paul Drinkwater, Colin Darlow for allowing trials on their farms, and Jan Wevers of IRS, Bergen op Zoom, Netherlands for Millieumeetlat system calculations. This work was partially funded by Monsanto plc. Rothamsted Research receives grant-aided support from the BBSRC.

REFERENCES

- Baars, M. A. 1979 Catches in pitfall traps in relation to mean densities of carabid beetles. *Oecologia* **41**, 25-26.
- Brants, I. & Harms, H. 1998 Herbicide tolerant sugar beet. *Proceedings of the 61st IIRB Congress, Brussels*, 195-204.
- Chamberlain, D. E., Fuller, R. J., Bunce, R. G. H., Duckworth, J. C. & Shrubbs, M. 2000 Changes in the abundance of farmland birds in relation to the timing of agricultural intensification in England and Wales. *Journal of Applied Ecology* **37**, 771-788.
- CLM. Centrum voor Landbouw en Milieu (Centre for Agriculture and Environment) 2000 Working Document: Milieumeetlat. Eds CLM Utrecht and IKC-Kerngroep MJPG, Ede. (supplements to working documents 1994, 1995 and 1997 and diskettes).
- Dewar, A. M., Haylock, L. A., Bean, K. M., May, M. J. 2000 Delayed control of weeds in glyphosate-tolerant sugar beet and the consequences on aphid infestation and yield. *Pest Management Science* **56(4)**, 345-350.
- Donald, P. F., Green, R. E. & Heath, M. F. 2001 Agricultural intensification and the collapse of Europe's farmland bird populations. *Proceedings of The Royal Society, Biological Sciences*, 7 January 2001 **268 (1462)**, 25-29.
- English Nature 1998 *Government Wildlife Advisor Urges Caution - Press release*. <http://www.english-nature.co.uk/news/story.asp?ID=139>, Peterborough.
- English Nature 2000 *Genetically modified organisms - position statement*. <http://www.english-nature.co.uk/news/statement.asp?ID=14>, Peterborough.
- Finch, S., Collier, R. H. 2000 Host-plant selection by insects – a theory based on 'appropriate/inappropriate landings' by pest insects of cruciferous plants. *Entomologia experimentalis et Applicata* **96**, 91-102.
- Firbank, L. G., Dewar, A. M., Hill, M. O., May, M. J., Perry, J. N., Rothery, P., Squire, G. R. & Woiwod, I. P. 1999 Farm-scale evaluation of GM crops explained. *Nature* **339**, 727-728.
- Hails, R.S. 2000 Genetically modified plants - the debate continues. *Trends in Ecology and Evolution* **15**, 14-18.
- Haughton, A. J., Bell, J. R., Boatman, N. D. & Wilcox, A. 1999 The effects of different rates of the herbicide glyphosate on spiders in arable field margins. *Journal of Arachnology* **27**, 249-254.
- Jansen, L. L. 1972 Extent and cost of weed control with herbicides and an evaluation of important weeds, 1968. ARS-H-1. Agricultural Research Service, US Department of Agriculture, Washington, DC, 227pp.
- Krebs, J. R., Wilson, J. D., Bradbury, R. B. & Siriwardena, G.M. 1999 The second silent spring? *Nature* **400**, 611-612.
- Kromp, B. 1999 Carabid beetles in sustainable agriculture: a review on pest control efficacy, cultivation impacts and enhancement. *Agriculture, Ecosystems & Environment* **74**, 187-228.
- Luff, M. L. & Woiwod, I. P. 1995 Insects as indicators of land-use change: a European perspective, focusing on moths and ground beetles. (In R. Harrington & N.E. Stork (eds) *Insects in a Changing Environment*). 399-422 Academic Press, London 576pp.
- Moll, S. 1997 Commercial experience and benefits from glyphosate tolerant crops. *Proceedings of the BCPC Conference – Weeds* **3**, 931-940.
- Moreby, S. J. & Southway, S. E. 1999 Influence of autumn applied herbicides on summer and autumn food available to birds in winter wheat fields in southern England. *Agriculture, Ecosystems and Environment* **72**, 285-297.

- Powell, W., Dean, G. J. & Dewar, A. M. 1985 The influence of weeds on polyphagous arthropod predators in winter wheat. *Crop Protection* **4**, 298-312.
- Purvis, G. & Curry, J. P. 1984 The influence of weeds and farmyard manure on the activity of Carabidae and other ground-dwelling arthropods in a sugar beet crop. *Journal of Applied Ecology* **21**, 271-283.
- Schweizer, E. E. & Dexter, A. G. 1987 Weed control in sugarbeets (*Beta vulgaris*) in North America. *Reviews of Weed Science* **3**, 113-33.
- Scott, R. K., Wilcockson, S. J. & Moisey, F. R. 1979 The effects of time of weed removal on growth and yield of sugar beet. *Journal of Agricultural Science, Cambridge* **93**, 693-709.
- Taylor, L. R., Kempton, R. A., Woiwod, I. P. 1976 Diversity statistics and the log-series model. *Journal of Animal Ecology* **45**, 255-271.
- Watkinson, A. R., Freckleton, R. P., Robinson, R. A. & Sutherland, W.J. 2000 Predictions of biodiversity responses to genetically modified herbicide-tolerant crops. *Science* **289**, 1554-1557.
- Wilson, J. D., Morris, A. J., Arroyo, B. E., Clark, S. E. & Bradbury, R.B. 1999 A review of the abundance and diversity of invertebrate and plant foods of granivorous birds in northern Europe in relation to agricultural change. *Agriculture Ecosystems and Environment* **75**, 13-30.
- Wilson, R. G. 1999 Response of nine sugarbeet (*Beta vulgaris*) cultivars to postemergence herbicide applications. *Weed Technology* **13**, 25-29.
- Wilson, R. G., Dean Yonts, C. & Smith, J. A. 2002 Influence of glyphosate and glufosinate on weed control and sugarbeet (*Beta vulgaris*) yield in herbicide-tolerant sugarbeet. *Weed Technology* **16**, 66-73.

Table 1 The effects of herbicide treatments on abundance and biodiversity of carabids, staphylinids and spiders caught (cumulative total) in pitfall traps in selected treatments at each of four sites.

Site	Sample weeks	Parameter	Treatment						Total (all tmts)
			Untreated	Conv	Glyph e+l	Glyph m+l	Glyph l+ll	Glyph eb+l	
1 1999	4	N	3346	2459	2353	2536	3131	-	12732
		No. of species	48	46	47	48	50	-	
		α index	7.94	8.03	8.32	8.40	8.45	-	9.80
		SE of α	1.25	1.30	1.34	1.33	1.31	-	1.26
6 1999	4	N	2752	2493	2402	2525	2646	-	12852
		No. of species	37	37	33	35	40	-	56
		α index	6.04	6.16	5.41	5.75	6.69	-	7.52
		SE	1.09	1.11	1.03	1.06	1.16	-	1.08
3 2000	9	N	3796	3403	3113	3690	3207	3528	31911
		No. of species	63	50	50	52	53	58	86
		α index	10.73	8.31	8.46	8.57	9.02	9.86	10.80
		SE of α	1.48	1.29	1.31	1.3	1.36	1.42	1.24
4 2000	7	No. of individuals	1694	894	755	796	943	781	9889
		No. of species	48	41	42	34	38	36	68
		α index	9.19	8.87	9.59	7.22	7.94	7.8	9.83
		SE of α	1.47	1.56	1.68	1.39	1.45	1.47	1.29

N = number of carabids + staphylinids + spiders per treatment (all 12 traps); SE = standard error; e = early; m = mid-timing; l = late; ll = later; b=band




Figure 1 Effects of the timing ($^{\circ}\text{Cd}$ after sowing) of the first overall sprays of glyphosate in a two spray regime (circles), and the second overall spray following a band spray (squares) on the cumulative biomass of weeds (left column figures) and sugar yield of sugar beet (right column figures) at five sites in 1999 and 2000. Data for conventional herbicides are given by orange circles.

Figure 2 Effect of weeds on abundance of arthropods in GMHT sugar beet

