

# Murgacashada Canqowga

## Ankle Sprain

The ankle is a joint where the bones of the legs and the feet meet. These bones are kept in place by ligaments.

A sprain occurs when the ligaments are suddenly stretched or torn.

Canqowga waa xubinta halka lafaha lugaha iyo cagta ay ku kulmaan. Lafahaan waxaa meel ku hayo seedaha.

Murgacashada waxay dhacdaa goortii seedaha ay si lama filaan ah ukala baxaan ama u jiidmaan.

### Signs

- Pain or tenderness
- Swelling
- Bruising
- Not able to move the ankle well

### Aastaamaha

- Xanuunka iyo danqashada
- Bararka
- Maruubka
- Uma suurtogaleysa inay si wanaagsan u dhaqaajiyaan canqowga

### Your care

Your doctor may want you to:

- Have an x-ray.
- Wear a splint or an ace wrap to support the injured ankle.
- Take over-the-counter pain medicine.
- Keep your foot raised above the level of your heart for the first 2 days. This will reduce swelling and pain.
- Put ice on the injured ankle for 15 to 20 minutes each hour for 1 to 2 days. Put the ice in a plastic bag and place a towel between the bag of ice and your skin.

### Daryeelkaaga

Dhaqtarkaaga wuxuu rabi karaa inaad:

- Raajo sameysid.
- Xiratid geedka kabka ama duubka taageero canqowga dhaawacan.
- Qaado daawada xanuunka dukaanka lagu iibiyo.
- Ku haay cagtaada inay ka koreyso heerka wadnahaaga ee 2 maalin ee ugu hooreeyo. Tani waxay yareyneysaa bararka iyo xanuunka.
- Saar baraf canqowga dhaawacan oo 15 illaa 20 daqiqo saacad walba oo 1 illaa 2 maalmood. Geli baraf bac oo saar shukumaan inta u dhaxeyso bacda iyo barafka iyo maqaarkaaga.

- Avoid putting weight on your foot for as long as your doctor tells you. Slowly begin putting weight on the injured ankle as you are able.
- Use crutches or a cane until you can stand on your ankle without having pain.

## **Call your doctor right away if your:**

- Bruising, swelling or pain gets worse
- Toes below the injury feel cold when you touch them, are numb or blue

**Talk to your doctor or nurse if you have any questions or concerns.**

- Iska ilaali inaad miisaan badan saarto cagtaada illaa xadka uu dhakhtarkaagu kuu sheego. Si tartiib ah u billoow inaad miisaankaaga saarto anqawga dhaawacan intaad awoodo.
- Isticmaal biraha ama gasaca illaa aadku joogsan kartid canqowgaaga adiga oo aan xanuun laheyn.

## **Soo wac dhaqtarkaaga isla markaas haddii:**

- Maruubkaaga, bararkaaga ama xanuunka ka sii daro
- Suulasha ka hoose dhaawaca waxay dareemaan qabow marka aad taabatid iyaga, waa kabuubasho iyo baluug

**La hadal dhaqtarkaaga ama kalkalisada haddii aad qabtid wax su'aalo ah ama walaacyo.**