

寶寶的聽力測試

Hearing Test for Your Baby

Babies learn to understand and speak words through hearing. If your baby has a hearing problem and it is not found, your baby will have a hard time learning words and how to talk. When hearing loss is found early, treatment can begin right away to prevent long-term problems.

Hearing Test

All babies have a hearing test before going home from the hospital. The test measures how your baby responds to sound. It takes about 10 minutes and can be done while he or she is sleeping.

- A small probe is placed in or near the baby's ear. It sends soft clicking sounds into the ear.
- Small pads may be put on your baby's head. These measure how the baby's brain responds to the sounds.

The Results

The results of the hearing test will be given to you and to your baby's doctor. If your baby does have hearing loss, you will get more information about treatment and resources, and more tests will be done.

If you have any questions about having your baby's hearing tested, talk to your baby's doctor or nurse.

嬰兒透過聽力，學會懂得辭語和說話。如果您的寶寶有聽力問題而未被發現，則可能很難學習辭語和說話。當聽力喪失能夠及早發現時，則可立即開始治療，以防止長期問題。

聽力測試

所有的寶寶在出院回家前都要做聽力測試。測試衡量您的寶寶對聲音的反應情況。測試約需10分鐘，並可在寶寶睡眠中進行。

- 測試人員將一個小探測儀放在寶寶耳朵內或耳朵附近，向耳朵內發出柔和的卡嗒聲。
- 測試人員可能將小墊子放在您的寶寶的頭上。這些墊子衡量寶寶的腦部對聲音的反應情況。

測試結果

您和您寶寶的醫生將獲得聽力測試的結果。如果您的寶寶確實聽力喪失，您將獲得治療和資源詳情，並將進行更多測試。

如果您對寶寶的聽力測試有任何疑問，請諮詢您寶寶的醫生或護士。

©2005 - February 1, 2021, Health Information Translations.

Unless otherwise stated, user may print or download information from www.healthinfotranslations.org for personal, non-commercial use only. The medical information found on this website should not be used in place of a consultation with your doctor or other health care provider. You should always seek the advice of your doctor or other qualified health care provider before you start or stop any treatment or with any questions you may have about a medical condition. The Ohio State University Wexner Medical Center, Mount Carmel Health System, OhioHealth and Nationwide Children's Hospital are not responsible for injuries or damages you may incur as a result of your stopping medical treatment or your failure to obtain treatment.