

COLD & FLU

MAX STRENGTH CAPSULES

paracetamol, caffeine, phenylephrine hydrochloride

Read all of this leaflet carefully before you take this medicine because it contains

important information you need to know.

This medicine is available without prescription,

however, you still need to use this product

carefully to get the best results from it.

Keep this leaflet as you may need to read it again

Ask your pharmacist if you need more

information or advice

I ■ If you get any side-effects, talk to your doctor or pharmacist. This includes any possible side-effects not listed in this leaflet.

You must talk to a doctor if you do not feel better or if you feel worse after 3 days

What is in this leaflet? I. What is Benylin Cold & Flu Max Strength Capsules and what is it

used for 12. What you need to know before you take Benylin Cold & Flu Max

Strength Capsules 3. How to take Benylin Cold & Flu Max Strength Capsules

4. Possible side-effects 5. How to store Benylin Cold & Flu **Max Strength Capsules** 6. Contents of the pack and other

information I What is Benylin Cold & Flu **Max Strength Capsules and** what is it used for

Benylin Cold & Flu Max Strength Capsules

relieve the symptoms of colds and flu, including headache, fever, nasal congestion, sore throat, fatigue and drowsiness, aches and pains. The capsules contain paracetamol, an analgesic

and antipyretic to relieve pain and fever; caffeine, a stimulant to help combat the tiredness associated with a cold and to help increase the

pain relief; and phenylephrine, a decongestant to relieve nasal congestion. 2 What you need to know before you take Benylin Cold

& Flu Max Strength Capsules This medicine is suitable for most people but a few people should not use it. If you are in any doubt, talk to your doctor or pharmacist. Do not use this medicine... If you are allergic to paracetamol,

> caffeine, phenylephrine or any of the other ingredients listed in section 6.

to any of the ingredients. If you have high blood pressure, severe heart disease, circulatory problems, or an

If you have ever had a bad reaction

overactive thyroid gland. If you have a history of stomach ulcers If you are taking, or have taken in the

last two weeks, drugs for depression known as Monoamine Oxidase Inhibitors (MAOIs).

If you are pregnant or breast-

feeding, unless advised by your doctor. If you are taking tricyclic antidepressants, such as imipramine, amitriptyline. In children under 16 years old.

A Talk to your doctor or pharmacist... If you suffer from liver or kidney

If you suffer from alcoholic liver disease or alcohol dependence If you have diabetes.

If you have a condition called Raynaud's Phenomenon (which results from

poor circulation in the fingers and toes). If you are taking **alcohol** or have alcohol dependence.

Other medicines and Benylin Cold & Flu Max Strength Capsules Tell your doctor or pharmacist if you are

taking, have recently taken or might take

any other medicines: colestyramine (used to lower blood cholesterol). metoclopramide or domperidone (used

to treat nausea and vomiting). tricyclic antidepressants (used to treat depression). barbiturates (a kind of sleeping tablet) such as phenobarbitone.

symbathomimetic drugs (used to treat congestion and asthma). medicines to treat migraines such as

ergotamine or methysergide. antihypertensives (drugs used to treat high

ephedrine or xylometazoline.

If you are not sure about any of the

medicines you are taking, show the pack

blood pressure such as beta blockers). anticoagulants (drugs used to thin the **blood** such as warfarin or coumarins). other decongestant drugs such as

to your pharmacist. If any of these bullet points apply, talk

to a doctor or pharmacist. A If you are pregnant or breast-feeding

Do not take this product if you are pregnant. Ask your doctor or pharmacist for advice before taking any medicine. Benylin Cold & Flu Max Strength

Capsules contains paracetamol. Do not take anything else containing paracetamol whilst

taking this medicine.

Check the table below to see how many capsules to take. For oral use only. Swallow whole with water. Do not chew. Do not take more medicine than the label tells you to. If you do not get better, talk to your doctor. Children and adolescents Do not give to children under 16 years. Adults, the elderly and children aged 16 years and over Age Dose Two capsules Adults, the elderly every 4 to 6 hours and children aged as required. 16 years and over Leave at least 4 to 6 hours between doses. Do not take more than eight capsules

Benylin Cold & Flu Max Strength

Do not drink alcohol (beer, wine,

spirits etc) while taking this product.

3 How to take Benylin Cold &

Flu Max Strength Capsules

Capsules with food, drink and alcohol

advised by your doctor. Taking this medicine with food and drink: You should avoid too much caffeine in drinks like coffee and tea. If you are also consuming other medicines or food that contains caffeine (coffee, tea, cola drinks, chocolate) you may have a high caffeine intake and might have difficulty sleeping, start shaking and/or have an uncomfortable feeling in the chest.

(4 doses) in any 24 hours period.

pharmacist.

If symptoms persist, talk to your doctor or

Do not take for more than 3 days unless

If anyone has too much Talk to a doctor at once if you take too much of this medicine even if you feel well. This is because too much paracetamol can cause delayed, serious liver damage. Go to your nearest hospital casualty department. Take your medicine and this leaflet with you.

If you forget to take the medicine

If you forget to take a dose, take the next

dose when needed provided that the last dose was taken at least 4 hours ago.

Do not take a double dose. 4 Possible side-effects

Benylin Cold & Flu Max Strength Capsules can have side-effects, like all medicines, although

these don't affect everyone and are usually mild and temporary. If you experience any of the following, stop using the medicine and tell your doctor or pharmacist: Unknown side-effects are:

bronchospasm

agitation

yellowing of the eyes and skin More rarely, the following side-effects can you may become more prone to bleeding, bruising, fever and infections, such as sore

fast, slow or irregular heart beat (palpitations)

throat and ulcers, due to changes in your blood. Very rare cases of serious skin reactions have

happen:

been reported.

Reporting of side-effects If you get any side-effects, talk to your doctor, pharmacist or nurse. This includes any possible side-effects not listed in this leaflet. You can also report side-effects directly via the Yellow Card Scheme at: www.mhra.gov.uk/yellowcard

Rare side-effects are:

headache, tremor, dizziness

difficulty sleeping (insomnia)

high blood pressure

or diarrhoea

glaucoma

allergic reactions such as skin rash nausea (feeling sick) or vomiting (being sick)

problems or pain passing urine

sudden weight loss, loss of appetite and

By reporting side-effects, you can help provide

more information on the safety of this medicine.

5 How to store Benylin Cold & Flu Max Strength Capsules Do not store this product above 25°C. Keep this medicine out of the sight and reach of children. Do not use your medicine after the date shown as an expiry date on the packaging.

6 Contents of the pack and other information What's in this medicine?

The active ingredients in Benylin Cold & Flu Max Strength Capsules are: Paracetamol 500 mg, caffeine 25 mg, and phenylephrine hydrochloride 6.1 mg. Other ingredients are: Maize starch,

magnesium stearate, talc, gelatin, quinoline yellow (E104), titanium dioxide (E171), Patent Blue V (E131) and erythrosine (E127). What the medicine looks like

croscarmellose sodium, sodium laurilsulfate,

Benylin Cold & Flu Max Strength Capsules are red/yellow capsules available in a 16 capsule pack. **Marketing Authorisation holder:**

Wrafton Laboratories Ltd., Wrafton, Braunton, Devon, EX33 2DL, UK. Manufacturer: Wrafton Laboratories Ltd., Wrafton, Braunton,

This leaflet was revised January 2017.

Benylin is a registered trade mark.

Devon, EX33 2DL, UK.

McNeil

restlessness, anxiety, nervousness, irritability, 17-0021 |5P73QM|I