

DRAMAMINE ORIGINAL FORMULA- dimenhydrinate tablet
Medtech Products Inc.

Disclaimer: Most OTC drugs are not reviewed and approved by FDA, however they may be marketed if they comply with applicable regulations and policies. FDA has not evaluated whether this product complies.

Dramamine Original Formula

Drug Facts

Active ingredient

(in each tablet)

Dimenhydrinate 50 mg

Purpose

Antiemetic

Use

for prevention and treatment of these symptoms associated with motion sickness:

- nausea
- vomiting
- dizziness

Warnings

Do not give to

children under 2 years of age unless directed by a doctor.

Ask a doctor before use if you have

- a breathing problem such as emphysema or chronic bronchitis
- glaucoma
- trouble urinating due to an enlarged prostate gland

Ask a doctor or pharmacist before use if you are taking sedatives or tranquilizers.

When using this product

- marked drowsiness may occur
- avoid alcoholic drinks
- alcohol, sedatives, and tranquilizers may increase dizziness

- be careful when driving a motor vehicle or operating machinery

If pregnant or breast-feeding,

ask a doctor before use.

Keep out of reach of children.

In case of overdose, get medical help or contact a Poison Control Center (1-800-222-1222) immediately.

Directions

- to prevent motion sickness, the first dose should be taken ½ to 1 hour before starting activity
- to prevent or treat motion sickness, see below:

Adults and children 12 years and over	<ul style="list-style-type: none"> • take 1 to 2 tables every 4-6 hours • do not take more than 8 tablets in 24 hours, or as directed by a doctor
Children 6 to under 12 years	<ul style="list-style-type: none"> • give ½ to 1 tablet every 6-8 hours • do not give more than 3 tablets in 24 hours, or as directed by a doctor
Children 2 to under 6 years	<ul style="list-style-type: none"> • give ½ tablet every 6-8 hours • do not give more than 1 ½ tablets in 24 hours, or as directed by a doctor

Other information

- store at room temperature 20-25°C (68-77°F)

Inactive ingredients

colloidal silicon dioxide, croscarmellose sodium, lactose, magnesium stearate, microcrystalline cellulose

Questions or comments?

Call **1-800-382-7219**

Dramamine.com

PRINCIPAL DISPLAY PANEL

Dramamine®
motion sickness
Original
Dimenhydrinate tablets/antiemetic
12 tablets (50 mg each)



Tamper Evident: Do not use if outer carton is open.

Important: Keep carton for future reference

Drug Facts		Drug Facts (continued)	
Active ingredient (in each tablet) Dimenhydrinate 50 mg	Purpose Antiemetic	Poison Control Center (1-800-222-1222) immediately.	
Use for prevention and treatment of these symptoms associated with motion sickness: ▪ nausea ▪ vomiting ▪ dizziness		Directions ▪ to prevent motion sickness, the first dose should be taken ½ to 1 hour before starting activity ▪ to prevent or treat motion sickness, see below:	
Warnings Do not give to children under 2 years of age unless directed by a doctor. Ask a doctor before use if you have ▪ a breathing problem such as emphysema or chronic bronchitis ▪ glaucoma ▪ trouble urinating due to an enlarged prostate gland Ask a doctor or pharmacist before use if you are taking sedatives or tranquilizers.		Adults and children 12 years and over ▪ take 1 to 2 tablets every 4-6 hours ▪ do not take more than 8 tablets in 24 hours, or as directed by a doctor	
When using this product ▪ marked drowsiness may occur ▪ avoid alcoholic drinks ▪ alcohol, sedatives, and tranquilizers may increase drowsiness ▪ be careful when driving a motor vehicle or operating machinery		Children 6 to under 12 years ▪ give ½ to 1 tablet every 6-8 hours ▪ do not give more than 3 tablets in 24 hours, or as directed by a doctor	
If pregnant or breast-feeding, ask a doctor before use. Keep out of reach of children. In case of overdose, get medical help or contact a		Children 2 to under 6 years ▪ give ¼ tablet every 6-8 hours ▪ do not give more than 1½ tablets in 24 hours, or as directed by a doctor	
		Other information store at room temperature 20°-25°C (68°-77°F)	
		Inactive ingredients colloidal silicon dioxide, croscarmellose sodium, lactose, magnesium stearate, microcrystalline cellulose	
		Questions? 1-800-382-7219 Dramamine.com	

DRAMAMINE ORIGINAL FORMULA

dimenhydrinate tablet

Product Information

Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:63029-901
Route of Administration	ORAL		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
DIMENHYDRINATE (UNII: JB937PER5C) (8-CHLOROTHEOPHYLLINE - UNII:GE2UA340FM)	DIMENHYDRINATE	50 mg

Inactive Ingredients

Ingredient Name	Strength
SILICON DIOXIDE (UNII: ETJ7Z6XBU4)	
CROSCARMELOSE SODIUM (UNII: M28OL1HH48)	
LACTOSE, UNSPECIFIED FORM (UNII: J2B2A4N98G)	
MAGNESIUM STEARATE (UNII: 70097M6I30)	
CELLULOSE, MICROCRYSTALLINE (UNII: OP1R32D61U)	

Product Characteristics

Color	orange	Score	2 pieces
Shape	ROUND	Size	8mm
Flavor	ORANGE	Imprint Code	
Contains			

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:63029-901-01	3 in 1 BOX	01/15/2012	
1		12 in 1 BLISTER PACK; Type 0: Not a Combination Product		
2	NDC:63029-901-02	1 in 1 BOX	01/15/2012	
2		12 in 1 BOTTLE; Type 0: Not a Combination Product		
3	NDC:63029-901-03	1 in 1 CARTON	01/15/2012	
3		2 in 1 POUCH; Type 0: Not a Combination Product		

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC monograph final	part336	01/15/2012	

Revised: 5/2023

Medtech Products Inc.