

**SENOKOT-S- standardized senna concentrate and docusate sodium tablet**  
**Atlantis Consumer Healthcare, Inc.**

-----  
**Senokot-S**  
**(standardized senna concentrate and docusate sodium)**

**Drug Facts**

**Active ingredients (in each tablet)**

Docusate sodium    50mg  
Sennosides            8.6mg

**Purpose**

Stool softener  
Laxative

**Uses**

- relieves occasional constipation (irregularity)
- generally produces a bowel movement in 6-12 hours

**Warnings**

**Do not use**

- if you are now taking mineral oil, unless directed by a doctor
- laxative products for longer than 1 week unless directed by a doctor

**Ask a doctor before use if you have**

- stomach pain
- nausea
- vomiting
- noticed a sudden change in bowel habits that continues over a period of 2 weeks

**Stop using and ask a doctor** if you have rectal bleeding or fail to have a bowel movement after use of a laxative. These may indicate a serious condition.

**If pregnant or breast-feeding,** ask a health professional before use.

## Keep out of reach of children.

In case of overdose, get medical help or contact a Poison Control Center rightaway.

### Directions

- take preferably at bedtime or as directed by a doctor

age	starting dosage	maximum dosage
adults and children 12 years of age and over	2 tablets once a day	4 tablets twice a day
children 6 to under 12 years	1 tablet once a day	2 tablets twice a day
children 2 to under 6 years	1/2 tablet once a day	1 tablet twice a day
children under 2 years	ask a doctor	ask a doctor

### Other information

- each tablet contains: **calcium 7 mg, sodium 4 mg VERYLOW SODIUM**
- store at 25°C (77°F); excursions permitted between 15°-30°C(59°-86°F)

### Inactive ingredients

**Inactive ingredients** croscarmellose sodium, dicalcium phosphate, FD&C Yellow #5 Lake\*, FD&C Yellow #6 Lake, hypromellose, magnesium stearate, maltodextrin, microcrystalline cellulose, mineral oil, polyethyleneglycol, sodium benzoate, sodium lauryl sulfate, starch, stearic acid, talc, titanium dioxide, triacetin

©2020

Avrio Health L.P.

A1023

Senokot-S 60 Tablets Label

NDC: 67618-310-60



Senokot-S 60 Tablets Leaflet  
NDC: 67618-310-60

Do not use if seal under cap is missing or damaged.

**Drug Facts**

Active Ingredients (in each tablet)	Purpose
Docusate sodium 50 mg	Stool softener
Sennosides 8.6 mg	Laxative

**Uses**

- relieves occasional constipation (irregularity)
- generally produces a bowel movement in 6–12 hours

**Warnings**

Do not use

- if you are now taking mineral oil, unless directed by a doctor
- laxative products for longer than 1 week unless directed by a doctor

Ask a doctor before use if you have

- stomach pain ■ nausea
- vomiting
- noticed a sudden change in bowel habits that continues over a period of 2 weeks

Stop use and ask a doctor if you have rectal bleeding or fail to have a bowel movement after use of a laxative. These may indicate a serious condition.

If pregnant or breast-feeding, ask a health professional before use.

Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away.

**Directions**

■ take preferably at bedtime or as directed by a doctor

age	starting dosage	maximum dosage
adults and children 12 years of age and over	2 tablets once a day	4 tablets twice a day
children 6 to under 12 years	1 tablet once a day	2 tablets twice a day
children 2 to under 6 years	½ tablet once a day	1 tablet twice a day
children under 2 years	ask a doctor	ask a doctor

**Other information**

■ each tablet contains: calcium 7 mg, sodium 4 mg VERY LOW SODIUM ■ store at 25°C (77°F); excursions permitted between 15°–30°C (59°–86°F)

Senokot-S 60 Tablets Carton  
NDC: 67618-310-60



Senokot-S 10 Tablets Carton  
 NDC: 67618-310-01



**Senokot-S** Effective, comfortable overnight constipation relief  
Standardized Senna Concentrate  
Docusate Sodium

**Drug Facts**

Active ingredient (in each tablet)	Purpose
Docusate sodium 50 mg	Stool softener
Sennosides 8.6 mg	Laxative

**Uses** • relieves occasional constipation (irregularity) • generally produces a bowel movement in 6 to 12 hours.

**Warnings**  
Do not use • if you are now taking mineral oil, unless directed by a doctor  
• laxative products for longer than 1 week unless directed by a doctor  
Ask a doctor before use if you have  
• stomach pain • nausea • vomiting • noticed a sudden change in bowel habits that continues over a period of 2 weeks  
Stop use and ask a doctor if you have rectal bleeding or fail to have a bowel movement after use of a laxative. These may indicate a serious condition.  
If pregnant or breast-feeding, ask a health professional before use.  
Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away.

**Directions**  
• take preferably at bedtime or as directed by a doctor

age	starting dosage	maximum dosage
adults and children 12 years and over	2 tablets once a day	4 tablets twice a day
children 6 to under 12 years	1 tablet once a day	2 tablets twice a day
children 2 to under 6 years	½ tablet once a day	1 tablet twice a day
children under 2 years	ask a doctor	ask a doctor

**Other information**  
• each tablet contains: calcium 7 mg, sodium 4 mg VERY LOW SODIUM  
• store at 25°C (77°F); excursions permitted between 15°-30°C (59°-86°F)

**Drug Facts (continued)**

**Inactive ingredients** croscarmellose sodium, dibasic calcium phosphate, FD&C Yellow #5 Lake<sup>®</sup>, FD&C Yellow #6 Lake, hypromellose, magnesium stearate, malic acid, microcrystalline cellulose, mineral oil, polyethylene glycol, sodium benzoate, sodium lauryl sulfate, starch, stearic acid, talc, titanium dioxide, triacetin

# SEKOKOT-S

standardized senna concentrate and docusate sodium tablet

## Product Information

<b>Product Type</b>	HUMAN OTC DRUG	<b>Item Code (Source)</b>	NDC:67618-310
<b>Route of Administration</b>	ORAL		

## Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
<b>SENNOSIDES</b> (UNII: 3FYP5M0JJX) (SENNOSIDES - UNII:3FYP5M0JJX)	SENNOSIDES	8.6 mg
<b>DOCUSATE SODIUM</b> (UNII: F05Q2T2JA0) (DOCUSATE - UNII:M7P27195AG)	DOCUSATE SODIUM	50 mg

## Inactive Ingredients

Ingredient Name	Strength
<b>CROSCARMELLOSE SODIUM</b> (UNII: M28OL1HH48)	
<b>CALCIUM PHOSPHATE, DIBASIC, ANHYDROUS</b> (UNII: L11K75P92J)	
<b>FD&amp;C YELLOW NO. 5</b> (UNII: I753WB2F1M)	
<b>FD&amp;C YELLOW NO. 6</b> (UNII: H77VEI93A8)	
<b>HYPROMELLOSE, UNSPECIFIED</b> (UNII: 3NXW29V3WO)	
<b>MAGNESIUM STEARATE</b> (UNII: 70097M6I30)	
<b>MALTODEXTRIN</b> (UNII: 7CVR7L4A2D)	
<b>CELLULOSE, MICROCRYSTALLINE</b> (UNII: OP1R32D61U)	
<b>MINERAL OIL</b> (UNII: T5L8T28FGP)	
<b>POLYETHYLENE GLYCOL, UNSPECIFIED</b> (UNII: 3WJQ0SDW1A)	
<b>SODIUM BENZOATE</b> (UNII: OJ245FE5EU)	
<b>SODIUM LAURYL SULFATE</b> (UNII: 368GB5141J)	
<b>STARCH, CORN</b> (UNII: O8232NY3SJ)	
<b>STEARIC ACID</b> (UNII: 4ELV7Z65AP)	
<b>TALC</b> (UNII: 7SEV7J4R1U)	
<b>TITANIUM DIOXIDE</b> (UNII: 15FIX9V2JP)	
<b>TRIACETIN</b> (UNII: XHX3C3X673)	

## Product Characteristics

<b>Color</b>	ORANGE	<b>Score</b>	no score
<b>Shape</b>	ROUND	<b>Size</b>	10mm
<b>Flavor</b>		<b>Imprint Code</b>	P
<b>Contains</b>			

## Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:67618-310-01	1 in 1 CARTON	10/01/1974	
1		10 in 1 BLISTER PACK; Type 0: Not a Combination		



1		Product		
2	NDC:67618-310-30	1 in 1 CARTON	10/01/1974	
2		30 in 1 BOTTLE, PLASTIC; Type 0: Not a Combination Product		
3	NDC:67618-310-60	1 in 1 CARTON	10/01/1974	
3		60 in 1 BOTTLE, PLASTIC; Type 0: Not a Combination Product		

## Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC Monograph Drug	M007	10/01/1974	

**Labeler** - Atlantis Consumer Healthcare, Inc. (118983925)

**Registrant** - Atlantis Consumer Healthcare, Inc. (118983925)

Revised: 12/2023

Atlantis Consumer Healthcare, Inc.