

**PREVALENCE OF POSTERIOR SEGMENT LESIONS IN DIABETES MELLITUS
USING USG B-SCAN**

Dr. Karuna Hazarika, Dr. Sima Malakar and Dr. Rishiraj Kalita*

Department of Radiology, Tezpur Medical College, Assam, India.

***Corresponding Author: Dr. Rishiraj Kalita**

Department of Radiology, Tezpur Medical College, Assam, India.

Article Received on 03/02/2023

Article Revised on 23/02/2023

Article Accepted on 14/03/2023

ABSTRACT

Ocular B-mode ultrasonography (US) has increasingly been useful in characterizing various ocular abnormalities. Though ophthalmoscopy is the most widely used and primary investigation in most of the posterior segment ocular lesions, the same can't be said all the time specially when the transparent media becomes opaque, mainly due to mature cataract or vitreous hemorrhages etc. Thus, US can guide the ophthalmologist in diagnosing disease and choosing treatment in presence of an opaque media. US is typically ideal for the eye due its superficial location and cystic configuration. Moreover, dynamic study helps distinguish between various conditions that would otherwise be difficult to differentiate in some clinical setting, such as vitreous, retinal, and choroidal detachment. The prevalence of visual impairment and blindness due to diabetic retinopathy (DR) and diabetic eye complications is on the rise worldwide and specially much more prevalent in the western world as a part of the metabolic syndrome. In diabetic patients with an opaque media due to any cause, US can be easily used to assess the prevalence of various ocular abnormalities mainly confined to the posterior segment. Thus, an understanding of the basic anatomy of the eye, the US technique, and common entities that affect the ocular globe will allow radiologists to offer this valuable imaging modality to patients and referring clinicians.

KEYWORDS: Opaque media Diabetes mellitus Diabetic retinopathy.

INTRODUCTION

Ultrasound is defined as sound that is above the range of human hearing (greater than 20 kHz). Discovery of modern ultrasound dates back to 1793, when it was observed by Lazzaro Spallanzani (Italy) that bats utilize echoes of their whistles to orient while flying in dark.^[1] The first use of ultrasound in ophthalmic diagnosis was reported in 1956 by Mundt and Hughes, who used industrial ultrasound equipment to examine enucleated normal eyes and eyes with intra-ocular tumors.^[2] The location and acoustic properties of the eyeball's contents makes it ideally suitable for ultrasound examination. The use of the intensity-modulated B-scan (Brightness modulation scan) in ophthalmic diagnosis was described by Baum and Greenwood in 1958.^[3] It is based on physical principles of pulse-echo technology and tissue acoustic impedance mismatch. It employs crystal transducers which produce ultrasonic pulses and retrieve echoes that are displayed after processing. When ophthalmoscopy is not possible, mainly due to opacification of the transparent media (e.g., mature cataract or vitreous hemorrhage), US can guide the ophthalmologist in diagnosing disease and choosing treatment. Recent research indicates that non-specialized eye scanners have a sensitivity of 93% and a specificity of 99%. Even in the absence of a specific eye scanner,

there is a strong association between sonographic data and clinic-pathological diagnosis when performed by qualified personnel.^[4] Compared to CT or MRI, which are unable to scan in real time and have poor spatial resolution, ultrasound makes a larger contribution to the tissue diagnosis.^[5] The transparent media often becomes opaque in diabetic patients and thus US in such circumstances can be quite valuable. Low vision due to DR occurs through a variety of mechanisms, including retinal detachment (RD), preretinal or vitreous hemorrhage (VH), associated neovascular glaucoma, and macular edema or capillary nonperfusion. The chronic hyperglycemia of diabetes is associated with long-term damage, dysfunction and failure of different organs, especially the eyes, kidneys, nerves, heart, and blood vessels.^[6] There is rapid rise in the prevalence of diabetes to 382 million individuals worldwide during the last 20 years and the expected rise to 592 million globally by 2030.^[7] people with diabetes have a higher risk of developing infections, cognitive impairment and dementia.^[8] Thus diagnosis, treatment and prevention of diabetes and its widespread micro and macrovascular complications is of utmost value. The present study is aimed to assess the prevalence of posterior segment ocular lesions both in clear and opaque media in diabetic patients.

Objective

1. Estimation of prevalence of posterior segment lesions in diabetes mellitus using USG B-scan.

MATERIALS AND METHOD

It is a hospital based prospective study. The study was conducted for duration of 6 months. The study was carried out in the department of radio diagnosis, Tezpur Medical College, Assam. Sample size of the study is according to standard statistical formula.

A group of selected diabetic patients was evaluated by ultrasonography B scan that will mostly come from Ophthalmology OPD. Before starting the ophthalmic ultrasound scanning, all patients underwent standard physical and ophthalmologic examinations where a detailed history was taken and eye examination, including slit lamp examination, visual acuity tests, pupillary reaction and fundoscopy were be done.

USG B-scan imaging

All ultra-sonography B scan imaging will be performed in a supine position. The diagnostic B scan sonography will be performed using a Samsung RS80A ultrasonic device, equipped with a high frequency (7.5-11MHZ)

direct contact linear transducer. For efficient and accurate diagnosis of ultrasound images, appropriate time gain compensation and dynamic range control of ultrasound echo signals will be automatically set by the system or manually adjusted by the sonographer to obtain the desired image quality on the screen.

Inclusion criteria

Patients with any type of diabetes mellitus (DM) (fasting serum glucose > 126 mg/dL on two independent determinations), with no previous history of ocular trauma or ocular surgery at any time and in the absence of any ocular infection or tumor and not undergoing insulin treatment.

Exclusion criteria


1. Patients with history of ocular trauma.
2. Patient with history of surgery.
3. Patients with associated orbital infection or orbital tumor.
4. Diabetic patients undergoing insulin treatment.


RESULTS AND OBSERVATION

The data has been analyzed and presented using appropriate methods.

1. Age distribution



Age	No. of cases	Percentage
0-10	00	00
10-20	00	00
20-30	00	00
30-40	02	5.8
40-50	04	11.4
50-60	11	20
60-70	11	31.4
70-80	07	31.4
total	35	100







2. Sex distribution

Sex	No. of cases	Percentage
Male	22	62.8
Female	13	37.2




3. Clinical presentation


Presenting symptoms	No. of abnormalities	Percentage
Blurred vision	17	48.6
Redness of eyes	12	34.3
Poor night vision	13	37.1
pain	06	17.1
Vision loss	07	20



4. Distribution of various ocular abnormalities


Ocular abnormalities	No. of abnormalities	Percentage
Vitreous	25	46.3
Lens	16	45.7
Retina	08	14.9
Choroid	03	5.5
Miscellaneous	02	3.7
Total	54	100






5. Laterality


Side affected	No. of cases	Percentage
Right eye	26	48.2
Left eye	16	29.6
Both eyes	12	22.2
total	54	100



6. Spectrum of lens abnormalities



Various abnormalities	no. of cases with lens abnormalities	percentage
Cataract	13	81.1
Dislocation	03	18.9
Subluxation	00	00
Others	00	00
Total	16	100






7. Spectrum of vitreous abnormalities

Vitreous abnormalities	No. of cases	Percentage
Vitreous degeneration	16	64
Vitreous hemorrhage	08	32
Vitreous detachment	01	04
Others	00	00
Total	16	100




8. Spectrum of retinal abnormalities

Various abnormalities	No. of cases	Percentage
Retinal detachment	04	50
Retinal hemorrhage	04	50
Total	08	100



9. Spectrum of choroidal abnormalities

Various abnormalities	No. of cases	Percentage
Choroidal detachment	02	66.7
Choroidal hemorrhage	01	33.3
Total	03	100



DISCUSSION

Ultrasonography has established itself as one of our most important diagnostic tools in the clinical practice of ophthalmology and has increased greatly our ability to detect and differentiate a wide variety of ocular and orbital disorders. Within the last decade, B-scan screening of opaque ocular media, particularly in eyes with cataract and vitreous hemorrhage as these are more common in diabetic patients, has constituted one of the most common indications for ocular ultrasound examination.

Ocular abnormalities are believed to be more common in males than in females. In a study done by OP Sharma the sex ratio was 2:1.^[9] In present study also male predominance was seen as 63% patients were males & 37% were females. The sex ratio was 1.7:1. In a study done by OP Sharma ocular abnormalities were observed maximum in 4th to 5th decades. In present study, maximum abnormalities were seen in 5th and 6th decade with 11 cases in each. In present study most common ocular abnormality detected was of vitreous with 25 cases. In a study by David Mcleod and Marie Restori 176 eyes were studied and maximum cases were of vitreous abnormalities.^[10] In another study by OP Sharma similar results were obtained. In the present study also maximum

cases were of vitreous abnormalities (46%). In another study by Jasmin Zvornicanin et al, maximum cases were of vitreous abnormalities (25,3%). In present study among vitreous abnormalities most abnormal eyes had vitreous hemorrhage (32%). Other vitreous lesion noted in abnormal eyes was vitreous detachment (4%). Study by David Mcleod and Marie Restori on 176 cases revealed increased prevalence of vitreous detachment (61%) than vitreous hemorrhage (57%). In our study also with diabetic patients, vitreous hemorrhage was the most common vitreous abnormality followed by vitreous detachment.

Cataract is the most encountered lens abnormality. In present study also cataract was the most common lens abnormality and 81% cases had cataract among total lens abnormalities followed by dislocation of lens which was 18.9%. Our study shows the prevalence of cataract in diabetic patients. In present study, retinal detachment and retinal hemorrhage cases were equal with 50% in each probably due to diabetic retinopathy induced proliferative retinopathy. studied. In a study by Lt col KK Sen and colleagues retinal detachment was the most common retinal pathology.^[11] In another study by Hassani and Bard retinal detachment was seen in 13.8% of total abnormalities.^[12] Choroidal abnormalities

included maximum cases of choroidal detachment (66.7%), while rest 33.3% cases were of choroidal hemorrhage.

CONCLUSION

Diagnosing and characterizing the abnormalities in diabetic patients with B-scan helps in preoperative cases as well as in the management and prevention of various diabetic ocular complications early and effectively. However, experience and understanding of the principles are essential for accurate diagnosis.

BIBLIOGRAPHY

1. Springer Handbook of Auditory Research [Internet].
2. S. Marchant. "The use of ultrasound," MIDIRS Essence, no. December, 2009, 1981; 443–446.
3. M. A. Qureshi and K. Laghari. "Role of B-scan ultrasonography in pre-operative cataract patients.," Int. J. Health Sci. (Qassim), 2010; 4, 1: 31–7.
4. Nzeh D, Owoeye J, Ademola-popoola D UI. Correlation of clinical and ultrasound findings in orbito-ocular disease using non-dedicated scanners: Experience at Ilorin, Nigeria. Eur J Sci Res, 2007; 16(3): 352–7.
5. Fielding JA. Ultrasound of the eye and orbit. In: Sutton D E. Textbook of radiology and imaging. 6th editio. London: Churchill Livingstone, 1998; 1551–71.
6. I. E. Mohamed, M. A. Mohamed, M. Yousef, M. Z. Mahmoud and BA. "Use of ophthalmic B-scan ultrasonography in determining the causes of low vision in patients with diabetic retinopathy," Eur. J. Radiol. Open, 2018; 5: 79–86.
7. International Diabetes Federation. IDF Diabetes Atlas.
8. Biessels GJ, Staekenborg S, Brunner E BC, ScheltensP. Risk of dementia in diabetes mellitus: a systematic review. Lancet Neurol, 2006; 5: 64-74.
9. A. AS. study of role of B. scan ultrasound in posterior segment pathology of eye. I. J. M. R. R, 2015; 3(9): 969-974.
10. Jasmin Zvornicanin1, Vahid Jusufovic1, Emir Cabric2, Zlatko Musanovic1, Edita Zvornicanin3 AP-B; Significance of Ultrasonography in Evaluation of Vitreo-retinal Pathologies; DOI: 10.5455/medarh.2012.66.3.
11. Lt Col KK Sen, Lt Col JKS Parihar, Maj Mandeep Saini SM BMR. Conventional B-mode Ultrasonography for Evaluation of Retinal Disorders. Med. J. Armed Forces India, 2003; 59: 310-312.
12. Aironi VD, Chougule SR SJ. Choroidal melanoma: A B-scan spectrum. Indian J. Radiol Imaging, 2007; 17: 8–10