



**EMBLICA OFFICINALIS (AMLA): A REVIEW FOR ITS PHYTOCHEMISTRY,
ETHNOMEDICINAL USES AND MEDICINAL POTENTIALS**

Alka Shukla*¹, Priyanka Pandey², S. Karmakar² and Wasim Raja²

¹Department of Biotechnology, AKS University, Sherganj, Satna (Madhya Pradesh).

²Central Laboratory Facility, Chhattisgarh Council of Science and Technology, Raipur, Chhattisgarh, India.

*Corresponding Author: Alka Shukla

Department of Biotechnology, AKS University, Sherganj, Satna (Madhya Pradesh).

Article Received on 28/02/2023

Article Revised on 20/03/2023

Article Accepted on 10/04/2023

ABSTRACT

The traditional Indian Ayurvedic medical system's most significant and widely researched plant, *Emblca Officinalis* (Amla), is discussed along with its preventive efficacies and safety. During an 16-week trial, 14 qualified healthy adult participants were randomly assigned to receive either amla or a placebo (400 mg daily). Vascular function, blood haematology, oxidative and inflammatory indicators, glucose and lipid profiles, urinalysis, and liver hepatotoxicity were the efficacy parameters examined. The main efficacy metric of blood fluidity significantly improved after Amla consumption. Along with a considerable increase in HDL cholesterol and a reduction in LDL cholesterol, benefits were also seen in the secondary endpoints, which included a decrease in von Willebrand factor (vWF), reduced 8-hydroxy-2'-deoxyguanosine (8-OHdG), and thrombin (TM), biomarkers of oxidative stress. After consuming amla, there were no appreciable differences in liver hepatotoxicity, urinalysis, or haematological as compared to baseline or placebo. After consuming amla, no negative side effects, changes in safety parameters, or problems with tolerance were noticed. As a result, Amla supplementation improved endothelial functioning, decreased oxidative stress, and had a tolerable palatability. The present review provided the more information regarding the scientific research and review of *Emblca Officinalis* (Amla).

KEYWORDS: After consuming amla, no negative side effects, changes in safety parameters, or problems with tolerance were noticed.

INTRODUCTION

Amla (*Emblca officinalis*) (EO) is revered in the ancient Indian medical system known as Ayurveda. Indian mythology holds that Amla, a member of the Euphorbiaceae family and also known as *Phyllanthus emblica* or Indian gooseberry, was the first tree to be created in the universe. Amla is a native of India and also grows in Pakistan, Uzbekistan, Sri Lanka, South East Asia, China, and Malaysia in tropical and subtropical climates.^[1] The fruits of the Amla tree are frequently employed in Ayurveda medicine and are thought to strengthen the body's resistance to illness.^[2] It plays a positive effect in the treatment of degenerative conditions such as cancer, diabetes, liver disease, ulcers, anaemia, and heart problems and is a crucial component of hepatoprotective formulations. Amla is incredibly nutrient-dense and one of the best sources of vitamin C, minerals, and amino acids.^[3] It has a variety of chemical components, including phenols, alkaloids, and tannins. Emblicanin A and B, gallic acid, and ellagic acid are two of the four hydrolysable tannins that have been reported to have biological activity. Ayurveda and traditional

medicine have both employed fruit as a potent rasayana to cure a variety of diseases, including diarrhoea, jaundice, inflammation, and several others.^[5] Nearly all portions also have therapeutic qualities. In the Indian medical system, amla fruit is frequently used as a diuretic, laxative, liver tonic, refrigerant, stomachic, restorative, anti-pyretic, hair tonic, and to avoid ulcer and dyspepsia. It is also used to treat fever and the common cold.

CLASSIFICATION

- **Kingdom:** Plantae
- **Division:** Angiospermae
- **Class:** Dicotyledonae
- **Order:** Geraniales
- **Family:** Euphorbiaceae
- **Genus:** *Emblca*
- **Species:** *officinalis* Geartn.
- **Vernacular names English:** Emblic myrobalan, Indian Goose berry
- **Sanskrit:** Aamalaki
- **Hindi:** Amla

MORPHOLOGY

The amla tree is a small to medium-sized deciduous tree that typically grows to a height of 8 to 18 metres. Its thin, light-grey bark exfoliates in tiny, uneven flakes, revealing the younger, lighter-colored surface beneath the older bark. The main stem is about 70 cm in diameter. The main trunk is typically split into 2 to 7 scaffolds very close to the base.^[22] The leaves are densely set in pinnate fashion and are 10–13 mm long

by 3 mm broad, giving the branches a fluffy look. Leaves form after the fruits have set. Flowers are unisexual, 4 to 5 mm long^[23], pale green in colour, and produced in clusters of 6 to 10 on the leaf axils. Fruits have a fleshy, almost globose shape, a diameter of 2.1–2.4 cm, a weight of 5.3–5.7 g, and a volume of 4.5–5.0 mL. The fruit's stone has six ribs and divides into three segments, each of which normally contains two seeds that are 4–5 mm long and 2–3 mm wide and weigh 572 to 590 mg.^[23-24]



Figure:- Amla(Emblica officinalis)

CHEMICAL CONSTITUENTS

Plants known for and extensively studied include amla. Research reveals that it contains a variety of biological elements, namely alkaloids, phenols, and tannins.^[23] Fruit contains about 28% of the plant's total tannic content. These tannins, which are antioxidants by nature, are found in two hydrolyzable forms: emblicanin A and emblicanin B.^[24] Emblicanin A hydrolyzes to produce ellagic acid, glucose, and gallic acid, whereas emblicanin B hydrolyzes to produce ellagic acid and glucose. Phyllembelin can also be found in this fruit.^[12] The additional fractionation revealed the presence of numerous other phytochemical components, including geraniin, corilagin, gallic acid, and furosin.^[25]

AMLA USE

Metabolic syndrome

A substantial amount of fructose-induced metabolic syndrome is present in the ethyl acetate-extracted extract of *E. officinalis*. According to this study's findings, *E. officinalis* has a significant amount of polyphenol.^[35]

Cardioprotective

Its primary advantage is protection from CVD, atherosclerosis, and other heart disorders, in addition to the other advantages. Only by reducing the oxidation of damaged or low-density lipoprotein (LDL) is an atherosclerosis cure conceivable. The Amla fruit's juice was confirmed to be high in polyphenol content. Also, the surgical pathological healing of the heart muscles ensured the *E. officinalis*' prophylactic activity. *E. officinalis* has heart-protective, antioxidant, and free radical scavenging activities, according to all the study and debate.^[36,37]

Diabetes and related complications

Foods consumed on a regular basis help to regulate the level of diabetes. Amla (*E. officinalis*), like garlic, onions, and turmeric, has been shown to reduce blood sugar levels. Around 2–3 g of *E. officinalis* powder effectively improves the amount of HDL cholesterol and regulates the level of LDL cholesterol. Moreover, Amla fruit is used by diabetic patients to prevent the development of neuropathy.^[38]

Immunostimulant

We are all familiar with a number of herbs that have immune-stimulating properties. Similar to this, ascorbic acid is best found in Amla, which increases immunoactivity (i.e., makes it two times more effective) by boosting immune cells and antibodies.^[39]

Antimicrobial

In America and the tropics, respectively, infectious diseases account for 50% and 20% of fatalities. Since more than a century ago, chemical components derived from medicinal plants have been employed to treat microbial infections.^[40] Amla (*E. officinalis*) extract in organic solvents (such as CHCl₃ and CH₃ OH) exhibits effective results against a small number of Gram-positive and Gram-negative bacteria.^[41] On the other hand, Gram-positive and Gram-negative bacteria were discussed by Vijayalakshmi *et al.* alongside the antibacterial properties of an aqueous extract of *E. officinalis* fruit pulp.^[42] Nonetheless, because of its antibacterial properties, *E. officinalis* medications will be used as affordable and secure medications in the future.

Anticancer

Because to the high concentration of polyphenol components in it, *E. officinalis* is superior for anticancer than other natural medicinal plants. Free radicals that damage skin are involved in the mechanisms of polyphenols' anticarcinogenic action, inflammatory response, and radiation resistance.^[43] Moreover, Amla (*E. officinalis*) is ideal for anti-aging and is used to make skin care cosmetics.^[48]

Eye disorders

E. officinalis and its tannoids are used to treat eye disease, which decreased the likelihood of oxidative stress as there was a reversal of changes with respect to lipid peroxidation, protein carbonyl content, and roles of antioxidant enzymes. A further defence against lens protein aggregation and insolubilization brought on by hyperglycemia was provided by amla.^[48]

CONCLUSION

Today, there is a renewed interest in studying the traditional medicinal herbs of India. Although the other medical systems are efficient, they also have a multitude of negative side effects that frequently result in life-threatening complications. Herbal medication alleviates all of these issues because it is natural. Amla, also known as *Emblica officinalis*, plays a significant role in Ayurveda, an ancient Indian medical system. Amla, which contains the maximum quantity of vitamin C and important nutrients, prevents a wide range of diseases due to its powerful antioxidant and biological capabilities. It can be applied to the nutraceuticals and biopharmaceuticals industries as well as as a potential food additive. Many studies have shown that different amla extracts and herbal formulations may have therapeutic benefits against a variety of disorders and have effects that are comparable to those of conventional medications. In this review, we attempted to establish the fundamental mechanisms behind the traditional and empirically supported usage of amla. Even though amla has been used for therapeutic purposes for many years, more research is urgently needed to prove its efficacy.

Research in medicinal plants has gained a renewed focus recently. The prime reason is that other system of medicine although effective come with a number of side effects that often lead to serious complications. Plant based system of medicine being natural does not pose this serious problems. Though *Emblica officinalis* has various medicinal applications, but it is the need of hour to explore its medicinal values at molecular level with help of various biotechnological tools and techniques. Further studies should be conducted to elucidate the molecular mechanism of interaction of various plant based drugs with human body in different diseases.

REFERENCES

1. Newman DJ, Cragg GM, Snader KM. The influence of natural products upon drug discovery. *Nat Prod Rep.*, 2000; 17: 215-34.
2. Cragg GM, Boyd MR, Khanna R, Kneller R, Mays TD, Mazan KD, et al. International collaboration in drug discovery and development: The NCI experience. *Pure Appl Chem.*, 1999; 71: 1619-33.
3. Sharma R, Thakur GS, Sanodiya BS, Savita A, Pandey M, Sharma A, et al. Therapeutic potential of *Calotropis procera*: A giant milkweed. *ISOR J Pharm Biol Sci.*, 2012; 4: 42-57.
4. Zhang LZ, Zhao WH, Guo YJ, Tu GZ, Lin S, Xin L. Studies on chemical constituents in fruits of Tibetan medicine *Phyllanthus emblica*. *Zhongguo Zhong Yao Za Zhi*, 2003; 28: 940-3.
5. Dasaroju S, Gottumukkala KM. Current trends in the research of *Emblica officinalis* (Amla): A pharmacological perspective. *Int J Pharm Sci Rev Res.*, 2014; 24: 150-9.
6. Dinesh M, Roopan SM, Selvaraj CI. Photocatalytic degradation of nitrophenol using biologically active *Phyllanthus emblica* seed extract. *J Photochem Photobiol B*, 2016; 161: 273-8.
7. Perianayagam JB, Sharma S, Joseph A, Christina A. Evaluation of anti-pyretic and analgesic activity of *Emblica officinalis* Gaertn. *J Ethnopharmacol*, 2004; 95: 83-5.
8. Nosal'ova G, Mokry J, Hassan KT. Antitussive activity of the fruit extract of *Emblica officinalis* Gaertn. (Euphorbiaceae). *Phytomedicine*, 2003; 10: 583-9.
9. Jeevangi S, Manjunath S, Sakhare PM. A study of anti-hyperlipidemia, hypolipidemic and anti-atherogenic activity of fruit of *Emblica officinalis* (Amla) in high fat fed albino rats. *Int J Med Res Health Sci.*, 2013; 1: 70-7.
10. Muruganandam A, Kumar V, Bhattacharya S. Effect of poly herbal formulation, EuMil, on chronic stress-induced homeostatic perturbations in rats. *Indian J Exp Biol*, 2002; 40: 1151-60.
11. Baliga M, Prabhu A, Prabhu D, Shivashankara A, Abraham A, Palatty P. Antidiabetic and cardioprotective effects of Amla (*Emblica officinalis*) and its phytochemicals: Preclinical observations. *J Ethnopharmacol*, 2013; 5: 583-600.
12. Chatterjee A, Chattopadhyay S, Bandyopadhyay SK. Biphasic effect of *Phyllanthus emblica* L. Extract on NSAID-induced ulcer: An antioxidative trail weaved with immunomodulatory effect. *Evid Based Complement Alternat Med*, 2011; 2011: 146808.
13. Yokozawa T, Kim HY, Kim HJ, Tanaka T, Sugino H, Okubo T, et al. Amla (*Emblica officinalis* Gaertn.) attenuates age-related renal dysfunction by oxidative stress. *J Agric Food Chem.*, 2007; 55: 7744-52.
14. Reddy VD, Padmavathi P, Kavitha G, Gopi S, Varadacharyulu N. *Emblica officinalis* ameliorates alcohol-induced brain mitochondrial dysfunction in rats. *J Med Food*, 2011; 14: 62-8.

15. Madhuri S, Pandey G, Khanna A. Oestrogen induced uterine damage in rats. *Toxicol Int.*, 2009; 16: 5-7.
16. Adil MD, Kaiser P, Satti NK, Zargar AM, Vishwakarma RA, Tasduq SA. Effect of *Embllica officinalis* (fruit) against UVB-induced photo-aging in human skin fibroblasts. *J Ethnopharmacol*, 2010; 132: 109-14.
17. Chularojmontri L, Suwatronnakorn M, Wattanapitayakul SK. *Phyllanthus emblica* L. Enhances human umbilical vein endothelial wound healing and sprouting. *Evid Based Complement Alternat Med*, 2013; 2013: 720728.
18. Sandhya T, Mishra K. Cytotoxic response of breast cancer cell lines, MCF 7 and T 47 D to triphala and its modification by antioxidants. *Cancer Lett.*, 2006; 238: 304-13.
19. Singh I, Sharma A, Jindal A, Soyad D, Goyal P. Fruit extract of *Embllica officinalis* (Amla) protects radiation induced biochemical lesions in the brain of Swiss albino mice. *Ann Neurosci*, 2010; 13: 65-71.
20. Ram MS, Neetu D, Yogesh B, Anju B, Dipti P, Pauline T, et al. Cyto-protective and immunomodulating properties of amla (*Embllica officinalis*) on lymphocytes: An in vitro study. *J Ethnopharmacol*, 2002; 81: 5-10.
21. Jain R, Pandey R, Mahant R, Rathore D. A review on medicinal importance of *Embllica officinalis*. *Int J Pharm Sci Res.*, 2015; 6: 72.
22. El-Desouky S, Ryu SY, Kim YK. A new cytotoxic acylated apigenin glucoside from *Phyllanthus emblica* L. *Nat Prod Res.*, 2008; 22: 91-5.
23. Rehman H, Yasin KA, Choudhary MA, Khaliq N, Rahman A, Choudhary MI, et al. Studies on the chemical constituents of *Phyllanthus emblica*. *Nat Prod Res.*, 2007; 21: 775-81.
24. Rajak S, Banerjee S, Sood S, Dinda A, Gupta Y, Gupta S, et al. *Embllica officinalis* causes myocardial adaptation and protects against oxidative stress in ischemic-reperfusion injury in rats. *Phytother Res.*, 2004; 18: 54-60.
25. Mirunalini S, Krishnaveni M. Therapeutic potential of *Phyllanthus emblica* (Amla): The ayurvedic wonder. *J Basic Clin Physiol Pharmacol*, 2010; 21: 93-105.
26. Variya BC, Bakrania AK, Patel SS. *Embllica officinalis* (Amla): A review for its phytochemistry, ethnomedicinal uses and medicinal potentials with respect to molecular mechanisms. *Pharmacol Res.*, 2016; 111: 180-200.
27. Ak T, Gülçin İ. Antioxidant and radical scavenging properties of curcumin. *Chem Biol Interact*, 2008; 174: 27-37.
28. Poltanov EA, Shikov AN, Dorman HD, Pozharitskaya ON, Makarov VG, Tikhonov VP, et al. Chemical and antioxidant evaluation of Indian gooseberry (*Embllica officinalis* Gaertn., syn. *Phyllanthus emblica* L.) supplements. *Phytother Res.*, 2009; 23: 1309-15.
29. Shivananjappa MM, Joshi MK. Influence of *Embllica officinalis* aqueous extract on growth and antioxidant defense system of human hepatoma cell line (HepG2). *Pharm Biol*, 2012; 50: 497-505.
30. Girish C, Pradhan SC. Indian herbal medicines in the treatment of liver diseases: Problems and promises. *Fundam Clin Pharmacol*, 2012; 26: 180-9.
31. Ghosh N, Ghosh R, Mandal V, Mandal SC. Recent advances in herbal medicine for treatment of liver diseases. *Pharm Biol*, 2011; 49: 970-88.
32. Golechha M, Bhatia J, Ojha S, Arya DS. Hydroalcoholic extract of *Embllica officinalis* protects against kainic acid-induced status epilepticus in rats: Evidence for an antioxidant, anti-inflammatory, and neuroprotective intervention. *Pharm Biol*, 2011; 49: 1128-36.
33. Dwivedi S, Aggarwal A. Indigenous drugs in ischemic heart disease in patients with diabetes. *J Altern Complement Med*, 2009; 15: 1215-21.
34. Yokozawa T, Kim HY, Kim HJ, Okubo T, Chu DC, Juneja LR. Amla (*Embllica officinalis* Gaertn.) prevents dyslipidaemia and oxidative stress in the ageing process. *Br J Nutr*, 2007; 97: 1187-95.
35. Kim HY, Okubo T, Juneja LR, Yokozawa T. The protective role of amla (*Embllica officinalis* Gaertn.) against fructose-induced metabolic syndrome in a rat model. *Br J Nutr*, 2010; 103: 502-12.
36. Patel SS, Goyal RK. Prevention of diabetes-induced myocardial dysfunction in rats using the juice of the *Embllica officinalis* fruit. *Exp Clin Cardiol*, 2011; 16: 87-91.
37. Zhao L, Zhang SL, Tao JY, Pang R, Jin F, Guo YJ, et al. Preliminary exploration on anti-inflammatory mechanism of corilagin (beta-1- O-galloyl-3, 6-(R)-hexahydroxydiphenoyl-D-glucose) in vitro. *Int Immunopharmacol*, 2008; 8: 1059-64.
38. Srinivasan K. Plant foods in the management of diabetes mellitus: Spices as beneficial antidiabetic food adjuncts. *Int J Food Sci Nutr*, 2005; 56: 399-414.
39. Kumar S, Gupta P, Sharma S, Kumar D. A review on immunostimulatory plants. *Zhong Xi Yi Jie He Xue Bao*, 2011; 9: 117-28.
40. Mahady GB, Huang Y, Doyle BJ, Locklear T. Natural products as antibacterial agents. In: *Studies in Natural Products Chemistry*. Vol. 35. Amsterdam: Elsevier, 2008; 423-44.
41. Rahman S, Akbor M, Howlader A, Jabbar A. Antimicrobial and cytotoxic activity of the alkaloids of amlaki (*Embllica officinalis*). *Pak J Biol Sci.*, 2009; 12: 1152-5.
42. Vijayalakshmi S, Arunkumar V, Anju D, Gunasundari P, Moorthy P, Chandrasekharan A. Comparative antimicrobial activities of *Embllica officinalis* and *Ocimum sanctum*. *Anc Sci Life*, 2007; 27: 1-6.
43. Priego S, Feddi F, Ferrer P, Mena S, Benlloch M, Ortega A, et al. Natural polyphenols facilitate elimination of HT-29 colorectal cancer xenografts by chemoradiotherapy: A Bcl-2-and superoxide

- dismutase 2-dependent mechanism. *Mol Cancer Ther.*, 2008; 7: 3330-42.
44. Penolazzi L, Lampronti I, Borgatti M, Khan MT, Zennaro M, Piva R, et al. Induction of apoptosis of human primary osteoclasts treated with extracts from the medicinal plant *Emblica officinalis*. *BMC Complement Altern Med*, 2008; 8: 59.
 45. Romano M, Vitaglione P, Sellitto S, D'Argenio G. Nutraceuticals for protection and healing of gastrointestinal mucosa. *Curr Med Chem.*, 2012; 19: 109-17.
 46. Mehmood MH, Siddiqi HS, Gilani AH. The antidiarrheal and spasmolytic activities of *Phyllanthus emblica* are mediated through dual blockade of muscarinic receptors and Ca²⁺ channels. *J Ethnopharmacol*, 2011; 133: 856-65.
 47. Baumann L, Woolery-Lloyd H, Friedman A. Natural ingredients in cosmetic dermatology. *J Drugs Dermatol*, 2009; 8: S5-9.
 48. Datta HS, Paramesh R. Trends in aging and skin care: Ayurvedic concepts. *J Ayurveda Integr Med*, 2010; 1: 110-3.
 49. Suryanarayana P, Saraswat M, Petrash JM, Reddy GB. *Emblica officinalis* and its enriched tannoids delay streptozotocin-induced diabetic cataract in rats. *Mol Vis*, 2007; 13: 1291.