



**ULTRASONIC VELOCITY, DENSITY MEASUREMENTS OF SUBSTITUTED  
HETEROCYCLIC DRUG IN 10% SODIUM CHLORIDE AT 303.15 K**

Samyekratna D. Kamble\*

\*Corresponding Author: Samyekratna D. Kamble

India

Article Received on 10/8/2023

Article Revised on 30/8/2023

Article Accepted on 20/9/2023

**ABSTRACT**

The acoustical properties have been investigated from the ultrasonic velocity and density measurements of Linezolid in 10% sodium chloride at 300.15K. The measurement have been done to evaluate acoustical parameter such as ultrasonic velocity, adiabatic compressibility (s), Partial molal volume (v), intermolecular free length (Lf), apparent molal compressibility (k), specific acoustic impedance (Z), relative association (RA), solvation number (Sn).

**KEY WORD:** - *adiabatic compressibility, apparent molal volume, Ultrasonic velocity.*

**INTRODUCTION**

In the recent years, measurements of the Ultrasonic velocity are helpful to interpreted solute-solvent, ion-solvent interaction in aqueous and non-aqueous medium.<sup>[1-6]</sup> Researchers have been studied the acoustical properties of complex in water. The author has studied the acoustical properties of four different drugs in methanol and she drawn conclusion from adiabatic compressibility. The four different drugs compress the solvent methanol to the same extent but it shows different solute-solvent interaction due to their different size, shape and structure.<sup>[6]</sup> Researcher studies the different acoustical properties of some substituted Pyrazolines in binary mixture acetone-water and observed variation of ultrasonic velocity with concentration.<sup>[7]</sup> Authors have investigated the measurement of ultrasonic velocity and density of amino acid in aqueous magnesium acetate at constant temperature.<sup>[8]</sup> The ion-dipole interaction mainly depends on ion size and polarity of solvent. The strength of ion-dipole attraction is directly proportional to the size of the ions, magnitude of dipole. But inversely proportional to the distance between ion and molecules. Voleisines has been studied the structural properties of solution of lanthanide salt by measuring ultrasonic velocity.<sup>[9]</sup> The investigators has been studied the ultrasonic velocity of PEG-8000, PEG- study of acoustical properties of substituted heterocyclic compounds under suitable condition.<sup>[10]</sup> Authors have studied the acoustical and thermodynamic properties of citric acid in water at different temperature.<sup>[11]</sup> Mishra et.al. have investigated ultrasonic velocity and density in non-aqueous solution of metal complex and evaluate acoustic properties of metal complex.<sup>[12-15]</sup>

After review of literature survey the detail study of substituted heterocyclic drug under identical set of experimental condition is still lacking. It was thought of interest to study the acoustical properties of substituted drug under suitable condition.

**EXPERIMENTAL AND METHODS**

In the present study, the drug linezolid was used. The double distilled solvent was used for preparation of different concentration of drug solution. The densities were determined by using specific gravity bottle by relative measurement method with accuracy  $1 \times 10^{-5}$  gm/cm<sup>3</sup>. The ultrasonic velocities were measured by using ultrasonic interferometer having frequency 3MHz. The constant temperature was maintained by circulating water through the double wall measuring cell, made up of steel.

In the present investigation, different properties such as adiabatic compressibility (s), apparent molal volume (v), intermolecular free length (Lf), apparent molal compressibility ( $\Phi_k$ ), specific acoustic impedance (Z), relative association (RA), solvation number(Sn), limiting apparent molal compressibility ( $\Phi_{k0}$ ), limiting apparent molal volume(v) and their constant (Sk, Sv).

**Theory**

Adiabatic compressibility ( $\beta_s$ ) is given by:

$$\text{Adiabatic compressibility}(\beta_o) = \frac{1}{\rho_o^2 d_o}$$

$$\text{Adiabatic compressibility}(\beta_s) = \frac{1}{\rho_s^2 d_s}$$

$$\text{Apparent molal volume}(\phi_v) = \left(\frac{M}{d_s}\right) \times \frac{(d_o - d_s) \times 10^3}{m \times d_s \times d_o}$$

$$\text{Apparent molal compressibility}(\phi_k) = 1000 \times \frac{(\beta_s d_o - \beta_o d_s) \times 10^3}{m \times d_s \times d_o} + \frac{\beta_s M}{d_o}$$

$$\text{Specific acoustic impedance (Z)} = \rho_s d_s$$

$$\text{Intermolecular free length (L}_f) = K\sqrt{\rho d}$$

$$\text{Solvation number (S}_n) = \frac{\phi_k}{\beta_o \left(\frac{M}{d_o}\right)}$$

**RESULTS AND DISCUSSION**

In the present investigation, different acoustical properties such as ultrasonic velocity, adiabatic compressibility, intermolecular free length, specific acoustic impedance, are listed in table-1. Partial molal volume, apparent molal compressibility, relative association, solvation number are listed in table-2. It was found that the ultrasonic velocity decreased with the increase in concentration for system (Table-1). Variation of ultrasonic velocity in solution depends upon the increase or decrease of molecular free length after mixing the component. This is based on a model for sound propagation proposed by Eyring and Kincaid.<sup>[13]</sup> Intermolecular free length increased linearly on increase in concentration of substituted drug. Hence, decreased in ultrasonic velocity with increase in concentration of drug. It happened because there was significant interaction between ions and solvent molecules suggesting a structure promoting behavior of the added electrolyte. The specific acoustic impedance decreased with the increase in concentration of drug. When concentration of electrolyte was increased, the thickness of oppositely charged ionic atmosphere increases due to decrease in ionic strength. This is suggested by decrease in acoustic impedance with increase in concentration in system. It was seen that the intermolecular free length

increased with the increase in concentration in system. The intermolecular free length increased due to greater force of attraction between solute and solvent by forming hydrogen bonding. The adiabatic compressibility increased with the increase in concentration of solution. It happened due to collection of solvent molecule around ions, this supporting weak ion-solvent interaction. This indicates that there is significant solute-solvent interaction.



It was observed that apparent molal volume increased with concentration in system. It indicates the existence of strong ion-solvent interaction. It was found that the value of apparent adiabatic compressibility was increased with the increase in concentration of drug. It shows strong electrostatic attractive force in the vicinity of ions. From the data, we were concluded that strong molecular association was found in drug. The value of relative association increased with the increase in concentration in system. It has been found that there was strong interaction between solute and solvent. There were regular increases in solvation number with the increase in concentration; it indicates the solvent molecule forms strong coordination bond in primary layer. It indicates the increase in size of secondary layer of Solvation.

**Table-1** Ultrasonic velocity, density, adiabatic compressibility (S), Specific acoustic impedance (Z) Intermolecular free length (L<sub>f</sub>).

Conc. Moles lit-1 (m)	Density (d <sub>s</sub> ) kg m <sup>-3</sup>	Ultrasonic velocity (U <sub>s</sub> ) m s <sup>-1</sup>	Adiabatic compressibility (β <sub>s</sub> ) ×10 <sup>-10</sup> m <sup>2</sup> N <sup>-1</sup>	Intermolecular free length (L <sub>f</sub> ) ×10 <sup>-11</sup> m	Specific acoustic impedance (Z×10 <sup>6</sup> ) kg m <sup>-2</sup> s <sup>-1</sup>
1×10 <sup>-3</sup>	1019.94	1554.93	4.0551	4.0500	1.5859
2×10 <sup>-3</sup>	1020.24	1549.65	4.081	4.0632	1.5810
3×10 <sup>-3</sup>	1021.53	1541.09	4.1219	4.0832	1.5743
4×10 <sup>-3</sup>	1021.84	1523.89	4.2142	4.1287	1.5572
5×10 <sup>-3</sup>	1022.02	1509.82	4.2923	4.1668	1.5431
6×10 <sup>-3</sup>	1022.39	1487.56	4.4201	4.2284	1.5209
7×10 <sup>-3</sup>	1022.51	1473.36	4.5052	4.2689	1.5065
8×10 <sup>-3</sup>	1022.74	1469.58	4.5274	4.2794	1.5030
9×10 <sup>-3</sup>	1022.89	1442.59	4.5275	4.2905	1.4756

**Table-2: apparent molal compressibility ( $\beta_{\text{app}}$ ), Relative association ( $R_A$ ), Apparent molal volume ( $V_{\text{app}}$ ), Solvation number ( $S_n$ ) –**

Conc. Moles lit-1 (m)	Apparent molal volume ( $V_{\text{app}}$ ) $\text{m}^3 \text{mole}^{-1}$	Apparent molal compressibility $\beta_{\text{app}}$ $\times 10^6 \text{m}^2 \text{N}^{-1}$	Relative association ( $R_A$ )	Solvation number ( $S_n$ )
1x10 <sup>-3</sup>	4.4799	1.341	1.0010	9.7893
2x10 <sup>-3</sup>	2.2188	1.349	0.9952	9.8456
3x10 <sup>-3</sup>	1.7820	1.361	0.9936	9.9332
4x10 <sup>-3</sup>	1.3283	1.391	0.9911	10.1522
5x10 <sup>-3</sup>	1.0311	1.417	0.9833	10.3420
6x10 <sup>-3</sup>	0.8634	1.459	0.9770	106485
7x10 <sup>-3</sup>	0.7094	1.487	0.9700	108529
8x10 <sup>-3</sup>	0.6070	1.494	0.9679	109039
9x10 <sup>-3</sup>	0.5189	1.494	0.9637	109039

**CONCLUSION**

The present study shows the experimental data for ultrasonic velocity and density at 300.15K for substituted heterocyclic drug in 10% sodium chloride. From experimental data, the acoustical properties were calculated. The solute-solvent interaction and ion-ion / solute-solute interaction existing between drug and solvent were also studied with the help of experimental data. Lastly it has been concluded from the experimental data, that the solute-solvent interaction in drug-solvent systems are weak.

**REFERENCES**

- S. Baluja and S. Oza, *Fluid phase equilibria.*, 2005; 200(1): 49-54.
- M. K. Rawat and Sangeeta, *Ind. J. pure Appl. Phy.*, 2008; 46: 187-192.
- A Ali. and A K Nain, *Acoustics Lett.*, 1996; 19: 53.
- H. Ogawa and S J Murakami, *J. Solution. Chem.*, 1987; 16: 315.
- D. Ubagaramary, Dr.P.Neeraja, *J. of Applied Chemistry.*, 2012; 2(5): 01-19.
- D. V. Jahagirdar , B. R. Arbad , S. R. Mirgane, M. K. Lande and A. G. Shankarvar , *J.Molecular Liq.*, 1998; 75: 33-43.
- Y. K. Meshram and M. L. Narwade , *Acta Ciencia Indica*, 2001,XXVII.C No.2 : 67-70.
- R. Palani and S. Saravanan, *Research J. Phy.*, 2008; 2(1): 13-21.
- B. Voleisiene and A. Voleisis, *J. Ultrasound*, 2008; 63(4): 7-18.
- V. K. Syal, A.Chauhan and S. Chauhan, *J. Pure Ultrasound.*, 2005; 27: 61-69.
- A. Tadkalkar, P.Pawar and G.K. Bichile, *J.Chem. Pharm.Res.*, 2011; 3(3): 165.
- A.P. Mishra and D.K. Mishra, *J.Chem. Pharm.Res.*, 2011; 3(3): 489.
- M. Arvinthraj, S. Venktesan and D. Meera, *J.Chem. Pharm.Res.*, 2011; 3(2): 623.
- S.K. Thakur and S.Chauhan, *J.Chem. Pharm.Res.*, 2011; 3(2): 657.
- Shilpa A. Mirikar, Pravina P. Pawar, Govind K. Bichile, *American Journal of Pharmacology and Pharmacotherapeutics*, 2015; 2(1): 2-7.