



STUDIES OF STABILITY CONSTANT OF MIXED LIGAND WITH TRANSITION METAL IONS AT 303.15 K.

J. P. Nehete*, R. D. Talele¹ and A. N. Sonar²

*S.V.P. College, Ainpur.

¹D.N. College, Faizpur.

²Shri. V. S. Naik College, Raver.



*Corresponding Author: J. P. Nehete

S.V.P. College, Ainpur.

Article Received on 25/06/2024

Article Revised on 15/07/2024

Article Accepted on 04/08/2024

ABSTRACT

The interaction of transition metal ion with mixed amino acids has been investigated by pH metric titration at 0.02 M ionic strength at 303.15K in 60 % Ethanol-Water mixture. The data obtained use to estimate the values of proton-ligand stability constant (P_k) and Metal -ligand stability constant (logK). It is observed that transition metal ion form 1:1, 1: 2 complexes with all the systems.

INTRODUCTION

The studies of metal-ligand complexes in solution of a number of metal ions with carboxylic acids, oximes, phenols etc. would be interesting which throw light on the mode of storage and transport metal ion in the biological kingdom.

In the earlier papers extensive data base on metal complexes with substituted heterocyclic drugs was presented. Narwade et.al.^[1] studies the Formation and Stability constant of thorium (IV) complex with some substituted pyrazolines. Mathieu W.A. Steenland et.al.^[2] studies stability constant of Cu (II) and Ni(II) complexes of trans -dioxopentaaza macrocycles in aqueous solution by different technique. Hong-Wen Gao et.al.^[3] has studied the stability constant of Cu (II) and Co (II) complexes with CNBAC in water sample spectrophotometrically. Tuba Sismanoglu^[4] have studied the stability constant of binary complexes of Nicotinamide with Mn (II) by pH metrically. He also determines change in free energy, change in enthalpy and change in entropy from stability constant at different temperature. Tekade et.al.^[5] have been studied complex formation of Cu (II) and Co (II) metal ion complex with substituted isoxazolines. O.Yamauchi et.al.^[6] studied stability constant of metal complexes amino acids with charged side chain by pH-metrically. Hayati Sari et.al.^[7] studied the stability constant of glyoxime derivative and their Nickel, Copper, Cobalt and Zinc complexes potentiometric and theoretically. Jaime M. Murphy^[10] et.al have been studied the stability constant of redox-active metals with amino acids. Naciye Türkel studied the

Stability Constants of Mixed Ligand Complexes of Nickel (II) with Adenine and Some Amino Acids.^[11]

After review of literature survey the detail study of complex under identical set of experimental condition is still lacking. It was thought of interest to study the chelating properties of mixed amino acids under suitable condition with metal ions by pH metrically.

MATERIAL AND METHOD

pH measurement were carried out with equip-tronic EQ-610pH meter (accuracy ± 0.01 units) using combine glass electrode at room temperature. Metal ions solution were prepared in triply distill water and concentration estimated by standard method.^[8] The solution of drug prepared in solvent. The pH metric reading in 50% ethanol – water mixture were converted to [H⁺] value by applying the correction proposed by Van Uitert Haas.


The overall ionic strength of solution was constant and calculated by the equation

$$\mu = 1/2 \sum X_i Z_i^2$$

The concentration of other ion in addition to Na⁺ and ClO₄⁻ were also taken into consideration.

RESULT AND DISCUSSION

Substituted heterocyclic drugs may be ionized as acid having replaceable H⁺ ion from -OH group. Therefore it is represented as HL i.e.



The titration data used to construct the curves between volume of NaOH and P^H . They are called acid-ligand titration curves.

It is observed from titration curves for all systems ligand start deviating from the free acid curves at P^H 2.5 and deviating continuously up to P^H =11. The deviation shows that dissociation of proton in substituted drugs. The average number of proton associated with the ligand (n_A) was determined from free acid and acid – ligand titration curves employing the equation of Irving and Rossotti.^[9] The P^k values were determined from formation curves ($n_A \text{ vs } P^H$) by noting the P^H at which $n_A=0.5$. The accurate values of pK were calculated by point wise calculations which are presented in table -1. The pK values are found.

Table 1: Proton ligand stability constant of ligand at 0.1M ionic strength.

System	Constant pK	
	Half integral	Point wise calculation
Ligand-1	6.52	6.590. $\pm$ 0.04

METAL -LIGAND STABILITY CONSTANT (Log k):-

Metal-ligand stability constant of transition metal ion chelate with drug were determined by employing Bjerrum calvin P^H metric titration method as adopted by Irving and Rossotti. The formation of chelate between transition metal ion with drug was indicated by the significant separation starting from pH =2.5 for transition metal ion with ligand -1.

Table 2: Determination of metal –ligand stability constant (log K) of transition metal ion with drug at 0.1M ionic strength.

System	$\text{Log}k_1$	$\text{Log}k_2$	$\text{Log}k_2 - \text{Log}k_1$	$\text{Log}k_2 / \text{Log}k_1$
Cu (II)-Ligand	3.55	5.35	1.80	1.0507
Fe (II)-Ligand	3.72	5.46	1.74	1.0677
Ni (II)-Ligand	3.65	5.57	1.92	1.5260
Mg (II)-Ligand	3.25	4.55	1.30	1.4000
Co (II)-Ligand	4.05	5.72	1.67	1.4124

The result shows the ratio of $\text{Log}k_2 / \text{Log}k_1$ is positive in all cases. This implies that there is little or no steric hindrance to the addition of secondary ligand molecule. The smaller difference may be due to trans structure.

REFERENCES

1. M. L. Narwade, V.S. Jamode, and S.K. Gudadhe, Stability constant of Co (II), Ni(II) and Cu (II) chelate with some substituted Pyrazolines, *Acta. CienciaIndica*, 1985; **XIc**: 234.
2. Mathieu W.A. steenland, Use dirack Gerrit G. turnan, Bart Devreese, Werner Lippens, Jozef Van Beeuman and Andre M. Goeminne, Potentiometric and spectrophotometric study of Cu(II) and Ni(II) complexes of trans –dioxopentaaza macrocycles in aqueous solution, *J.Chem. Soc. Dalton trans*, 1997; 3637-3647.
3. Hong Wen Gao and Jian-Fu-Zhao, Stability constant of Co(II) and Cu(II) complexes with CNBAC and selective determination of Cu(II) by completion coordination, *Croatica Chemica Acta*, CCACAA, 2003; 76(1): 1-6.
4. Tuba sismanoglu, Thermodynamic of stability constant of binary complex of Nicotinamide with Mn (II), *Chinese Chemical Letters*, 2003; 14(11): 1207-1210.
5. P.V. Tekade, K. N. Patil, M.L. Narwade, P.S. Bodake, Y.K. Meshram, Complex formation of Co (II) and Cu (II) metal ion complexes with substitutes isoxazolines. *Asian J. Chem.*, 2006; 18: 2657.
6. O. Yamauchi and A. Odani, Stability constant of metal complexes of amino acids with charged side chain, *Pure and Applied chemistry*, 1996; 68(2): 469-496.
7. Hayati Sari, Muzaffer Can and Mustafa macit, Potentiometric and theoretical studies of stability constant of glyoxime derivative and their Nickel, Copper, Cobalt and Zinc complexes, *Acta Chim Slov.*, 52: 317-322.
8. A.I. Vogel, Longmans Green A Text Book of Quantitative Inorganic Analysis, London, 1975: 589.
9. H.S. Irving and H.S. Rossotti, J. Chem. Society, 1954; 2904-2913.
10. Jaime M. Murphy, Brian A. Powell, Julia L. Brumaghim, *coordination chemistry review*, June 2020; 412: 1213253.
11. Naciye Türkel, Hindawi Publishing Corporation *Bioinorganic Chemistry and Applications*, 2015; 1-9.