



ETHNOPHARMACOLOGICAL STUDY OF ANTIHYPERLIPIDEMIC IN THE SUKASARI REGION, SUBANG, WEST JAVA, INDONESIA

**Zafira Fatiha Nurulhadi, Amira Ramadani, Anita, Aprilia Kuswanti, Della Putri Valentina,
Erna Sri Rahmawati, Ferdyan Pranata Saputra, Fitri Nur Nathania, Helsen, Iin Kurniawati, Irdiyani Fariha,
Renita Hamjah, Richa Putri Immelia, Silvia Ismayanti, Siti Mudrikah, Tanti Amelia and Maulana Yusuf
Alkandari***

Faculty of Pharmacy, Universitas Buana Perjuangan Karawang, Karawang, West Java, Indonesia.



***Corresponding Author: Maulana Yusuf Alkandari**

Faculty of Pharmacy, Universitas Buana Perjuangan Karawang, Karawang, West Java, Indonesia.

Article Received on 22/09/2024

Article Revised on 12/10/2024

Article Accepted on 01/11/2024

ABSTRACT

Hyperlipidemia is a secondary metabolic dysregulation related to increased levels of triglycerides, cholesterol and LDL in serum, which is a major risk factor for premature cardiovascular disease such as atherosclerosis, hypertension, coronary heart disease, etc. This research aims to document and preserve the use of ethnomedicinal to treat hyperlipidemia by communities in the Sukasari Region, Subang, West Java, Indonesia. Fieldwork was carried out from March to April 2024 using direct interviews, questionnaires, and discussions. Plant species are identified based on standard taxonomic methods, flower morphological characteristics, and where possible, using samples for comparison, as well as consultation with experts and the literature. The plant types obtained were grouped into families according to the Cronquist classification system. Plant names were checked against the Plant List (www.plantlist.org) and the International Plant Name Index (www.ipni.org). This research reports that 30 plant species are commonly used by people in the Sukasari Region to treat hyperlipidemia. Among the various plant parts used, leaves (50.0%) are most frequently used in making medicines, followed by rhizomes (13.3%), fruit (13.3%), flowers, stem, and seed (respectively 6.7%) and rind (3.3%). Meanwhile, the most frequently used preparation methods were decoction (76.7%) and infusion (23.3%). The results of this research confirm that people in the Sukasari Region still rely heavily on medicinal plants for their health care system, especially for the treatment of hyperlipidemia with the most frequently used parts of the leaves and their use in decoctions and infusions.

KEYWORDS: Traditional medicine, Ethnomedicinal plants, Sukasari Region, Antihyperlipidemic.

INTRODUCTION

Hyperlipidemia is a predictor of coronary artery disease (CAD). The prevalence of this disease is quite high and increasing in both developed and developing countries in the world.^[1] Hyperlipidemia is an important risk factor in the initiation and progression of atherosclerosis. The main manifestations of this disorder include increased plasma concentrations of total cholesterol (TC), triglycerides (TG), low-density lipoprotein cholesterol (LDL-C), and low concentrations of high-density lipoprotein cholesterol (HDL-C).^[2] Therefore, the main consideration in the therapy of hyperlipidemia and arteriosclerosis is to reduce the increase in blood serum/plasma lipid levels.^[3] Currently available hypolipidemic drugs have been reported to have several worrying side effects including hyperuricemia, muscle damage, impotence, memory loss, peripheral neuropathy, body aches, gynecomastia, skin rashes, and others. In addition, there is also an increased risk of myopathy and

rhabdomyolysis which commonly occurs when used in combination with other drugs.^[4] Most hypolipidemic drugs can be effective if used for several weeks but may result in worsening side effects such as liver damage.^[5] Therefore, it is necessary to search for new antihyperlipidemic agents derived from natural ingredients in the form of medicinal plants.

Medicinal plants have been widely used in traditional practices by different ethnic populations throughout the world for the prevention and/or treatment of several chronic diseases. Despite the development of newer technologies and advances in modern medicine, most of the world's population still relies on traditional systems of medicine to fulfill their medical care, especially in Indonesia.^[6] One of the Region in Indonesia that still uses herbal plants as an alternative treatment, especially to treat hyperlipidemia, is the Sukasari Region. This research aims to obtain detailed information about the

use of herbal plants for alternative therapy for hyperlipidemia in Sukasari Region, Subang, West Java, Indonesia using a field survey method.

MATERIALS AND METHODS

Study Area

Sukasari is located in Subang Regency, West Java, Indonesia, with an area of 51.79 km². This area has an altitude of 3 meters above sea level with an average maximum air temperature of 33°C and a minimum of 23°C. Moreover, it is located between 06°34' 11" South Latitude and 107°45' 46" East Longitude. This area is a tropical climate area that is mostly inhabited by Sundanese tribes (98%) and other tribes (2%). Vegetation in the study area is in humid conditions with an average rainfall of 2000 mm/year.

Data Collection

An extensive field survey was carried out to obtain information about medicinal plants from the Sundanese tribe in the study area. To document existing information about medicinal plants from tribal practitioners, several field visits were conducted from March to April 2024 in the Sukasari Region, Subang, West Java, Indonesia. During the research, ethnomedicinal information was collected from middle-aged and older tribal practitioners in their local language (Sundanese), through direct interviews, questionnaires, and discussions. Information on local names of plants, plant parts used, preparation methods and administration routes (e.g., infusion, paste, juice and decoction) of all ethnomedicinal plants collected were recorded during the survey period.

Botanical Identification

Plant species are identified based on standard taxonomic methods, flower morphological characteristics, and

where possible, using samples for comparison, as well as consultation with experts and the literature.^[7] The plant types obtained were grouped into families according to the Cronquist classification system, except for Pteridophyta and Gymnospermae.^[8] Plant names were checked against the Plant List (www.plantlist.org) and the International Plant Name Index (www.ipni.org).

Ethics Statement

All participants provided verbal consent before the interview and gave consent to publish the information they provided.

RESULTS AND DISCUSSION

This research revealed that 30 plant species are commonly used by local people to treat hyperlipidemia (Table 1). This shows that the study location is affordable in terms of biodiversity. Among the various plant parts used, leaves (50.0%) are most frequently used in making medicines, followed by rhizomes (13.3%), fruit (13.3%), flowers, stem, and seed (respectively 6.7%), and rind (3.3%). The use of leaves is reported to be easier to prepare and easier to extract active substances from them for treatment. At the same time, leaves have less effect on the mother plant.^[9] Meanwhile, the most frequently used preparation methods were decoction (76.7%) and infusion (23.3%). These results are in line with previous research which reported that the forms of traditional medicine most widely used by the community were decoctions and infusions.^[7]

Table 1: Ethnomedicinal plants, local name, part used, mode of administration, and dosage uses in Sukasari, Subang, West Java, Indonesia.

No	Species	Family	Local name	Parts used	Mode of administration	Dosage of use
1	<i>Allium sativum</i> L.	Alliaceae	Bawang Putih	Rhizome	Infusion	15 grams once a day
2	<i>Aloe vera</i> Burm.f.,	Asphodelaceae	Lidah Buaya	Stem	Decoction	200 grams once a day
3	<i>Andrographis paniculata</i> Nees	Acanthaceae	Sambiloto	Leaf	Decoction	25 grams once a day
4	<i>Annona muricata</i> L.	Annonaceae	Sirsak	Leaf	Infusion	150 grams once a day
5	<i>Anredera cordifolia</i> (Ten.) Steenis	Basellaceae	Binahong	Leaf	Decoction	50 grams once a day
6	<i>Averrhoa carambola</i> L.	Oxalidaceae	Belimbing	Leaf	Infusion	40 grams once a day
7	<i>Carica papaya</i> L.	Caricaceae	Pepaya	Leaf	Decoction	300 grams once a day
8	<i>Cinnamomum verum</i> J.Presl	Lauraceae	Kayu Manis	Stem	Decoction	80 grams once a day
9	<i>Citrus aurantifolia</i> (Christm) Swingle	Rutaceae	Jeruk Nipis	Fruit	Decoction	50 grams once a day
10	<i>Clitoria ternatea</i> L.	Fabaceae	Bunga Telang	Flower	Decoction	8 grams once a day
11	<i>Cosmos caudatus</i> Kunth	Asteraceae	Kenikir	Leaf	Decoction	50 grams once a day
12	<i>Curcuma longa</i> L.	Zingiberaceae	Kunyit	Rhizome	Infusion	120 grams once a day
13	<i>Garcinia mangostana</i> L.	Clusiaceae	Manggis	Rind	Infusion	500 grams once a day
14	<i>Hibiscus sabdariffa</i> L.	Malvaceae	Rosela	Flower	Decoction	40 grams once a day

15	<i>Kaempferia galanga</i> L.	Zingiberaceae	Kencur	Rhizome	Infusion	300 grams once a day
16	<i>Mangifera indica</i> L.	Anacardiaceae	Mangga	Leaf	Decoction	150 grams once a day
17	<i>Momordica charantia</i> L.	Cucurbitaceae	Pare	Leaf	Decoction	90 grams once a day
18	<i>Morinda citrifolia</i> L.	Rubiaceae	Mengkudu	Fruit	Infusion	8 grams once a day
19	<i>Moringa oleifera</i> Lamk.	Moringaceae	Kelor	Leaf	Decoction	70 grams once a day
20	<i>Nigella sativa</i> L.	Ranunculaceae	Jinten Hitam	Seed	Decoction	150 grams once a day
21	<i>Pandanus amaryllifolius</i> Roxb.	Pandanaceae	Pandan Wangi	Leaf	Decoction	50 grams once a day
22	<i>Persea Americana</i> Mill.	Lauraceae	Alpukat	Seed	Decoction	100 grams once a day
23	<i>Phaleria macrocarpa</i> (Scheff.) Boerl)	Thymelaceae	Mahkota Dewa	Fruit	Decoction	500 grams once a day
24	<i>Phyllanthus niruri</i> L.	Phyllanthaceae	Meniran	Leaf	Decoction	30 grams once a day
25	<i>Piper betle</i> L.	Piperaceae	Sirih	Leaf	Decoction	60 grams once a day
26	<i>Psidium guajava</i> L.	Myrtaceae	Jambu Biji	Leaf	Decoction	60 grams once a day
27	<i>Solanum torvum</i> Sw.	Solanaceae	Tokakak	Fruit	Decoction	65 grams once a day
28	<i>Syzygium polyanthum</i> (Wight) Walpers	Myrtaceae	Salam	Leaf	Decoction	100 grams once a day
29	<i>Tinospora crispa</i> L.	Menispermaceae	Baratawali	Leaf	Decoction	100 grams once a day
30	<i>Zingiber officinale</i> Rosc.	Zingiberaceae	Jahe	Rhizome	Decoction	20 grams once a day

CONCLUSIONS

The results of this research confirm that people in the Sukasari Region still rely heavily on medicinal plants for their health care system, especially for the treatment of hyperlipidemia with the most frequently used parts of the leaves and their use in decoctions and infusions.

REFERENCES

1. Sahebkar A, Beccuti G, Simental-Mendía LE, Nobili V, Bo S. *Nigella sativa* (black seed) effects on plasma lipid concentrations in humans: A systematic review and meta-analysis of randomized placebo-controlled trials. *Pharmacol Res*, 2016; 106: 37-50.
2. Alkandahri MY, Kusumiyati K, Renggana H, Arfania M, Frianto D, Wahyuningsih ES, et al. Antihyperlipidemic activity of extract and fractions of *Castanopsis costata* leaves on rats fed with high cholesterol diet. *RASĀYAN J Chem*, 2022; 15(4): 2350-2358.
3. Ghule BV, Ghante MH, Saoji AN, Yeole PG. Antihyperlipidemic effect of the methanolic extract from *Lagenaria siceraria* Stand. fruit in hyperlipidemic rats. *J Ethnopharmacol*, 2009; 124(2): 333-337.
4. Bidkar JS, Ghanwat DD, Bhujbal MD, Dama GY. Anti-hyperlipidemic activity of Cucumis melo fruit peel extracts in high cholesterol diet induced hyperlipidemia in rats. *J Complement Integr Med*, 2012; 9(1): 1-18.
5. Carvalho AAS, Lima ÜWP, Valiente RA. Statin and fibrate associated myopathy: study of eight patients. *Arquivos de Neuro-Psiquiatria*, 2004; 62(2a): 257-261.
6. Politi FA, de Mello JC, Migliato KF, Nepomuceno AL, Moreira RR, Pietro RC. Antimicrobial, cytotoxic and antioxidant activities and determination of the total tannin content of bark extracts *Endopleura uchi*. *Int J Mol Sci*, 2011; 12(4): 2757-2768.
7. Bieski IGC, Santos FR, de Oliveira RM, Espinosa MM, Macedo M, Albuquerque UP, de Oliveira Martins DT. Ethnopharmacology of medicinal plants of the Pantanal Region (Mato Grosso, Brazil). *Evid Based Complement Alternat Med*, 2012; 2012: 1-36.
8. Cronquist A. The evolution classification of flowering plants. The New York Botanical Garden, New York, NY, USA, 2nd edition, 1988.
9. Ahmed S, Ahmad M, Swami BL, Ikram S. A review on plants extract mediated synthesis of silver nanoparticles for antimicrobial applications: Agreen expertise. *J Adv Res*, 2016; 7(1): 17-28.