



ADVANCED SYNTHESIS AND STUDY OF SUBSTITUTED 1,3-THIAZINE AND ITS NANOPARTICLES ON PHYTOTIC GROWTH OF SOME FLOWERING PLANTS

*Chhaya D. Badnakhe

Department of Chemistry, Dr.Manorama and Prof.H.S.Pundkar, Arts, Commerce and Science College, Balapur, Dist. Akola.



***Corresponding Author: Chhaya D. Badnakhe**

Department of Chemistry, Dr.Manorama and Prof.H.S.Pundkar, Arts, Commerce and Science College, Balapur, Dist. Akola.

DOI: <https://doi.org/10.5281/zenodo.21292517>



How to cite this Article: *Chhaya D. Badnakhe. (2026). Advanced Synthesis And Study Of Substituted 1,3-Thiazine And Its Nanoparticles On Phytotic Growth Of Some Flowering Plants. European Journal of Biomedical and Pharmaceutical Sciences, 13(7), 432-437.

This work is licensed under Creative Commons Attribution 4.0 International license.

Article Received on 15/06/2026

Article Revised on 05/07/2026

Article Published on 10/07/2026

ABSTRACT

The synthesis, spectral analysis and biological activities of 4-phenyl-2-hydroxy-chlorosubstituted-2-imino-1,3-thiazines have been carried out. In this case 4-(2'-hydroxy-3',5'-dichlorophenyl)-6-(hexyl)-2-iminophenyl-3,6-dihydro-1,3-thiazine (H) has been screened. The compound H was synthesized from 2'-hydroxy-3',5'-dichlorophenyl-4-(hexyl)chalcone (a'') by the action of phenylthiourea. The compound (a'') was synthesized from 2'-hydroxy-3',5'-dichloroacetophenone by the action of heptanaldehyde in ethanol and 40% NaOH. The nanoparticles of the compound H has been prepared by using ultrasonic technique. The titled compound and its nanoparticles were screened for their growth promoting activity on some flowering plants viz. *Crysanthemum coronarium*, *Dahlia pinnata*, *Verbena officinalis*, *Iberis amara*.

KEYWORDS: Chalcone, thiazine, phenyl thiourea, growth promoting activities.

INTRODUCTION

Thiazine is a six membered ring system, which contains two hetero atom [N and S] placed in the heterocyclic ring at 1, 3 positions. Many workers have synthesized different 1,3-thiazines.^[1-3] Thiazines are very useful units in the field of medicinal and pharmaceutical chemistry and have been reported to exhibit a variety of biological activities such as antiserotonin,^[4] antipsychotic,^[5] anti-HIV,^[6] antimicrobial,^[7] antiviral,^[8] blood platelet aggregation inhibitors^[9] and Ca⁺⁺ antagonist.^[10] Moreover thiazine nucleus is a pharmacophore of cephalosporin that occupy a very important place in the antibiotics and antifungal activity of thiazine nucleus is due to the presence of thiourea linkage in the structure. Chalcones and their analogues having α,β -unsaturated carbonyl system are very versatile substrates for the evolution of various reactions and physiologically active compounds. The reaction of thiourea with α,β -unsaturated ketones results in 1,3-thiazine. It has been well focused that the presence of 4-phenyl-chlorosubstituted moieties as well as 2-substituted amino group present in thiazine ring is an important structural feature and the resulting molecule would exhibit

promising biological activities.^[11-13] In the present study, various 4-phenyl-2-substituted-imino-thiazines have been synthesized from chalcones by using thiourea, phenyl thiourea and diphenyl thiourea. The synthesized compounds were evaluated for their growth promoting activity on some flowering plants viz. *Crysanthemum coronarium*, *Dahlia pinnata*, *Verbena officinalis*, *Iberis amara*.

EXPERIMENTAL

All the glasswares used in the present work were of pyrex quality. Melting points were determined in hot paraffin bath and are uncorrected. The purity of compounds was monitored on silica gel coated TLC plate. IR spectra were recorded on Perkin-Elmer spectrophotometer in KBr pellets, ¹H NMR spectra on spectrophotometer in CDCl₃ with TMS as internal standard. UV spectra were recorded in nujol medium. The analytical data of the titled compounds was highly satisfactory. All the chemicals used were of analytical grade. All the solvents used were purified by standard methods. Physical characterisation data of all the compounds is given in Table 1.

2'-Hydroxy 3',5'-dichloroacetophenone

2-Hydroxy-5-chloroacetophenone was dissolved in acetic acid (5 ml), Sodium acetate (3g) was added to the reaction mixture and then chlorine in acetic acid reagent (40 ml; 7.5 w/v) was added dropwise with stirring. The temperature of the reaction mixture was maintained below 20°C. The mixture was allowed to stand for 30 minutes. It was poured into water with stirring. A pale yellow solid then obtained was filtered, dried and crystallized from ethanol to get the compound 2'-hydroxy 3',5'-dichloroacetophenone.

Preparation of 2'-hydroxy-3',5'-dichlorophenyl-4-hexylchalcone (a'')

2'-Hydroxy-3',5'-dichloroacetophenone (0.1 mol) was dissolved in ethanol (50 ml) and heptanaldehyde (0.1 mol) was added gradually to the above solution and the mixture was heated to boiling temperature. Aqueous sodium hydroxide solution [40%; 40 ml] was added dropwise with constant stirring. The mixture was stirred mechanically at room temperature for about half an hour and kept overnight. It was then acidified by hydrochloric acid solution (10%). The solid product separated, was filtered, and washed with sodium bicarbonate (10%) followed by water. The crude product was crystallized from ethanol acetic acid mixture (a'').

Preparation of 4-(2'-hydroxy-3',5'-dichlorophenyl)-6-(hexyl)-2-iminophenyl-3,6-dihydro-1,3-thiazine (H)

2'-Hydroxy-3',5'-dichlorophenyl-4-hexylchalcone (a'')(0.01 mol) and phenyl thiourea (0.01 mol) were

dissolved in ethanol (25 ml). To this aqueous solution of KOH (0.02 mol) was added. The reaction mixture was refluxed for three hours cooled, diluted with water and acidified with 1:1 HCl. The product thus separated was filtered and crystallized from ethanol to get the compound (H).

The newly synthesized compound was characterised on the basis of elemental analysis, molecular determination, UV, IR, NMR. spectral data.

The UV, IR, and NMR spectral data**Compound (H)****UV: Spectrum No. 1**


The UV-Vis spectrum of the compound 5b reported in dioxane showed $\lambda_{\max}$ value 355 nm corresponding to $n \rightarrow \pi^*$ transition.

IR KBr: Spectrum No. 2

3421 cm^{-1} (O-H phenolic), 2857.60 cm^{-1} (aliphatic -C-H stretching), 2954.57 cm^{-1} (aromatic C-H stretching), 3786 cm^{-1} (-NH stretching), 1595.95 cm^{-1} (-C=N-stretching), 1030.61 cm^{-1} [(C-N) (C-NO₂) stretching], 748.64 cm^{-1} [C-Cl stretching in aliphatic], 1177.4 cm^{-1} [C-Cl stretching in aromatic].

PMR: Spectrum No. 3

δ 0.82 (t, 3H, -CH₂-CH₃); δ 1.01 (envelope of -CH₂, 10H, -(CH₂)₅-CH₃); δ 2.2 (s, 1H, =N-H); δ 3.1 (d, 1H, -C=C-H); δ 6.9 to 7.3 (m, 7H, Ar-H); δ 9.4 (s, 1H, O-H).

Scheme

Where

- 1) R₁ = -H
- 2) R₂ = -C₆H₅

Growth Promoting Effect on some Flowering Plants

The experimental set up of the study was divided into two parts:

- (i) Seed treatment (ii) Field experiment.

(i) Seed treatment

With a view to safeguard dormant seed's potential from harmful external agencies, the seeds of the test plants were treated by test compounds before sowing.

(ii) Field experiment

Pregerminated quality seeds of *Crysanthemum coronarium*, *Dahlia pinnata*, *Verbena officinalis*, *Iberis amara*. were procured from Department of Horticulture, Dr. PDKV, Akola.

The beds of cotton soil, 2.5 x 2.5 m size were prepared in an open field. The sowing of seeds of all four test flowering plants were done in separate beds and irrigated periodically.

The plants from each bed were divided into two groups i.e. A and B and designated as "Control" and "Treated" group plants respectively.

The plants from group B were sprayed with the solution of test compounds at weekly intervals. The field

experiments were conducted to compare the treated plants of group B with untreated plants of controlled group A. In this context, the observations were recorded on 7, 14, 21, 28, 35, 42, 45, 56, 63, 70, 77, 84, 91 days after sowing corresponding to early vegetative, late vegetative, flowering, pod filling and pod maturation, with special reference to number of leaves and height of shoots.

The results of field's experiments are tabulated in the tables 2, 3 and 4.

Table 1: Characterisation data of newly synthesized compounds.


Compounds	Molecular formula	M.P. in °C	% of yield	% of element			
				C	H	N	S
	C ₈ H ₆ O ₂ Cl ₂	54	80	47.90/48	2.95/3		
a''	C ₁₅ H ₁₈ O ₂ Cl ₂	103	70	59.80/59.25	53.10/60.26		
H	C ₂₂ H ₂₃ ON ₂ Cl ₂ S	160	50	61.141/51.50	5.32/5.40	6.48/6.52	7.40/7.48

Activity of the test compound H


Table No. 3: 4-(2'-Hydroxy-3',5'-dichlorophenyl)-6-(hexyl)-2-iminophenyl-3,6-dihydro-1,3-thiazine (H).

Periodicity of Observations [in days]	<i>Crysanthemum coronarium</i>				<i>Dahlia pinnata</i>				<i>Verbena officinalis</i>				<i>Iberis amara</i>			
	Shoot height		No. of leaves		Shoot height		No. of leaves		Shoot height		No. of leaves		Shoot height		No. of leaves	
	C	T	C	T	C	T	C	T	C	T	C	T	C	T	C	T
7	1.0	1.0	1	1	2.5	1.5	2	2	4.4	4	2	2	2	2	2	3
14	1.2	1.2	2	2	7.5	7	2	2	10	8	2	2	2.1	2.5	2	3
21	1.3	1.4	7	10	8	12	2	4	15	11	3	5	2.3	2.8	3	4
28	1.5	1.6	9	11	9	19	3	6	16	18	4	6	2.5	2.7	4	5
35	1.6	1.8	10	12	11	26	4	7	18	19	5	9	2.8	3.0	5	7
42	1.8	2.0	12	15	17	42	5	8	19	20	7	12	3.0	3.4	6	8
49	2.3	3.5	14	18	25	48	6	8	20	21	8	14	3.5	3.9	8	9
56	3.6	4.0	16	22	28	52	7	9	23	25	10	17	3.8	4.5	10	12
63	5.5	6.7	18	24	31	55	8	10	25	30	12	18	4.2	5.0	12	14
70	7	12	20	28	34	60	9	11	27	32	14	20	4.6	5.4	14	16
77	14	19	22	30	36	63	10	13	28	35	16	23	5.5	7.0	16	18
84	20	24	24	32	38	65	11	15	29	36	18	25	7.2	14	20	24
91	24	30	26	36	40	68	12	17	36	38	20	27	8.2	17	25	29

Impact of the 4-(2'-Hydroxy-3',5'-dichlorophenyl)-6-(hexyl)-2-
iminophenyl-3,6-dihydro-1,3-thiazine (H) on phytotic growth of *Crysanthemum coronarium*




Impact of the compound 4-(2'-Hydroxy-3',5'-dichlorophenyl)-6-(hexyl)-2-
iminophenyl-3,6-dihydro-1,3-thiazine (H) on phytotic growth of *Dahlia pinnata*




Impact of the compound 4-(2'-Hydroxy-3',5'-dichlorophenyl)-6-(hexyl)-2-
iminophenyl-3,6-dihydro-1,3-thiazine (H) on phytotic growth of *Verbena officinalis*




Impact of the 4-(2'-Hydroxy-3',5'-dichlorophenyl)-6-(hexyl)-2-
iminophenyl-3,6-dihydro-1,3-thiazine (H) on phytotic growth of *Iberis amara*






Spectrum No. 1.

RC SAIF PU, Chandigarh



Spectrum No. 2.



Spectrum No. 3.

RESULT AND DISCUSSION

The titled compound and its nanoparticles were screened for their growth promoting activity on test flowering plants viz, *Crysanthemum coronarium*, *Dahlia pinnata*, *Verbena officinalis*, *Iberis amara*.

When a comparison of morphological characters was made between those of treated and control group plants, it was interesting to note that all the treated plants exhibited significant shoot growth and considerable

increase in the number of leaves as compared to those of untreated ones.

ACKNOWLEDGEMENTS

The authors are thankful to the Principal, Dr.D.H.Pundkar, Dr.Manorama & Prof.H.S.Pundkar, Arts, Commerce & Science College, Balapur, Dr.B.B.Wankhade, Principal, Malkapur Vidnyan Mahavidyalaya, Malkapur for providing necessary facilities to carry out the research work.

REFERENCES

1. Kartrizky A.R., and Boulton J.A., "*Advances in heterocyclic chemistry*", vols 1 to 2 Academic press, New York, 1963-1980.
2. Gilchrist, T.L., "*Heterocyclic Chemistry*", Pitman, London, 1985.
3. Weissberger, [Ed.] A., "*The chemistry of Heterocyclic Compound*", vols. 1 to 29, Wiley Interscience, New York, 1950 to 1975.
4. H. Saldowska, A.B. Malik and T. Tawisza, *Farmaco Ed. Sci.*, 40: 58, 1985.
5. M.E. Wolff, "*Burger Medicinal Chemistry*", 116 [Wolff vol. IV part III] 889.
6. I.A. Shehata, H.I. Elsubbagh, A.M. Abdelal, M.A. Sherbeny and A.A. Aaobaid, *Med. Chem. Res.*, 1996; 148.
7. A.K. Bhatt, H.G. Karadia and P.R. Shab., *Indian J. Heterocycl. Chem.*, 2004; 13: 281.
8. V.K. Pandey, S.K. Saxena and S.K. Bajpai, *Indian J. Chem.*, 43 B, 2004; 1015.
9. C. Brown and R.N. Davidson, *Adv. Heterochcl. Chem.*, 1985; 38: 135.
10. a) A.A. Fodda, A.M. Khalil and M.M. El. Habbal, *J. Indian Chem. Soc.*, 68, 1991, 393. b) R.L. Shriner and H.R. Todd., *Org. Synth.*, 1943; 11: 200.
11. M.S.K. Youself, *Indian J. Chem.*, 19 B, 1980; 796.
12. G. Bernath, Z. Szakoni, F. Fulop and P. Sohar, *Acta Pharma, Hung.*, 1994; 64: 153.
13. W. Peter and B.H. Hgregory, *Tetrahedron*, 1998; 54: 6987.