EUROPEAN JOURNAL OF BIOMEDICAL AND PHARMACEUTICAL SCIENCES
http://www.ejbps.com

ISSN 2349-8870
Coden USA: EJBPBO
Volume: 13
Issue: 11
Year: 2026

SPECTROSCOPIC AND COMPUTATIONAL ANALYSIS OF THREONINE: A REVIEW

Dr. Santosh Kumar Srivastava*

Department of Physics, Singhania University, Pacheri Bari, Jhunjhunu (Rajasthan)-333515, India.

***Corresponding Author: Dr. Santosh Kumar Srivastava**

Department of Physics, Singhania University, Pacheri Bari, Jhunjhunu (Rajasthan)-333515, India.

DOI: https://doi.org/10.5281/zenodo.23805060

How to cite this Article: Dr. Santosh Kumar Srivastava*. (2026). Spectroscopic and Computational Analysis of Threonine: A Review. European Journal of Biomedical and Pharmaceutical Sciences, 13(11), 75–82.

This work is licensed under Creative Commons Attribution 4.0 International license.

Article Received on 04/10/2026 **Article Revised on 24/10/2026** **Article Published on 01/11/2026**

ABSTRACT

Threonine is an essential proteinogenic amino acid whose molecular structure combines amino, carboxyl, hydroxyl, and methyl groups. This combination makes it an informative model for studying the relationship between molecular conformation, hydrogen bonding, protonation state, and vibrational spectra. This review surveys experimental and computational approaches used to investigate L-threonine in crystalline, isolated-molecule, and aqueous environments. The spectroscopic methods discussed include infrared absorption, Fourier-transform infrared (FTIR), Raman, far-infrared, vibrational circular dichroism (VCD), rotational microwave spectroscopy, and inelastic incoherent neutron scattering. Computational approaches include molecular mechanics conformational searches, density functional theory (DFT), post-Hartree-Fock calculations, implicit and explicit solvation, normal-mode analysis, and periodic calculations for crystals. Published work has demonstrated that threonine cannot be represented adequately by a single geometry in every environment: multiple conformers may contribute, and cationic, zwitterionic, and anionic forms differ in their bonding and spectra. The low-frequency region can be particularly informative about torsional and skeletal motions, while VCD adds sensitivity to molecular chirality. In the solid state, intermolecular hydrogen bonding and crystal packing affect vibrational and electronic properties. The review also considers limitations in frequency prediction, solvent modelling, conformer populations, and comparisons between isolated molecules and crystals. Finally, it identifies opportunities for reproducible workflows that combine measured spectra, explicit solvent modelling, conformational sampling, and uncertainty-aware assignments.

KEYWORDS: L-Threonine, FTIR, Raman Spectroscopy, Far-Infrared Spectroscopy, Vibrational Circular Dichroism, Density Functional Theory, Conformational Analysis, Hydrogen Bonding, Molecular Modelling.

INTRODUCTION

Amino acids are central to chemistry and biology because they form the building blocks of peptides and proteins. Their structures contain an amino group, a carboxyl group, and a side chain whose composition influences polarity, conformational flexibility, and intermolecular interactions. Threonine is an essential amino acid, meaning that humans must obtain it through dietary sources. Its side chain contains a hydroxyl group and a methyl group, making it a useful system for examining the combined effects of polarity and steric constraints.[1] L-threonine is conventionally identified as (2S, 3R)-2-amino-3-hydroxybutanoic acid. It contains two stereogenic centres and must be distinguished from L-allo-threonine, which has a different configuration at one centre. This stereochemical distinction matters when interpreting chiroptical measurements and when constructing computational starting geometries.[1, 2] Spectroscopic methods report different aspects of a molecule. Infrared spectroscopy responds to changes in dipole moment during vibration, while Raman spectroscopy responds to changes in polarizability. VCD measures differences in infrared absorption of left- and right-circularly polarized light and is sensitive to chirality. Rotational spectroscopy can identify gas-phase conformers with high structural specificity. Computational chemistry helps connect these measurements to geometries, normal modes, relative energies, and interactions. A useful review must distinguish solid-state, solution-phase, and isolated-molecule results. Crystals contain extended hydrogen-bond networks; aqueous solutions involve hydration and

protonation equilibria; gas-phase experiments may reveal conformers that are not dominant in condensed phases. The same molecule may therefore show different spectra under different conditions.

CHEMICAL IDENTITY AND MOLECULAR STRUCTURE




Figure 1: Skeletal formula of threonine.[8]

The molecular formula of threonine is $C_4H_9NO_3$, and its molar mass is approximately 119.12 g mol^{-1}. A condensed representation of the neutral connectivity is CH_3-CH (OH)-CH (NH_2)-COOH. Near neutral pH in water, the predominant form is generally zwitterionic, with a protonated amino group and a deprotonated carboxyl group: CH_3-CH (OH)-CH (NH_3^+)-COO^-. The exact distribution among protonation states depends on pH and the relevant acid-base equilibria. The amino group contributes N-H stretching and deformation modes. The carboxyl group in its protonated form has a carbonyl vibration, while the carboxylate form has coupled asymmetric and symmetric COO^- stretching modes. The side-chain hydroxyl group contributes O-H stretching and bending vibrations and can participate in hydrogen bonding. The methyl group produces C-H stretching, bending, and rocking modes. These motions are not completely independent: normal modes can involve coordinated displacement of several atoms. The molecule has rotatable bonds around the backbone and side chain. Rotation changes the relative orientation of the amino, carboxyl, and hydroxyl groups, affecting intramolecular hydrogen bonding and the molecule's dipole moment. The two stereogenic centres constrain the accessible geometries and distinguish threonine from its stereoisomers. In a crystalline sample, the molecular geometry is influenced by packing and intermolecular hydrogen bonds. In water, hydration shells and solvent fluctuations alter the relative stability of conformations. For these reasons, chemical identity, stereochemistry, protonation state, physical phase, temperature, and sample preparation should always be reported when presenting a spectrum.

PROTONATION STATES AND ACID-BASE BEHAVIOUR

Threonine can exist as cationic, zwitterionic, or anionic species depending on solution pH. Under strongly acidic conditions, the amino group is protonated and the carboxyl group may remain protonated, giving a net positive charge. Near neutral pH, the carboxyl group is generally deprotonated while the amino group remains protonated, producing the zwitterion. Under strongly basic conditions, the ammonium group can lose a proton, yielding a net negatively charged species. These forms differ in charge distribution and in their vibrational signatures. Protonation of the carboxylate changes the balance between C=O and C-O bonding. Conversion of NH_3^+ to NH_2 changes the number and character of N-H vibrations. Electrostatic interactions also affect conformation and hydration. Thus, a spectrum measured at one pH should not be compared uncritically with a calculation representing a different ionic form. A notable experimental and computational study examined L-threonine solutions at pH 1.00, 5.70, and 13.00, representing cationic, zwitterionic, and anionic conditions, respectively. The authors recorded mid-IR, far-IR, Raman, and VCD spectra and compared them with conformational searches and quantum-chemical calculations. Their conformational search identified numerous candidate structures before higher-level calculations were applied to selected conformers.[5] For reproducible work, pH should be measured rather than inferred from a recipe, and solution concentration, temperature, counterions, and solvent composition should be recorded. Strong acid or base can also affect ionic strength and may introduce spectral contributions from other species. Theoretical comparisons should use a consistent protonation state and should consider whether the model includes solvent stabilization.

FUNDAMENTALS OF MOLECULAR VIBRATIONS

Molecular vibrations are motions of atoms about an equilibrium structure. In the harmonic approximation, each normal mode is treated as an independent oscillator. The approximate vibrational frequency depends on the effective mass of the moving atoms and the force constant associated with the mode. A stronger bond generally vibrates at a higher frequency than a weaker bond involving similar masses, although mode coupling and molecular environment complicate simple rules. For a molecule with N atoms, a nonlinear isolated molecule has 3N-6 vibrational normal modes. Threonine has 13 atoms in its neutral formula representation, so the formal count is 33 vibrational modes for a nonlinear isolated structure. The practical number of observed bands may differ because modes overlap, some are weak, symmetry and selection rules affect activity, and condensed-phase spectra can broaden or split features. An infrared-active mode requires a change in dipole moment during vibration. A Raman-active mode requires a change in polarizability. A mode can be active in both methods, in one, or weak in both. The two techniques are therefore complementary rather than interchangeable. Real molecular vibrations are anharmonic. O-H and N-H stretching modes, in particular, may be affected by anharmonicity and hydrogen bonding. Harmonic DFT calculations often overestimate frequencies and are commonly compared with experiment after applying a scaling factor or a more sophisticated treatment. Scaling factors depend on the computational method and basis set; a universal factor should not be assumed. In amino

acids, internal coordinates such as bond stretches and angle bends mix into collective normal modes. Assignments should therefore be based on displacement vectors and calculated intensities, not solely on a frequency range. Isotopic substitution, polarization measurements, and comparison across phases can help test assignments.

FTIR AND INFRARED SPECTROSCOPY

FTIR spectroscopy records the absorption of infrared radiation and is widely used to identify functional groups and characterize molecular interactions. An interferometer collects an interferogram, which is transformed mathematically to produce a spectrum. FTIR instruments can provide rapid acquisition, good wavenumber reproducibility, and access to a broad spectral region. For threonine, important regions include X-H stretching, C-H stretching, carboxyl/carboxylate vibrations, and fingerprint-region modes involving C-N, C-O, and skeletal motions. The following are broad guide ranges rather than definitive assignments for a particular sample: O-H/N-H stretching may occur across roughly 3000-3700 cm^{-1}, C-H stretching often appears around 2800-3100 cm^{-1}; carboxylate asymmetric stretching may occur in the approximate 1550-1650 cm^{-1} region; and symmetric carboxylate stretching commonly occurs in a lower region around 1300-1450 cm^{-1}. Exact values depend on sample form, hydrogen bonding, and experimental conditions. A solid sample may be measured using an appropriate transmission pellet or an attenuated total reflectance (ATR) accessory. Aqueous measurements require attention to strong water absorption and path length. Background collection, atmospheric water-vapour correction, spectral resolution, number of scans, and sample handling should be reported. Broad bands are not necessarily evidence of a single vibration: hydrogen-bond distributions and overlapping modes can contribute. Assignments should be cross-checked against Raman data and quantum-chemical calculations. When comparing samples at different pH, changes in the carboxylate and amino regions may reflect altered protonation rather than a conformational change alone.

RAMAN SPECTROSCOPY

Raman spectroscopy probes inelastic scattering of monochromatic light. A small fraction of incident photons exchange energy with molecular vibrations, producing Stokes and anti-Stokes shifts. The intensity of a Raman band depends on the change in polarizability along the normal coordinate. Raman measurements can complement IR spectroscopy, particularly for modes whose polarizability changes are substantial. Published work has characterized L-threonine in solid form and in aqueous solution using Raman methods, and computational models have been used to interpret monomeric and dimeric structures. Guzzetti and colleagues studied zwitterionic L-threonine in aqueous media, comparing experimental Raman results with B3LYP calculations and implicit-solvent models. They also examined hydrogen bonding using atoms-in-molecules and natural-bond-orbital analyses.[4] Raman experiments can be influenced by fluorescence, laser heating, sample degradation, and the optical properties of the sample holder. For crystals, polarization and crystal orientation may change relative band intensities. For solutions, concentration and solvent background matter. Laser power should be kept low enough to avoid heating or decomposition, and replicate measurements are valuable. A Raman spectrum should not be interpreted by matching each peak to a generic table alone. The strongest assignments are supported by calculated Raman activities, normal-mode visualizations, isotopic evidence where available, and comparison with known spectra. It is also important to state whether the reported spectrum is normalized, baseline-corrected, or otherwise processed. When IR and Raman spectra are analyzed together, the combined evidence can help distinguish overlapping vibrations and clarify how a conformational change affects dipole moment and polarizability differently.

FAR-INFRARED AND LOW-FREQUENCY MODES

The far-infrared region is useful for examining low-frequency motions that may be difficult to characterize using conventional mid-IR measurements. These modes can include torsions, skeletal deformations, coupled intermolecular motions, and lattice vibrations in crystals. Because the energy differences between conformers can be small, low-frequency spectral features may provide clues about molecular flexibility and intermolecular organization. The study of L-threonine under acidic, neutral, and basic conditions reported that the low-frequency region, approximately 30-700 cm^{-1}, was particularly useful for examining the zwitterionic conformational landscape.[5] This result illustrates that conformational analysis should not be limited to the most familiar functional-group region. Low-frequency spectra require careful instrumental setup and background treatment. Atmospheric absorption, window materials, detector response, and sample thickness can affect measurements. In a crystalline solid, modes can be collective and depend on the unit cell, so an isolated-molecule calculation cannot fully reproduce all lattice vibrations. Computational interpretation should distinguish intramolecular normal modes from lattice modes. Periodic DFT can be appropriate for crystalline threonine, while molecular calculations are more suitable for isolated conformers or small hydrated clusters. Comparing these models directly without accounting for their different physical assumptions can lead to misleading assignments. Far-IR measurements are particularly valuable when combined with mid-IR, Raman, and calculated normal-mode displacement patterns. The combined data can constrain possible conformations more strongly than a single spectral region.

VIBRATIONAL CIRCULAR DICHROSIM

Vibrational circular dichroism is a chiroptical technique that measures the differential absorption of left- and right-circularly polarized infrared light. The sign and magnitude of VCD features depend on molecular chirality, conformation, and the coupling of vibrational transition moments. VCD is especially useful for molecules with stereogenic centres and for distinguishing candidate conformations that may have similar conventional IR spectra. L-threonine has two stereogenic centres. Its stereochemical identity must be specified accurately, because the calculated VCD spectrum depends on configuration. The stereoisomer L-allo-threonine is not interchangeable with L-threonine. A comparison between experiment and theory should use the correct configuration, protonation state, solvent model, and conformer ensemble. A published study measured VCD spectra for L-threonine solutions at three pH values and compared them with quantum-chemical calculations.[5] The work demonstrates how VCD can be combined with conventional vibrational spectra to investigate the conformational landscape across protonation states. VCD interpretation is challenging because individual bands can be weak and overlapping, and the spectrum may represent a weighted contribution from multiple conformers. A single optimized geometry may not adequately reproduce the experimental signal. Conformer populations are often estimated using relative free energies, but those estimates depend on the quality of the conformational search, solvation model, and treatment of entropy. Reliable reporting should include the experimental conditions, baseline treatment, calculated method, frequency scaling, and the method used to combine conformer spectra. VCD is most persuasive when its interpretation is consistent with independent structural evidence.

ROTATIONAL AND NEUTRON SPECTROSCOPY

Rotational spectroscopy can provide exceptionally precise structural information for molecules in the gas phase. Laser-ablation molecular-beam Fourier-transform microwave spectroscopy has been used to characterize multiple conformers of L-threonine. Alonso and colleagues reported the identification of seven gas-phase conformers by comparing measured rotational and nuclear quadrupole coupling constants with theoretical predictions.[6] These measurements are significant because the polar side chain creates a network of possible intramolecular hydrogen-bond arrangements. Rotational constants are sensitive to the distribution of mass in the molecule, allowing candidate geometries to be tested. However, gas-phase conformer populations need not match those in aqueous solution or a crystal, where solvent interactions and packing change the energy landscape. Inelastic incoherent neutron scattering (IINS) is sensitive to vibrational motions, particularly those involving hydrogen atoms. Deuteration can help distinguish modes and support assignments. A combined investigation of neutron scattering, infrared, Raman, and ab initio calculations for L-threonine was published in 2001.[7] Such multimodal approaches are valuable because no single technique provides complete information about all molecular motions. Rotational spectroscopy, IINS, IR, and Raman methods operate through different physical interactions and are therefore complementary. Rotational data constrain geometry, neutron scattering helps characterize hydrogen-involving vibrations, and optical spectroscopy provides broader vibrational fingerprints. Computational calculations offer a common framework for interpreting these diverse observations.

CONFORMATIONAL LANDSCAPE

A conformer is a structure accessible through rotation about single bonds without changing the molecule's connectivity or stereochemical configuration. Threonine's backbone and hydroxyl-bearing side chain allow multiple conformers, with stability influenced by intramolecular hydrogen bonding, steric interactions, electrostatics, and solvent stabilization. A conformational search should generate a broad set of candidate structures before detailed electronic-structure calculations are performed. Molecular mechanics methods can be used for initial screening, followed by geometry optimization using DFT or another quantum-chemical method. Duplicate structures should be removed using geometric criteria, and optimized structures should be checked to ensure that they are genuine minima rather than transition states. The 2013 pH-dependent study began with conformational searches for the cationic, zwitterionic, and anionic forms. It reported 9 candidate zwitterions, 27 cations, and 52 anions before quantum-chemical optimization; the number of non-redundant structures was lower after optimization.[5] These values are results of that particular search protocol, not universal counts of all possible conformers. The lowest electronic-energy structure is not always the dominant conformer at a finite temperature. Population estimates depend on Gibbs free energies, solvent conditions, and entropy. A Boltzmann distribution is often used as an approximation, but the results are only as reliable as the calculated free energies and the completeness of the conformational search. For spectral simulation, the spectrum of each conformer can be weighted by its estimated population. This is preferable to selecting one structure without justification, especially when multiple conformers have comparable free energies.

DENSITY FUNCTIONAL THEORY

Density functional theory is widely used to predict molecular geometries, vibrational frequencies, electronic properties, and relative energies. In DFT, the electronic energy is expressed in terms of the electron density, and the accuracy depends on the exchange-correlation functional, basis set or numerical representation, and treatment of environmental effects. For molecular threonine, hybrid functionals such as B3LYP and meta-hybrid functionals such as M06-2X have been used with basis sets including 6-311++G(d,p). A pH-dependent conformational study applied B3LYP to candidate

conformers and used M06-2X and MP2 calculations for selected stable structures.[5] Different methods can predict slightly different relative energies, so cross-checking selected structures is useful.

Geometry optimization produces bond lengths, angles, and torsions. Frequency calculations provide harmonic normal modes and intensities. A minimum should normally have no imaginary vibrational frequencies; imaginary frequencies may indicate a transition state or an incomplete optimization. Calculated spectra should be scaled or otherwise corrected before direct frequency comparison with experiment. DFT limitations include functional dependence, incomplete treatment of dispersion in some models, harmonic approximation errors, and sensitivity to conformer choice. A method that performs well for one property may not be optimal for another. For example, an approach suited to molecular vibrations may not give the best crystal lattice parameters or electronic band gap. A reproducible computational report should identify the program, version, functional, basis set, integration settings, convergence thresholds, solvent model, frequency scaling procedure, and all assumptions used to select conformers.

SOLVATION AND HYDROGEN BONDING

Water affects threonine through electrostatic stabilization, hydrogen bonding, and solvent fluctuations. The polar amino and carboxylate groups are strongly hydrated, while the side-chain hydroxyl group can both donate and accept hydrogen bonds. Hydration changes molecular conformational preferences and can shift vibrational frequencies. Implicit solvent models, such as the integral equation formalism version of the polarizable continuum model (IEF-PCM), represent the solvent as a dielectric environment. They are computationally economical and can capture some bulk polarization effects. However, they do not explicitly represent individual water molecules or discrete hydrogen-bond arrangements. Explicit hydration models add water molecules around threonine. These models can describe specific donor-acceptor contacts but require sampling of many possible arrangements. A hybrid approach may combine explicit water molecules near the solute with an implicit continuum for the surrounding solvent. Guzzetti and colleagues investigated a threonine monomer, dimers, and hydrated complexes involving one to four discrete water molecules, with implicit-solvent effects included.[4] Their work illustrates how both molecular association and hydration can influence structural and vibrational properties. Hydrogen-bond analysis can use geometric criteria, electron-density topology, natural bond orbital analysis, or energy decomposition. No single descriptor fully captures hydrogen-bond strength in every context. Claims about hydrogen bonding should specify the analysis method and avoid treating a short contact as definitive proof of a strong interaction. A robust solution-phase model should consider protonation state, solvent representation, and conformational sampling together rather than treating them as independent choices.

DIMERIZATION AND INTERMOLECULAR ASSOCIATION

Threonine molecules can interact through intermolecular hydrogen bonds, forming dimers and larger associations under suitable conditions. The carboxylate, ammonium, and hydroxyl groups provide multiple possible interaction sites. Association can alter vibrational frequencies and intensities because hydrogen bonding changes bond strengths and couples motions between molecules. Theoretical dimer calculations require careful comparison of total energies. If two monomers are combined into a dimer, the binding energy is commonly estimated from the dimer energy minus the energies of the isolated monomers, with consistent computational settings. Basis-set superposition error may need to be considered for localized basis sets. Solvent effects can change the relative stability of dimers and monomers, so gas-phase binding energies should not be assumed to represent aqueous association. Experimental spectra may contain overlapping contributions from monomers, dimers, and solvent. However, observing a spectral feature alone does not uniquely establish a particular dimer geometry. A convincing assignment should compare plausible structures, predicted spectra, and the experimental conditions. Published Raman and computational work on aqueous zwitterionic L-threonine investigated monomeric and two dimeric structures, as well as hydrated complexes.[4] This research is relevant to understanding how local hydrogen-bond networks can influence the vibrational fingerprints of amino acids. Future work could combine concentration-dependent Raman or IR measurements with molecular dynamics and enhanced sampling. Such an approach may help distinguish persistent association from transient solvent-mediated contacts.

SOLID-STATE AND CRYSTAL CALCULATIONS

The solid state differs from an isolated molecule because threonine molecules are organized in a crystal lattice and interact through an extended network of hydrogen bonds. Crystal packing can affect bond lengths, molecular conformation, lattice vibrations, optical response, and thermal properties. A molecular calculation that omits neighbouring molecules cannot fully capture these collective effects. Araujo and colleagues investigated anhydrous orthorhombic L-threonine crystals using first-principles density functional theory with local-density and dispersion-corrected generalized-gradient approaches.[3] Their study considered structural, electronic, optical, vibrational, and thermodynamic properties. The reported calculations compared theoretical lattice parameters with X-ray data and examined vibrational and optical behaviour. Periodic DFT calculations treat the crystal as a repeating unit cell. They can predict optimized lattice parameters, electronic band structures, dielectric response, and phonon modes. Dispersion corrections may be important when

describing intermolecular packing, although the best method depends on the material and property. Experimental comparison should use the same phase and conditions where possible. Temperature, pressure, hydration, polymorphism, defects, and crystal orientation can affect measured spectra. A theoretical unit cell should be matched to the correct crystallographic structure and stoichiometry. Solid-state and solution-phase calculations answer different questions. A periodic calculation is appropriate for lattice and band properties, while molecular DFT is often more convenient for conformational analysis and solution-phase vibrations. The results should be compared only after accounting for these differences in physical model.

ELECTRONIC STRUCTURE AND OPTICAL PROPERTIES

Computational electronic-structure analysis can provide information about molecular orbitals, electron density, electrostatic potential, polarizability, and optical transitions. These descriptors can support structural interpretation, but their meaning depends on the computational method and the property being calculated. The highest occupied molecular orbital (HOMO) and lowest unoccupied molecular orbital (LUMO) are frequently visualized to inspect regions associated with occupied and unoccupied orbital density. The orbital-energy difference is often written as $\Delta E = E(LUMO) - E(HOMO)$. This difference is a computational descriptor and should not automatically be equated with an experimental optical absorption energy. Excited-state calculations or periodic electronic-structure methods may be needed for more direct comparisons. Molecular electrostatic potential maps show the electrostatic potential around a molecular electron density. They can help identify regions likely to interact with charged or polar species. In threonine, the carboxylate, ammonium, and hydroxyl regions are chemically important, but the map depends on the protonation state and geometry. For crystalline L-threonine, periodic calculations have been used to estimate electronic band gaps and optical properties.[3] The published work reported a calculated direct gap close to an experimental value cited by the authors. Such results apply to the crystal model and should not be transferred directly to isolated threonine molecules. Optical calculations should document the functional, basis or plane-wave cutoff, pseudopotentials, k-point sampling, polarization direction, and any correction applied to the band gap. Comparisons with experiment must also consider sample quality and measurement geometry.

DATA ANALYSIS AND SPECTRAL ASSIGNMENT

Spectral interpretation benefits from a structured workflow. First, sample identity and conditions should be documented. Second, instrumental background and baseline effects should be evaluated. Third, peaks or bands should be identified using reproducible criteria rather than manual selection alone. Finally, assignments should be tested against theoretical calculations and complementary experiments. For FTIR and Raman spectra, peak positions, widths, intensities, and relative shifts can all contain useful information. Broadening can arise from unresolved modes, hydrogen-bond distributions, thermal motion, and instrumental resolution. A single fitted peak may not represent a single normal mode, particularly in complex regions. Calculated normal modes should be inspected through their displacement vectors. A label such as “C-O stretching” is often an approximation because real modes can mix C-O stretching with bending, skeletal motion, and hydrogen displacement. The assignment is stronger when the predicted intensity is consistent with the experimental feature and when nearby modes are considered. Frequency scaling should be applied consistently. Scaling one region differently from another without a stated rationale can create misleading agreement. Where possible, comparisons should include both raw and scaled frequencies and explain the method used to select the scale factor. For a rigorous article, spectral tables should distinguish observed values from calculated values. Each row should identify the method, sample phase, protonation state, and assignment confidence. Approximate guide ranges should never be presented as exact experimental peaks. Where a published value is not available, the table should say “not reported” rather than filling the gap with an estimate.

LIMITATIONS AND SOURCES OF UNCERTAINTY

Several sources of uncertainty affect spectroscopic and computational studies of threonine. Experimentally, uncertainties may arise from calibration, sample purity, water content, temperature, pH, concentration, crystal orientation, baseline correction, and instrumental resolution. Repeated measurements and explicit reporting of conditions improve reproducibility. Computational uncertainties include incomplete conformational sampling, method and basis-set dependence, the harmonic approximation, imperfect solvation models, and errors in calculated relative free energies. The lowest-energy conformer predicted by one method may not remain lowest under another method or in a different solvent model. Small energy differences should therefore be interpreted cautiously. A further issue is the distinction between electronic energy and free energy. Conformer populations at finite temperature depend on entropy and environmental effects, not solely on optimized electronic energies. A population-weighted spectrum is only as reliable as the estimated populations. Comparing gas-phase calculations with solution measurements can be informative, but the differences should be discussed explicitly. Similarly, an isolated-molecule spectrum cannot reproduce all lattice modes of a crystal. Periodic and molecular calculations should be selected according to the research question. Finally, literature reviews can be limited by incomplete access to full text, inconsistent terminology, and differences in reported experimental conditions.

PROPOSED INTEGRATED RESEARCH WORKFLOW

A future experimental-computational study of L-threonine could use a staged workflow. The first stage would define the research question: for example, whether pH changes the conformational population, whether hydration shifts a particular band, or how the crystal spectrum differs from that of aqueous threonine. A clearly defined question prevents the study from collecting data without a focused interpretation. The second stage would document sample source, purity, stereochemical identity, concentration, solvent, pH, temperature, and physical phase. FTIR and Raman spectra could then be recorded under controlled conditions, with repeated measurements and appropriate background controls. If chirality or conformational populations are central, VCD could provide additional constraints. Far-IR or rotational spectroscopy may be considered where suitable instrumentation is available. The computational stage would begin with a broad conformational search for the correct protonation state. Candidate structures would be optimized, checked by frequency calculations, and ranked using consistent energy and solvation methods. Selected structures could be re-evaluated with a second functional or a higher-level method. Explicit hydration clusters or molecular dynamics would be considered if discrete solvent interactions are essential. The analysis stage would compare calculated and measured frequencies, intensities, and spectral patterns. Conformer contributions should be combined using justified populations rather than arbitrary averaging. Uncertainty and alternative assignments should be reported. Finally, all input structures, computational settings, scripts, raw spectra, processed spectra, and assignment tables should be archived. This makes the study easier to reproduce and allows later researchers to test alternative methods.

RESEARCH GAPS AND FUTURE DIRECTIONS

The existing literature provides a substantial foundation for understanding threonine, but several avenues remain valuable. One is improved comparison of phase-specific spectra using consistent computational protocols. Solid-state, solution, and gas-phase spectra reflect different physical environments and should be treated as complementary evidence rather than interchangeable measurements. Another direction is more systematic conformational sampling. Conventional searches may miss structures separated by high barriers or stabilized only by particular solvent arrangements. Enhanced-sampling molecular dynamics, automated conformer generation, and carefully validated free-energy methods may improve estimates of conformer populations. A third opportunity is to integrate explicit water molecules with continuum solvent models and compare the resulting spectra with solution experiments. This may be particularly important for the side-chain hydroxyl group and the charged amino and carboxylate groups. Isotopic substitution, including deuteration, can further assist in identifying hydrogen-involving modes. Machine-learning approaches may help accelerate spectral prediction or identify patterns in large datasets, but training data quality and chemical coverage are critical. Predictions should be validated against measurements and quantum-chemical calculations for the specific protonation state and environment. Reproducibility is also an important future priority. Authors should share optimized geometries, conformer energies, frequency lists, scaling factors, raw spectra where possible, and detailed metadata. A carefully documented data set would support fair comparisons among functionals, basis sets, and solvent models. These opportunities should be described as future research directions unless a targeted literature search establishes the current extent of work in each area.

CONCLUSION

Threonine is a useful model for examining how molecular structure, stereochemistry, protonation, hydration, and intermolecular interactions influence spectroscopic behaviour. Its amino, carboxyl, hydroxyl, and methyl groups contribute characteristic vibrational modes, while conformational flexibility and two stereogenic centres make structural interpretation more complex than a simple functional-group assignment. FTIR and Raman spectroscopy provide complementary information about vibrational transitions. Far-IR measurements can reveal low-frequency motions, VCD can constrain chiral conformations, rotational spectroscopy can identify gas-phase structures, and neutron scattering can support the analysis of hydrogen-involving vibrations. Together, these techniques provide a richer structural picture than any one method alone. Computational methods connect spectra to molecular structures. DFT and related approaches can optimize candidate conformers, predict vibrational frequencies and intensities, and examine hydrogen bonding and electronic properties. Solvent models are needed for solution studies, while periodic DFT is more appropriate for crystal packing, lattice vibrations, and band properties. The chosen model must match the physical question. Published research demonstrates that protonation state, solvent environment, conformational sampling, and hydrogen-bond networks all influence the observed behaviour of L-threonine. A robust interpretation therefore requires experimental conditions to be reported carefully and calculated spectra to be compared with the correct species and phase. The most reliable future studies will combine complementary spectroscopy, validated computational methods, transparent uncertainty reporting, and accessible research data. This integrated approach can improve vibrational assignments and contribute to broader understanding of amino-acid structure and interactions.

REFERENCES

1. Forbes KJ, Merheb E. Protein Structure and Function. StatPearls Publishing, 2026; Available from: https://www.ncbi.nlm.nih.gov/books/NBK562260/

2. Hernandez B, Pfluger F, Adenier A, Nsangou M, Kruglik SG, Ghomi M. Energy maps, side chain conformational flexibility, and vibrational features of polar amino acids L-serine and L-threonine in aqueous environment. J Chem Phys., 2011; 135: 055101.
3. Araujo RL, Vasconcelos MS, Barboza CA, Lima Neto JX, Albuquerque EL, Fulco UL. DFT calculations of the structural, electronic, optical and vibrational properties of anhydrous orthorhombic L-threonine crystals. Comput Theor Chem., 2019; 1170: 112621.
4. Guzzetti KA, Brizuela AB, Romano E, Brandan SA. Structural and vibrational study on zwitterions of L-threonine in aqueous phase using the FT-Raman and SCRF calculations. J Mol Struct, 2013; 1045: 171-179.
5. Moreno MMQ, Garcia AAM, Moreno JRA, Gonzalez JJL. Conformational landscape of L-threonine in neutral, acid and basic solutions from vibrational circular dichroism spectroscopy and quantum chemical calculations. Tetrahedron: Asymmetry, 2013; 24(24): 1537-1547.
6. Alonso JL, Perez C, Sanz ME, Lopez JC, Blanco S. (2009). Seven conformers of L-threonine in the gas phase: A LA-MB-FTMW study. Phys Chem Chem Phys, 2009; 11: 617-627.
7. Pawlukojc A, Leciejewicz J, Tomkinson J, Parker SF. Neutron scattering, infra red, Raman spectroscopy and ab initio study of L-threonine. Spectrochim Acta A Mol Biomol Spectrosc, 2001; 57(12): 2513-2523.
8. https://en.wikipedia.org/wiki/Threonine