



**SYNTHESIS, CHARACTERIZATION AND STUDY OF MICROBIOLOGICAL
ACTIVITY OF COBALT ION COMPLEX WITH [2 - (2 - METHOXYBENZYLIDINE)
AMINO PHENOL] (NOMBAP)**

Vasant Dnyandeo Barhate* and Ritika Manoj Makhijani

V.E. S. College of Arts, Science and Commerce, Sindhi Society Chembur, Mumbai - 400071, India.

***Author for Correspondence: Dr. Vasant D. Barhate**

V.E. S. College of Arts, Science and Commerce, Sindhi Society Chembur, Mumbai - 400071, India.

Article Received on 19/12/2015

Article Revised on 11/01/2016

Article Accepted on 31/01/2016

ABSTRACT

A Schiff base ligand derived from o - methoxy benzaldehyde and 2 - amino phenol and its transition metal complexes with Co(II) have been synthesized which was characterized by elemental analysis, molar conductance, magnetic susceptibility, electronic spectra, IR and ESR spectroscopy. The metal complex is coloured and solid in nature. The Schiff base ligand exhibits bidentate nature, coordination through azomethine nitrogen, phenolic oxygen to the metal ions. On the basis of electronic spectral analysis and magnetic susceptibility values, geometry of complex was proposed to be tetrahedral. The molar conductivity data of complex suggests its non - electrolytic nature. The ligand and metal complex have been screened for their microbiological activity.

KEYWORDS: Schiff base (NOMBAP), Spectra, Metal Complex and Microbiological activity.

INTRODUCTION

Schiff bases are an important class of ligand in the field of coordination chemistry.^[1] During the past two decades, considerable attention has been paid to the chemistry of the metal complexes of Schiff bases containing nitrogen and other donor atoms.^[2-5] A large number of Schiff bases and their metal complexes have been found to possess important biological and catalytic activity.^[6-7] Due to their great flexibility and diverse structural aspects, wide ranges of Schiff bases have been synthesized and their complexation behaviour was studied.^[8] Schiff bases are also used as an efficient reagent in trace analysis of some metal cations.^[9-10]

In the present paper, metal complexes of Co (II), with Schiff base ligand derived from o-methoxy benzaldehyde and 2- amino phenol have been synthesized and these complexes were characterized by elemental analysis, magnetic susceptibility, molar conductivity measurements, electronic spectra, IR, and ESR spectrum to determine the mode of bonding and geometry. The microbiological activities of the ligand and their metal complexes have also been studied.

MATERIALS AND METHODS


Instrumentation

All the used chemicals and solvents were of AR grade and were used without further purification. All the reagents used for the preparation of Schiff base and metal salts were obtained from Lobachemie. The percentage compositions of the elements (CHNO) of the compounds were determined using an element analyzer

CHNO model Flash EA 1112 series. The metal contents of the complexes were analyzed by ICPAES (Inductive coupled plasma atomic emission spectroscopy) in an Iris Intrepid II XSP model instrument. The Infrared spectra were recorded as potassium bromide (KBr) discs using a 8400 FTIR Perkin Elmer Spectrophotometer. The electronic Spectrum of complexes were recorded on ELICO SL -159 UV-Vis Spectrophotometer. The Electron spin resonance spectra (ESR) were recorded on E-112 ESR Spectrometer at I.I.T. Mumbai. Melting point of ligand and their metal complexes were recorded using Differential Scanning Calorimeter (DSC) instrument. The molar conductivity measurements were made on ELICO SL - 303 model using methanol as the solvent. Magnetic susceptibility of complex was measured by Gouy's method using mercury tetra isothio cyanato cobaltate as the callibrant.

Synthesis of ligand with [2 - (2 - methoxybenzylidene) amino phenol] (NOMBAP)

The Schiff base ligand NOMBAP was synthesized by refluxing equimolar amount of ethanolic solution of o - methoxybenzyldehyde with 2 - amino phenol for 5 - 6 hours. On cooling the reaction mixture, a sharp yellow crystal product separated out (80%, yield m.p.87⁰-88⁰C) which was collected by filtration. The resulting NOMBAP was recrystallised using aqueous ethanol as the procedure recommended by Vogel.^[11] The product was characterized by elemental and spectral analysis.



Preparation of complexes with Schiff base ligand (NOMBAP)

The complex of Co (II) was prepared by refluxing hydrated cobalt chloride in ethanolic solution of ligand in molar ratio of 1:2 at pH 8 - 9. The resulting mixture was refluxed on water bath for 2 - 3 hrs. After cooling at room temperature, dark brown colored solid separated out which was filtered, washed with water and then with ethanol. The resulting product was recrystallised and dried in vacuum desiccator over CaCl_2 anhydrous (yield 60 - 70%).

Biological Studies

Table:1 The Analytical and physical data of ligand and Metal complexes

Compounds (Colour)	Molecular Weight	M.P.	% Elemental Analysis Found (Calculated)				μ_{eff} B.M.	λ_m $\text{Ohm}^{-1} \text{cm}^2 \text{mol}^{-1}$
			C	H	N	M		
NOMBAP (Yellow)	227.28	87°C	73.12 (73.91)	5.14 (5.719)	6.41 (6.159)	-		
[Co(NOMBAP) ₂] H ₂ O (Dark Brown)	531.496	93°C	63.16 (63.217)	5.195 (5.268)	5.48 (5.26)	12.09 (11.087)	4.41	24.10

Infrared Spectral Analysis

Infrared spectra were recorded with a Perkin Elmer FTIR -8400S Spectrometer (4000-400 cm^{-1}) using KBr pellets. On the basis of this, the presence of important functional groups in the compound can be identified. The Infrared spectrum of the ligand was compared with the spectra of Co(II) complex. The typical IR spectra were shown in Fig. 2 and 3. Data was summarized in Table-2 along their assignment.

The IR spectrum of the ligand shows broad band at 1618.05 cm^{-1} , which was assigned to due ν (C=N) stretching of azomethine group.^[12] In complexes this band was shifted to lower regions, 1601.94 cm^{-1} for Co(II) complex, suggesting the coordination of azomethine nitrogen to metal atom in complexation.^[13] This was due to the donation of electron density from Nitrogen to metal.

IR spectrum of the ligand shows a weak band around 3011.3 cm^{-1} due to intramolecular hydrogen bonded -OH group^[14-15] This band is absent in the spectra of complexes indicating the dissociation of the phenolic proton on complexation and involvement of phenolic anionic oxygen in coordination.^[16] Moreover, the strong

Antibacterial and antifungal activity of ligand and their complexes have been determined by cup plate method. The stock solution of Co (II) compound was prepared by dissolving 20 mg of compound in methanol and the volume was made up to 10ml with same solvent. The stock solution of 2000ppm of each compound thus prepared on active ingredient basis was kept at room temperature till used. Sterile Mueller Hinton agar plates for bacterial test cultures and Sabourauds agar plates for fungus *Candida albicans* were seeded with 1ml of 24 hour old, 0.1 O.D. cultures. Sabourauds agar plates were surface spread with 0.2 ml of 48 hour old *Aspergillus* culture. The compounds to be tested (50 μl) were added to the wells punched in the above media. The plates were incubated at 37°C / R.T. for 48 hours depending on the cultures. The zone of inhibition around the wells was measured in millimeters.

RESULTS AND DISCUSSION

The metal complex is colored, solid, stable at room temperature and hygroscopic in nature. The Analytical and physical data of ligand and Co (II) complex is recorded in Table - 1. The metal complex exhibits 1:2 (metal-ligand) stoichiometry.

band at 1251 cm^{-1} due to C - O (phenolic) in the ligand has been shifted to the 1260 - 1287 cm^{-1} in the spectra of complex.^[16]

New bands were observed between 400 - 500 cm^{-1} region in the complex, which were absent in the spectrum of ligand. The bands between 439 - 444 cm^{-1} were assigned to stretching frequencies of $\nu(\text{M-O})$ ^[17-18], the band between 480 - 487 cm^{-1} have been assigned to the stretching frequencies $\nu(\text{M-N})$ respectively^[19-21]

The IR spectrum of the ligand and complex shows band at 2936 - 2933 cm^{-1} , which was assigned to due $\nu(\text{C-H})$ stretching of benzene group.^[22]

The IR spectrum of Co (II) metal complexes exhibit a broad and strong absorption band in the region 3491.94 cm^{-1} confirms the presence of ν (OH) absorption for lattice water^[22]

Therefore from IR spectra it is concluded that ligand behaves as {uni negatively} bidentate ligand co-ordinated to metal ion via azomethine (N) and phenolic (O).

Table-2: The Important IR bands of Ligand and Their Metal Complexes

Compound	$\nu(-OH)$ cm^{-1}	$\nu(OCH_3)$ cm^{-1}	$\nu(C=N)$ cm^{-1}	$\nu(C-O)$ stretching cm^{-1}	$\nu(M-N)$ cm^{-1}	$\nu(M-O)$ cm^{-1}
NOMBAP	3011.36phenolic	1020.39	1618.05	1251.8 ⁸ .	-	-
Co(NOMBAP) ₂ H ₂ O	3451.94 H ₂ O	1029.40	1601.94	1261.01	485.22	439.39




Fig 1: Spectra of Ligand


Figure-4: IR Spectrum of Co(NOMBAP)₂·H₂O complex.

Fig 2: IR Spectra of CO (II) Complex

Molar Conductance

The molar conductance values of 1×10^{-3} M solution of metal complex in methanol were measured at room temperature (Table - 1) using an ELICO Conductivity meter (cell constant 1.0 cm^{-1}). These values were compared with known molar conductivities^[23-24] which indicates the non - electrolytic nature^[25] of the complex.

Electronic Spectra and Magnetic Susceptibility Measurements of Complexes

In UV-Visible electromagnetic radiation, the transitions are associated with the electronic energy levels of the compound under investigation. The electronic spectra were recorded on ELICO SL - 159 UV-Visible Spectrophotometer. The transition metal ions occur in a

variety of structural environments. Because of this, the electronic structures are extremely varied. The electronic structures have been identified with UV-Visible spectroscopy. The nature of ligand field around the metal ion and the geometry of complexes have been deduced from the electronic spectra. Electronic spectra of complexes were recorded by using solution of complexes

The Co (II) complex display band at 20408.1632 cm^{-1} (490 nm) and which may be assigned to ${}^4\text{A}_2 \rightarrow {}^4\text{T}_1$ transition^[26] suggesting tetrahedral geometry of complexes. The room temperature magnetic moment value of and Co (II) complex lies in the range of 4.41 B.M. attributed to Tetrahedral structure. Electronic Spectra and Magnetic Moment of the Complexes are listed in Table - 4.

Table - 4. Electronic Spectra and Magnetic Moment of the Complexes

Compound	Band, λ max (nm)	Assignments	μ_{eff} (B.M.)
[Co (NOMBAP) ₂].H ₂ O	490	${}^4\text{A}_2 \rightarrow {}^4\text{T}_1$	4.41

Electronic Spin Resonance (ESR) Spectra

ESR spectra of complexes provide information about the extent of the delocalization of unpaired electron. In the present study, the ESR spectra of complexes have been recorded on X - band at frequency 9.5 GHZ under the magnetic field strength 3400 gauss in DMF at room temperature (Fig. 4) and their g_{\parallel} , g_{\perp} , g_{avg} , Δg , and G values have been calculated. The values of ESR parameters for Co(II) complex viz. g_{\parallel} , g_{\perp} , g_{avg} , Δg , and G presented in Table - 5.

ESR spectra of the complex revealed two g values (g_{\parallel} and g_{\perp}). the trend $g_{\parallel} > g_{\perp}$ shows that the unpaired electron is delocalized in $dx^2 - y^2$ orbital in the ground state of metal and spectra are characteristics of axial symmetry. The parameter g_{avg} was obtained by equation

$[(g_{\text{avg}}) = 1/3 (g_{\parallel} + 2g_{\perp})]$. Kvelson&Neiman^[27] have reported that, $g_{\parallel} > 2.3$ is the characteristics of an ionic environment and $g_{\parallel} < 2.3$ indicates a covalent character in metal ligand bonding. Applying this criterion, in the present compounds g_{\parallel} indicate the prevalence of covalent character in metal - ligand bond.^[28] The exchange coupling interaction has been explained by Hathaway expression $G = (g_{\parallel} - 2) / (g_{\perp} - 2)$.

According to Hathaway if the value of G is greater than four ($G > 4$), the exchange interaction is negligible whereas when the value of G is less than four ($G < 4$), a considerable exchange interaction is indicated in the complex. It is observed that G value for this complex is less than four ($G < 4$), indicating considerable exchange interaction the complex^[29-30]

Table – 5. Electronic Spin Resonance Parameters of The Complex

Metal Complexes	g_{\parallel}	g_{\perp}	g_{avg}	Δg	G
[Co (NOMBAP) ₂].H ₂ O	2.11	2.05	2.09	0.06	2.2





Fig 3 Electronic Spin Resonance (ESR) Spectra of Co

Proposed Structure of Complexes

The present study clearly indicates that the ligand forms stable coordinate complexes with the metal ions under investigation which is evidenced by micro - analytical,

magnetic and spectroscopic data. The probable structures of the complexes under investigation on the basis of the above experimental evidence can be shown as (Fig. 4)

Fig 4 $\text{H}_2\text{O M} = \text{Co (II)}$

CONCLUSIONS

It can be concluded from the results of above study that Schiff base ligand NOMBAP [N - (o - methoxybenzaldehyde) 2- amino phenol] acts as a very good bidentate complexing agent towards transition metal ions during complexation. Analytical data of complexes are in good agreement with their molecular formula. The involvement of azomethine (N), phenolic (O) of the bidentate Schiff bases to the metal ions was confirmed by comparing the IR data of the ligands with those of the metal complexes. Further conclusive evidence of the coordination of these Schiff-bases with the metal ions was shown by the appearance of new bands due to $\nu(\text{M}-\text{N})$ and $\nu(\text{M}-\text{O})$ in the metal complexes. The electronic spectral and magnetic susceptibility measurements were used for assigning the geometry of metal complexes. Complexes of Co (II) with NOMBAP were proposed to be tetrahedral in nature.

ESR spectra of complex provide information about the extent of the delocalization of unpaired electron. On the basis of these findings the structures have been proposed

for the complex which are in good agreement with theoretical consideration. (Fig.4).

Biological activity

Schiff base ligand and their metal complexes were screened for antibacterial activity against gram positive bacteria (staphylococcus aureus and corynebacterium diphtheria), gram negative bacteria (Escherichia coli and Klebsiella pneumonia) and antifungal activity against (Candida albicans and aspergillus species) by cup plate method^[31]. The results of the biological activity of the metal complexes indicated the following facts.

A comparative study of the ligand and their complexes indicates that the metal chelates exhibited higher antibacterial and antifungal activity than that of the free ligand recorded in Table - 5. The increase in the biological activity of metal chelates was found due to the effect of metal ion on the metal chelates which could be explained on the basis of overtones concept and chelation theory. On chelation the polarity of the metal ion reduced to a greater extent due to the overlap of the ligand orbital and partial sharing of positive charges of metal ion with donor groups. It was further noted that the delocalization of electrons over the whole chelate ring enhanced the lipophilicity of the complexes. This increased lipophilicity^[32-33] enhanced the penetration of the complexes into lipid membrane and blocking the metal sites on enzymes of microorganism. The activity was compared with zone of inhibition was measured in millimeters.

Table 6: Zone of inhibition of growth in millimeters after 48 hours of incubation

Complexes	Cultures					
	S.a	C.d	E.c	K.p	C.a	Asp.
NOMBAP	21	8	09	8	-	10
$[\text{Co}(\text{NOMBAP})_2]\text{H}_2\text{O}$	34	42	-	NG	NG	15
SOLVENT	-	-	-	-	-	-

ACKNOWLEDGEMENT

The authors are thankful to SAIF, I.I.T. Mumbai and Dr. Y.S.Lakshmi Narasimham, Asst. Director, Lupinpharma, Pune, providing spectral and analytical data. We are also thankful to the principal, Dr. (Mrs.) J. K. Phadnis, V. E. S. College, Chembur, Mumbai for providing necessary research facilities and Shewta Patil, Assistant prof. Dept. of Microbiology, for providing biological studies.

REFERENCES

- Shirodkar SG, Mane PS, Chondhekar TK (Synthesis and fungitoxic studies of Mn(II), Co(II), Ni(II) and Cu(II) with some heterocyclic Schiff base ligands). Indian j. Chem., 2001; 40: 1114-1117.
- Djebbar SS, Benali BO, Deloume JP, (Synthesis, characterization and electrochemical behavior of copper (II) complexes with linear and tripodal tetradentate ligands derived from Schiff base). Polyhedron, 1997; 16: 2175-2182.
- Bhattacharya P, Parr J, Rossa AT, (First synthesis of a unique lead Schiff base complex) J. Chem. Soc., Dalton, 1998: 3149.
- He L, Gou SH, Shi QF, (The formation of a Schiff base intermediate: a nickel (II) complex of an asymmetric tripodal ligand). J. Chem. Crystallogr, 1999; 29: 207.
- Wu J C, Tang N, Liu WS, Tan MY, Chan ASC. (Intramolecular hydrogen bond self-template synthesis of some new Robson-type macrocyclic ligands) Chin. Chem. Lett., 2001; 12: 757-760.
- Wahhenri li kam. Tagenine J, Minu B. (Synthesis and antibacterial study of Schiff base complexes 2,3-diamino pyridine and o vanillin). Indian j. Chem, 2001; 40: 999-1003.
- Soliman AA, Mohammed GG, (Study of the ternary complexes of copper with salicylidene-2-

- aminothiophenol and some amino acids in the solid state). *Thermochim. Acta*, 2004; 421: 151-159.
8. Vaghasiya Y K , Nair R , Soni M , Baluja S, Chanda S (Synthesis, structural determination and antibacterial activity of compounds derived from vanillin and 4-aminoantipyrine). *J. serb. Chem. Soc.*, 2004; 69: 991-998.
 9. Mandhare DB, Barhate VD. (Development of extractive spectrophotometric method for the determination of nickel (II) with Schiff base 2-[(2-hydroxyphenylimino)methyl]-4-nitrophenol). *J Chem Pharm Res.*, 2015; 7: 1069-73.
 10. Barhate VD, Patil MR. (Extraction and spectrophotometric determination of cobalt (II) with p-methylisonitrosoacetophenone). *current science*, 1989; 58: 6.
 11. Vogel A I. *Practical Organic chemistry*, 5th Ed., London; Longman group limited, 1978.
 12. Lever M. *Anal.Chem.Acta*, 1973; 65: 311.
 13. Mahmoud NA, Hafidh FR, Majeed AR (synthesis and characterization of some transition metal complexes with tridentate N3donor schiff base derived from 2-aminothiazole) *Eur. Chem. Bull.*, 2014; 3(6); 559-562.
 14. BellamyLJ, *infrared spectra of complexes molecules*, New York; wiley, 1954.
 15. Silverstein RM & Webster FX, *Spectrometric Identification of organic compounds*, 6th ed., John wiley & sons: 87 3.6.8
 16. Esmadia F, Khabourb OF, Albarqawib AI., Ababneha M, Talib MA. (Synthesis and Characterization of Some Transition Metal Complexes of Thiocarbohydrazone Schiff Bases) *Jordan Journal of Chemistry*, 2013; 8(1): 31-43.
 17. Joshi KR, Rojivadiya AJ, Pandya JH (Synthesis and Spectroscopic and Antimicrobial Studies of Schiff Base Metal Complexes Derived from 2-Hydroxy-3-methoxy-5-nitrobenzaldehyde) *International Journal of Inorganic Chemistry*, 2014; 8 .
 18. MabryTJ, Markham KR , *Flavonoids*, edited by Harborne JB, Mabry H , Chapman and Hall, Landon, 1975; 78.
 19. Mahmoud NA, Hafidh FR .Majeed AR (synthesis and characterization of some transition metal complexes with tridentate N3donor schiff base derived from 2-aminothiazole). *Eur. Chem. Bull.*, 2014; 3(6): 559-562.
 20. Jadeja RN, Parmar NJ. (Oxovanadium(IV), Cr(III), Fe(III), Fe(II), Ni(II), Cu(II), Zn(II), and UO₂(VI) chelates from onno donor Schiff base ligand) *Synthesis and Reactivity in Inorganic and Metal-Organic Chemistry*, 2005; 35: 111-117.
 21. Anitha C, Sheela CD, Tharmaraj P, Shanmugakala R (Studies on Synthesis and Spectral Characterization of Some Transition Metal Complexes of Azo-Azomethine Derivative of Diaminomaleonitrile). *International Journal of Inorganic Chemistry*, 2013; 10.
 22. Esmadia F, Khabourb OF, Albarqawib AI, Ababneha M , Talib MA.(Synthesis and Characterization of Some Transition Metal Complexes of Thiocarbohydrazone Schiff Bases) *Jordan Journal of Chemistry*, 2013; 8(1): 31-43.
 23. Quagliano JV, Fujita J, Franz G, Phillips DJ, Walmsley JA, Tyree SY. (The Donor Properties of Pyridine N-Oxide). *J. Am. Chem. Soc.*, 1961; 83: 3770 -3773
 24. Thaker BT, Surati KR, Oswal S, Jadeja RN, GuptaVK. (Synthesis, spectral, thermal and crystallographic investigations on oxovanadium (IV) and manganese (III) complexes derived from heterocyclic β -diketone and 2-amino ethanol). *Structural Chemistry*, 2007; 18: 295-310.
 25. El- Tablas, Ayad ML. Synthesis and reactivity in inorganic, metal organic and nano metal chemistry, 2003; 33: 369.
 26. Mishra AP., Tiwari A., Gupta SK., and Jain R. (Synthesis, Spectral and Antimicrobial Studies of some Co (II), Ni (II) and Cu (II) Complexes Containing 2-Thiophenecarboxaldehyde Moiety) *E-Journal of Chemistry* 2012; 9(3): 1113-1121.
 27. Maki AH, Mcgarvey BR. (Calculation of NMR and EPR Parameters: Theory and Applications edited by Martin Kaupp, Michael Bühl, Vladimir G. Malkin) *J.Chem. Phys.*, 1958; 29: 31- 35.
 28. Halcrow MA, Chia LML, Liu X, McInnes EJJ, Davies JE. (Spectroscopic characterisation of a copper (II) complex of a thioether-substituted phenoxyl radical: a new model for galactose oxidase). *Chem. Commun.*, 1998; 2465-2466.
 29. Mishra AP, Khare M, Gautam SK. (Synthesis, physico-chemical characterization, and antibacterial studies of some bioactive Schiff bases and their metal chelates). *Synthesis and Reactivity in Inorganic and Metal-Organic Chemistry*, 2002; 32(8): 1485-1500.
 30. Abragam A, Bleaney B. *EPR of Transition ions*, oxford; charedon, 1970
 31. Sandhar RK, Sharma JR, Manrao MR.(Synthesis and Fungitoxicity of C-phenyl Substituted Benzal-4-fluoroanilines). *Pesticide Research Journal*, 2005; 17: 9-11.
 32. Sandhar RK, Hothi HS, Sharma JR, Manrao MR (Synthesis and Antifungal Potential of Fe (III) Complexes of 2-Hydroxy-1 -naphthalanilines). *Pesticide Research Journal*, 2006; 18: 4-6.
 33. Sheikh J' Juneja H, Ingle V, Ali P, Hadda TB. (Synthesis and in vitro biology of Co (II), Ni (II), Cu(II) and Zinc(II) complexes of functionalized beta-diketone bearing energy buried potential antibacterial and antiviral O,O pharmacophore sites). *Journal of Saudi Chemical Society*, 2013; 17(3): 269-276.