



“COMPARISON OF VARIOUS PERIODS OF PREOXYGENATION”

Dr. Ranjeeta Aske^{*1}, Dr. Aditya Agarwal² and Dr. Yogita Nimje³

¹Assistant Professor, Department of Anaesthesiology, Gandhi Medical College and Hamidia Hospital, Bhopal (M.P.)

²Professor, Department of Anaesthesiology, Gandhi Medical College and Hamidia Hospital, Bhopal (M.P.)

³Resident, Department of Anaesthesiology, Gandhi Medical College and Hamidia Hospital, Bhopal (M.P.)

***Corresponding Author: Dr. Ranjeeta Aske**

Assistant Professor, Department of Anaesthesiology, Gandhi Medical College and Hamidia Hospital, Bhopal (M.P.).

Article Received on 05/03/2016

Article Revised on 26/03/2016

Article Accepted on 17/04/2016

ABSTRACT

Background: The present study compare the effects of varying periods of preoxygenation on oxygen saturation and arrive at a value of the optimal duration of preoxygenation to tide over the period of hypoxia that occurs during the process of intubation. **Methods:** 60 ASA grade I and II patients in the age group of 20 to 50 years were studied under three equal groups. Group I patients received preoxygenation in the form of four vital capacity breaths. Group II and III patients received preoxygenation for 3 and 5 minutes respectively. Oxygen saturation was monitored by pulse oximeter. **Results:** The results were analysed using the “analysis of variance” method. It was observed that patients in group I after one minute of apnea and period of laryngoscopy and intubation had a fall in saturation to a mean of $84.32\% \pm 1.2$ in comparison to $95.92\% \pm 0.8$ and $99.0\% \pm 0.2$ in group II and III respectively. It was also observed that for the saturation to come to 100% in the post intubation phase, the patients took an average of 20.80, 20.75 and 20.25 seconds in the three groups respectively. **Conclusion:** Routine preoxygenation of 5 minutes must be carried out in all patients scheduled for elective surgery. The degree of desaturation occurring in patients who are adequately pre-oxygenated for five minutes is significantly lower than that in the four vital capacity breaths and three minutes pre-oxygenated groups.

KEYWORDS: Preoxygenation, various periods.

INTRODUCTION

Preoxygenation with 100% oxygen replaces nitrogen in the functional residual capacity and increases oxygen stores thus prolonging the safe duration of apnea after induction of general anaesthesia and administration of muscle relaxants. Thorpe et al in their study have shown that 98% of patients not preoxygenated developed hypoxia during intubation following induction of general anaesthesia.^[1] Numerous studies have also emphasized the need for routinely pre-oxygenating any patient undergoing general anaesthesia.^[2-7] The present study sought to compare the effects of varying periods of preoxygenation on oxygen saturation and arrive at a value of the optimal duration of pre-oxygenation to tide over the period of hypoxia that occurs during the process of intubation.

MATERIAL AND METHOD

After obtaining institutional ethical committee clearance and written informed consent from all of the patients. 60 patients of ASA grade I and II, age between 20-50 years scheduled for elective surgery. The patients were randomly allocated into the three groups of 20 each, Patients in group I received preoxygenation in the form of four vital capacity breaths, group II patients received

preoxygenation for three minutes and group III patients received five minutes of preoxygenation with 100% oxygen at 10 liters with the aid of an anatomical face mask prior to the induction of general anaesthesia. Smokers, pregnant women, patients with cardiorespiratory diseases, patients with a difficult airway, patients with neuromuscular disorders, patients with haemoglobinopathies excluded from the study.

In the operation theatre after securing an intravenous access the pulse rate, ECG, blood pressure and SpO₂ values were recorded on room air. All patients were pre-medicated with intravenous injection Glycopyrrolate 0.2 mg, intravenous injection Midazolam 0.02 mg/kg and intravenous injection Fentanyl 2 µg/kg. Depending on the group the patients belonged to preoxygenation. Patients in all the three groups were induced with injection of Propofol 2.5 mg/kg intravenously followed by injection Suxamethonium 2 mg/kg. The patients were then kept apneic for a period of one minute. At the end of one minute, time took for laryngoscopy and intubation were noted and the patients were intubated with an adequate sized endotracheal tube. The patients were then ventilated with 100% oxygen and 0.5% Halothane. The heart rate, ECG, blood pressure and SpO₂ values of all


the patients were recorded every 15 seconds till the saturation reached 100% following this nitrous oxide was administered and the surgery was commenced.

RESULTS


A total of 60 patients of ASA physical status I and II were enrolled in this study. The mean age in years was 30.96 ± 7.97 . The mean weight of the patients was 47.98 ± 5.15 kilograms with male to female ratio of 40:60. It was observed that in patients of all groups after one minute of apnea post induction time took for laryngoscopy and intubation was 20.6 seconds. In group

I patients, there was a fall in oxygen saturation to a mean of $84.32\% \pm 1.2$ whereas it was $95.92\% \pm 0.8$ in group II and $99.0\% \pm 0.2$ in group III. It was also observed that for the saturation to come to 100% in the post intubation phase the patients took an average of 20.80, 20.75 and 20.25 seconds in the three groups respectively. It was also observed that there was no statistically significant variation in heart rate and blood pressure between these three groups. The results were analyzed using the "Analysis of variance" (ANOVA) method. All data were presented as means $\pm$ standard deviation.


OBSERVATION CHARTS




Line diagram -1: SpO₂ values in the pre-induction, apnea periods and post intubation periods.



Line diagram -2: Pulse rate values in the pre-induction, apnea periods and post intubation periods.



Line diagram -3: Systolic blood pressure values in the pre-induction, apnea periods and post intubation periods.



Line diagram -4: Diastolic blood pressure values in the pre-induction, apnea periods and post intubation periods.

DISCUSSION

Pre-oxygenation before induction of general anaesthesia is used less frequently despite the fact that hypoxemia does occur during the period of laryngoscopy. The reasons for hypoxia included a decreased functional residual capacity, respiratory depression, decreased cardiac output and ventilation perfusion mismatch. Pre-oxygenation does protect against hypoxia during intubation and has been recommended by several authors for all cases. There are many possible reasons for the reluctance to pre-oxygenate routine cases which may include a perceived lack of time on a busy operating list and the fact that holding an anatomical face mask is a labour intensive task preventing the anaesthesiologist or his assistant from carrying out any other activity.

This study was carried out to evaluate the need for pre-oxygenating a patient prior to the induction of general anaesthesia and also to assess the speed of desaturation and resaturation in patients who receive varying periods of pre-oxygenation.

Thopre et al determined the incidence of hypoxia using pulse oximetry during induction of anaesthesia.^[1] It was concluded that 98% of the patients not pre-oxygenated adequately developed hypoxia during intubation. Skea et al observed that in pre-oxygenated patients the SpO₂ values remained at a mean of 98% during the apneic period of intubation whereas it fell to a mean of 86% in the non pre-oxygenated group.^[2] So, it was suggested that serious consideration be given to the routine use of pre-oxygenation prior to the induction of general anaesthesia. Hett et al observed that 97% of patients in whom no adequate pre-oxygenation was done had a fall in saturation to a mean of 86% and in patients in whom three minutes of pre-oxygenation was done had a mean saturation of 98% during the apneic period of intubation. In our study it was found that in group I in whom pre-oxygenation was instituted in the form of four vital capacity breaths, after a minute of apnea in the induction phase resulted in a fall of saturation to a mean of 84.32% $\pm$ 1.2. This was in sharp contrast to the groups in whom pre-oxygenation was given for three and five minutes respectively. In these latter groups the saturation fell to a mean of 95.92% $\pm$ 0.8 and 99.0% $\pm$ 0.2 respectively.

The results in our study correlate well with those obtained in the study conducted by Hett et al, wherein the patients who did not receive adequate pre-oxygenation had a fall in saturation to a mean of 84% and patients who had three minutes of pre-oxygenation had a fall in saturation to a mean of 98%.^[3, 4] These results are also in accordance with the study done by Skea et al who reported a mean fall in oxygen saturation to 81% in the group not adequately pre-oxygenated to a mean of 98% in the pre-oxygenated group. Motoyama et al reported that a reasonable amount of desaturation was noticed in patients in whom routine adequate pre-oxygenation was not carried out.^[5] Laycock et al and Weitzner et al, also noticed a significant amount of desaturation in patients in whom adequate pre-oxygenation was not advocated prior to induction of general anaesthesia.^[6, 7]

Campbell IT, Beauti PG inferred that denitrogenation does not equate exactly with preoxygenation in terms of prolonging acceptable apnoea time and that 1 min of 100% oxygen does allow 3 minutes of apnoea 3 min of oxygen, almost doubles this time.^[8]

Jaideep Pandit et al compared 3 minutes of tidal breathing with four Deep Breaths (DB) in 300 sec/8 DB in 60 sec and found out that 3 min Tidal Volume (TV) or 8 DB are superior & equally effective that 4 DB in 30 Sec.^[9] McGowan P, Skinner A. et al also studied adequate preoxygenation by telling the importance of a good face mask seal.^[10]

Usharani Nimmagadda et al in their study found out that preoxygenation via circle absorber & nasooral system when compared with five types of reservoir bags and inferred that these bags should not be used for preoxygenation.^[11] J.W. Mc Crory et al studied four different techniques of preoxygenation in 20 volunteers by continuous analysis of respired gases. They compared 3 min of Tidal Volume (TV) from McGill's circuit or four Vital capacity (VC) breathing from a non breathing system and was found to be equally effective. Maximum expiration before application of facemask results in improved oxygenation when four VC breaths were taken from McGill system.^[12]

Gambie Market al Compared 3 min of spontaneous preoxygenation and 4 full deep breaths. Similarly Tanoubi et al did a randomized double blind cross over trial. They applied inspiratory supports & in other group they applied positive end expiratory pressure (PEP) and studied how they increase the effectiveness of preoxygenation.^[13]

A. Herringer told the importance of PEEP during induction of anaesthesia may prevent atelectasis formation in the lungs. This technique increases the functional residual capacity (FRC) and prolongs non hypoxic apnoea duration by >2 min.^[14] Tanoubi I, Drolet P, Fortier LP, Donati F. did a randomized, double blind, cross-over trial in which they studied inspiratory support versus spontaneous breathing during preoxygenation in healthy subjects.^[15]

Chintophe Baillard studied that preoxygenation using non invasive ventilation (NIV) is more effective at reducing arterial oxyhaemoglobin desaturation than usual method.^[16]

Baillard C, Fosse JP, Sebbane M, Chanques G, Vincent F, Courouble P et al. found out that noninvasive ventilation improves preoxygenation before intubation of hypoxic patients Nimmagadda U, Chiravuri SD, Salem MR, Joseph NJ, Wafai Y, Crystal GJ et al. also explained the role of preoxygenation with tidal volume and deep breathing techniques and the impact of duration of breathing and fresh gas flow.^[17, 18] The above observations were similar to what we found out in our study.

CONCLUSION

In conclusion we suggest that serious consideration should be given to the routine use of pre-oxygenation of five minutes before induction of general anaesthesia. As the degree of desaturation occurring in patients who are adequately pre-oxygenated for five minutes is significantly lower than that in the four vital capacity breaths and three minutes pre-oxygenated groups. Also the vital parameters such as the heart rate and blood pressure remain closer to the baseline values in the pre-oxygenated group which could make a significant degree of difference in a compromised patients.

REFERENCES

1. Thorpe CM, Gauntlet IS. Arterial oxygen saturation during the induction of anaesthesia. *Anaesthesia*, 1990; 45: 1012-1015.
2. Skea G, Jones A, Snowdon SL. Routine pre-oxygenation. *Anaesthesia*, 1991 Jun 1; 46(6): 510-1.
3. Hett DA, Geraghty IF, Radford R. Routine Pre-oxygenation. *Anaesthesia*, 1993; 48: 76-77.
4. Hett DA, Geraghty IF. Routine Pre-oxygenation. *Anaesthesia*, 1994; 49: 157-159.
5. Motoyama. Hypoxemia during induction of general anaesthesia. *Anaesth Analg*, 1986; 654: 267-272.
6. Laycock. Hypoxemia during induction of anaesthesia for routine elective surgery. *Anaesthesia*, 1988; 43: 981-984.
7. WEITZNER SW, KING BD, IKEZONO E. The rate of arterial oxygen desaturation during apnea in humans. *The Journal of the American Society of Anesthesiologists*. 1959 Sep 1; 20(5): 624-7.
8. Campbell IT, Beatty PC. Monitoring Preoxygenation. *Br J Anaesth*, 1994; 72: 3-4.
9. Pandit JJ, Duncan T, Robbins PA. Total Oxygen Uptake with Two Maximal Breathing Techniques and the Tidal Volume Breathing Technique A Physiologic Study of Preoxygenation. *The Journal of the American Society of Anesthesiologists*, 2003 Oct 1; 99(4): 841-6.
10. McGowan P, Skinner A. Preoxygenation- the importance of a good face mask seal. *Br J Anaesth*, 1995; 75: 777-8.
11. Nimmagadda U, Salem MR, Joseph NJ, Lopez G, Megally M, Lang DJ et al. Efficacy of preoxygenation with tidal volume breathing. Comparison of breathing systems. *Anaesthesiology*, 2000; 93: 693-8.
12. McCrory JW, Matthews JN. Comparison of four methods of preoxygenation. *British journal of Anaesthesia*, 1990 May 1; 64(5): 571-6.
13. Gambie AM, Hertzka RE, Fisher DM. Preoxygenation techniques: comparison of three minutes and four breaths. *Anaesth Analg*, 1987; 66: 468-70.
14. Herriger A, Frascarolo PH, Spahn DR, Magnusson L. The effect of positive airway pressure during pre-oxygenation and induction of anaesthesia upon duration of non-hypoxic apnoea. *Anaesthesia*, 2004 Mar 1; 59(3): 243-7.
15. Tanoubi I, Drolet P, Fortier LP, Donati F. [Inspiratory support versus spontaneous breathing during preoxygenation in healthy subjects. A randomized, double blind, cross-over trial]. *In Annales francaises d'anesthesie et de reanimation*, 2010 Mar; 29(3): 198-203.
16. Baillard C, Depret F, Levy V, Boubaya M, Beloucif S. Incidence and prediction of inadequate preoxygenation before induction of anaesthesia. *In Annales francaises d'anesthesie et de reanimation*, 2014 Apr 30; 33(4): e55-e58. Elsevier Masson.
17. Baillard C, Fosse JP, Sebbane M, Chanques G, Vincent F, Courouble P et al. Noninvasive ventilation improves preoxygenation before intubation of hypoxic patients. *Am J Respir Crit Care Med*, 2006; 174: 171-7.
18. Nimmagadda U, Chiravuri SD, Salem MR, Joseph NJ, Wafai Y, Crystal GJ et al. Preoxygenation with tidal volume and deep breathing techniques: the impact of duration of breathing and fresh gas flow. *Anaesth Analg*, 2001; 92: 1337-41.