



**CYTOTOXICITY SCREENING OF METHANOLIC BULB EXTRACT OF  
*BELLICORYNE PLUMBAGINIFOLIA* AGAINST VARIOUS MAMMALIAN CANCER  
CELL LINES**

**\*V. Ramamurthy and K. L. Naveen**

P.G & Research Department of Biochemistry, Marudupandiyar College, Vallam, Thanjavur, 613 403, Tamil Nadu, India.

**\*Corresponding Author: V. Ramamurthy**

P.G & Research Department of Biochemistry, Marudupandiyar College, Vallam, Thanjavur, 613 403, Tamil Nadu, India.

Article Received on 17/05/2017

Article Revised on 07/06/2017

Article Accepted on 28/06/2017

**ABSTRACT**

Cancer is a major public health burden in both developed and developing countries. Anticancer activity is the effect of natural and synthetic or biological and chemical agents to reverse, suppress or prevent carcinogenic progression. Several synthetic agents are used to cure the disease but they have their toxicity and hence the research is going on to investigate the plant derived chemotherapeutic agents. Therefore an attempt has been made to review different in vitro method for estimating anticancer properties of natural products from medicinal plants. In this study the anticancer potentials of plants was investigated against Dalton's lymphoma (DL), MCF-7 and HeLa cells in vitro. Cytotoxicity of various plant extracts was determined by MTT assay. The results showed that the methanolic bulb extract of *Bellincoryne plumbaginifolia* possessed a moderate amount of anticancer activity and the IC<sub>50</sub> value was recorded. The most potent anticancer activity was observed with the methanolic bulb extract of *Bellincoryne plumbaginifolia* with IC<sub>50</sub> values of 51.2 $\mu$ g/ml and 53.1 $\mu$ g/ml and 48.7 $\mu$ g/ml on DL, MCF-7 and HeLa cells respectively. Phytochemical analyses revealed the presence of large amount of alkaloids and flavonoids in the potent plant extracts which may be suggested to play an important role in their anticancer activities. The methanolic extract would be studied further for isolation and characterization of active components for lead optimization studies.

**KEYWORDS:** Anticancer, cytotoxicity, DL, MCF-7 and HeLa cells, MTT assay.

**INTRODUCTION**

Ayurveda, a traditional Indian medical practice using plant drugs has been successful from very early times in using these natural drugs and preventing or suppressing various tumours with different lines of treatment (Balachandran and Govindarajan, 2005). In India, people of different ethnic groups inhabiting various terrains, possess their own distinct culture, religious rites, food habit and a rich knowledge of traditional medicine (Parinitha et al., 2005). They practice herbal medicine to cure a variety of diseases. Natural products, especially plants have been used in the treatment of various diseases for thousands of years.

Since time immemorial man has used parts of plants in treatment and prevention of many ailments (Chah et al., 2006). Our ancestors made new discoveries of the healing power of plants through trial and error. Although some of the therapeutic properties attributed to plants have proven too erroneous, medicinal plant therapy is based on the empirical finding of hundreds and thousands of years (Gurib- Fakim, 2006). WHO has

estimated that at least 80% of all the global inhabitants rely on traditional systems of medicine for their primary health needs and these systems are largely plant based. Ethno medicines have received renewed global attention of scientists in India and abroad because of their wide local acceptability, and providing leads to the discovery of new drugs of plant origin.

Cancer is one of the major human diseases and causes large suffering and economic loss world-wide. Chemotherapy is one of the methods of treating cancer. However the chemotherapeutic drugs are highly toxic and have devastating side effects. Various new strategies are being developed to control and treat several human cancers (Modha and Modha, 2007). Over 60% of anticancer drugs available in the market are of natural origin. Natural products are also the lead molecules for many of the drugs that are in use (Cragg et al., 1997). Therefore, the phytochemicals present in several herbal products and plants may have the potential to act as preventive or therapeutic agents against various human cancers (Modha and Modha, 2007). The increased

popularity of herbal remedies for cancer therapy perhaps can be attributed to the belief that herbal drugs provide benefit over that of allopathy medicines while being less toxic (Gupta et al., 2004). Since the conventional therapies have devastating side effects, there is a continuous need for search of new herbal cures of cancer (Aquil et al., 2006).

Apoptosis, or programmed cell death, is one of the most finely coordinated regulatory functions for maintenance of the homeostasis in the living organism. It involves the continuous checking of the cellular integrity and cascade-like events of self destruction when the integrity of the organism is endangered. Morphological hallmarks of apoptosis are nuclear condensation, cell shrinkage, membrane blebbing and the formation of apoptotic bodies. These changes are accompanied by biochemical features, including DNA fragmentation and the proteolytic cleavage of a variety of intracellular substrates. Caspases are the best-characterized enzymes that perform this, while the Bcl-2 family is the principal set of proteins that regulate the apoptotic cascade.

*Bellicoryne plumbaginifolia* (Liliaceae) is extensively used in most of the Indian herbal pharmaceuticals and nutraceuticals (Brisca Renuga and Mary Mettilda Bai, 2013). Juice of pounded bulb is given every half an hour duration (depends on the severity of bites) for snake bite. Leaf paste is applied on the spider bitten area. Paste of bulb is applied over the head for mental disorder. The present investigation was taken up for evaluating the antiproliferative potential possessed by the 50% ethanol extract of Bulbs of *Bellicoryne plumbaginifolia* against various human cancer cell lines.

## MATERIALS AND METHODS

### Collection of plant samples

The bulb of the medicinal plants was used for the present study, the plants of *Bellicoryne plumbaginifolia* was collected from Kannikars of Kanyakumari District, Tamil Nadu, India. The plant parts were identified taxonomically and authenticated according to various literatures, Flora of Madras Presidency and Wealth of India including other pertinent taxonomic literature.

### Phytochemical Analysis

The collected bulbs samples were washed thoroughly two times with running tap water and once with sterile water, air-dried, powdered using a pulverizer and used for extraction. About 50 grams of air-dried and coarsely powdered plant material was extracted successively with

250 ml of methanol using a Soxhlet extractor for at least 15 refluxs. After complete extraction, the extract in the round bottom flask were removed and condensed using rotary evaporator. After solvent evaporation, extracts were weighed for the percentage yield calculation. The thick syrup plant extract were labeled and stored at 5°C in sterile screw-capped vials for further use.

Preliminary phytochemical screening of methanol extract of *Bellicoryne plumbaginifolia* Bulbs was carried out to detect the phyto-constituents using standard conventional protocols (Trease and Evans, 1989; Sofowara, 1993 and Harbone, 1998). Alkaloids, carbohydrates, tannins and phenols, flavonoides, gums and mucilage, fixed oils and fats and saponins were qualitatively analyzed.

### Tumour cell lines

Cell lines of different tissue origin such as MCF-7 (human breast tumor), HeLa (human cervical cancer) and Dalton's lymphoma cells (mouse ascites tumor) were used. Cells were cultured in minimum essential medium (MEM) supplemented with 10% fetal calf serum, 2 mM L-glutamine, 100 IU/ml penicillin and 50 µg/ml streptomycin in a humidified atmosphere of 5% CO<sub>2</sub> in air at 37°C in a CO<sub>2</sub> incubator.

### MTT assay (Mossman et al., 1983)

Antiproliferative effects were measured in vitro by using MTT ([3-(4,5-dimethylthiazol-2-yl)-2,5-diphenyltetrazolium bromide] assays. After treatment, the living cells were assayed by the addition of 20 µl of 5 mg/ml MTT solution. Finally, the reduced MTT was assayed at 545 nm; wells with untreated cells were utilized as controls. Antiproliferative and cytotoxic effects were distinguished by cell number and the duration of treatment (72 h, 5000 cells/w, and 24 h, 25000 cells/w, respectively). Stock solutions of the tested materials were prepared with dimethyl sulfoxide (DMSO). The highest DMSO concentration (0.3%) of the medium did not have any significant effect on the cell proliferation. Extracts which demonstrated potent activity (growth inhibition > 50%) were selected for further in vitro testing (dose-response curve and cytotoxicity). To study the interactions between acridones and doxorubicin, a checkerboard method was applied. A series of 2-fold dilutions of the acridones was tested in combination with 2-fold dilutions of doxorubicin. The cell growth rate was determined with MTT staining; drug interactions were evaluated according to the following system (fractional inhibitory index = FIX):

FIX < 0.5	Synergism	1 < FIX < 2	Indifferent effect
FIX = 0.51-1	Additive effect	FIX > 2	Antagonism

## RESULTS AND DISCUSSION

Qualitative phytochemical analyses for alkaloids, carbohydrates, tannins, phenols, gums and mucilage, fixed oils and fats, saponins, proteins, volatile oils, flavonoids and steroids were screened in methanolic

extracts of *Bellicoryne plumbaginifolia* Bulbs. The screening of the extract indicated the presence of alkaloids, tannins and saponin in the methanolic extracts of bulb (Table 1). The plant extract yield percentage on the usage of methanol agreed with the earlier reported by

Jamuna et al. (2014) obtained in *Hypochaeris radicata* L. The plant extract obtained using soxhlet is varied among the herbal plants to plant. In a plant, different parts having differently yielded (Krishna et al., 2013). The plant extract yield percentage on the usage of methanol agreed with the earlier reported by Ramamurthy and Durgadevi (2017) obtained in *Brassica oleracea*.

**Table 1: Phytochemical screening of methanolic extract of *Bellicoryne plumbaginifolia***

Phyto-constituents	Observation
Alkaloids	+
Flavonoids	+
Terpenoids	+
Phenolic Compounds	+
Saponins	+
Tannins	+
Glycosides	+
Cardiac Glycosides	+
Coumarins	-

+: presence; -: absence

Anticancer activity of *Bellicoryne plumbaginifolia* was studied in different mammalian cell line. Anticancer activity of methanolic extract of *B. plumbaginifolia* as well as standard was determined through MTT cytotoxicity assay. In the preliminary study, the methanolic extract showed the good yielding capacity of phytocompounds activity. In this regards, the present investigation the methanolic extract of *B. plumbaginifolia* was studied in DL, HeLa and MCF-7

cell lines and its result labeled in the table 2 and also made with standard drug 5-Fluorouracil.

The minimum cell viability (21.4%) and maximum cell inhibition (85.6%) were noted in 100  $\mu$ g/ml concentration of *B. plumbaginifolia*. The IC<sub>50</sub> value (48.7 $\mu$ g/ml) was calculated for anticancer activity of methanolic extract of *B. plumbaginifolia* against DL, HeLa and MCF-7 cell lines. The 5-Fluorouracil used as a standard for this study. In the standard, the minimum cell viability (18.5%) and maximum cell inhibition (87.6%) were observed in higher concentration. The percentage of cell inhibition was noted in the different concentrations of methanolic extract of *B. plumbaginifolia* ranges from 10 to 100  $\mu$ g/ml. The lowest cell inhibition (32.1%) was recorded in the lowest concentration and highest cell inhibition (85.6%) was noted in the higher concentration of methanolic extract of *B. plumbaginifolia*.

Anticancer properties of many natural compounds isolated from different Indian plant extracts have been reported. Research is being carried out throughout the world to find a lead compound which can block the development of cancer in humans. Nature has always been a great contributor towards this goal. Plant-derived natural products such as flavonoids, terpenoids and steroids have received considerable attention due to their diverse pharmacological properties, which include cytotoxic and chemopreventive effects (Abdullaev, 2001). They were the first agents to advance into clinical use for the treatment of cancer (Cragg and Newman, 2005).

**Table 2. Survival analyses of cancer cells treated with extracts of *Bellicoryne plumbaginifolia*.**

Concentrations ( $\mu$ g ml <sup>-1</sup> )	DL cells		HeLa		MCF7	
	Cell viability (%)	Cell inhibition (%)	Cell viability (%)	Cell inhibition (%)	Cell viability (%)	Cell inhibition (%)
10	62.7	35.4	63.5	32.1	69.8	36.4
20	51.5	46.2	53.2	42.7	62.3	48.6
40	43.6	53.4	45.4	51.2	54.4	56.3
60	36.5	64.7	35.7	62.4	42.3	69.7
80	31.7	75.5	29.4	76.5	37.1	78.1
100	23.5	82.1	22.2	80.3	21.4	85.6
Vehicle control (DMSO)	100	0	100	0	100	0

In vitro cytotoxicity against four human cancer cell lines was determined by Monks et al. (1991) using 96-well tissue culture plates. One hundred microlitres of cell suspension was added to each well of the 96-well tissue culture plate. The cells were allowed to grow in a carbon dioxide incubator (37°, 5% CO<sub>2</sub>, 90% RH) for 24h. Test materials in complete growth medium (100  $\mu$ l) were added after 24 h of incubation to the wells containing cell suspension. The plates were further incubated for 48 h. The cell growth was stopped by gently layering trichloroacetic acid (50%, 50  $\mu$ l) on top of the medium in all the wells. The plates were incubated at 4°C for one hour to fix the cells attached to the bottom of the wells.

The liquid of all the wells was gently pipetted out and discarded. The plates were washed five times with distilled water to remove trichloroacetic acid, growth medium low molecular weight metabolites and serum proteins and then air-dried. The plates were stained with sulphorhodamine B dye (0.4 % in 1% acetic acid, 100  $\mu$ l) for 30 min. The plates were washed five times with 1% acetic acid and then air-dried (Skehan et al., 1990).

*Withania somnifera* as a potential source of new molecules that can curtail cancer growth were studied by Dredge et al. (2003). *Bellicoryne plumbaginifolia* bulbs have also been shown to inhibit the growth of human

cancer cell lines comparable to that produced by 5-Fluorouracil. The bulb extract produced antiproliferative activity on MCF-7 (human breast tumor), HeLa (human cervical cancer) and Dalton's lymphoma cells (mouse ascites tumor) tumor cell lines. The inhibitory concentrations obtained was  $25.1 \pm 0.91$  against colon cell line HCT-116 (Jayaprakasam et al., 2003), but in this study bulb extracts from different cancer cell treatments of *B. plumbaginifolia* cultivated in fly ash containing soil had shown more than 82% inhibition against human cell lines. Furthermore this study has reported growth inhibitory importance in *B. plumbaginifolia* against various cancer cell lines i.e. MCF-7, HeLa and Dalton's lymphoma cells tumor cell lines. Hence, this study has revealed remarkable anticancer potential in the pulbs of *Bellideryne plumbaginifolia*.

## REFERENCES

1. Abdullaev, FI. 2001. Plant derived agents against cancer. In: Gupta, S. K., editor. Pharmacology and therapeutics in the new millennium. Narosa Publishing House: New Delhi, India, p. 345-354.
2. Aquil, F., Ahmad, I., Mehmood, Z. 2006. Antioxidant and free radical scavenging properties of twelve traditionally used Indian medicinal plants. *Truk J Biol.* 30: 177-183.
3. Balachandran, P., Govindarajan, R. 2005. Cancer an ayurvedic perspective. *Pharmacol Res.* 51: 19-30.
4. Brisca Renuga F and S. Mary Mettilda Bai. 2013. Natural products used by the Kanikkars of Kanyakumari district, Tamil Nadu, India. *J. Pharmacogn. Phytochemis.* 2(1): 255-261.
5. Chah, K.F., Eze C.A., Emuelosi, C.E., Esimon, C.O. 2006. Antibacterial and wound healing properties of methanolic extracts of some Nigerian medicinal plants, *J. Ethnopharmacol.* 104: 164-167.
6. Cragg, G.M., Newman, D.J., Snader, K.M. 1997. Natural products in drug discovery and development. *J Nat Prod.* 60: 52-60.
7. Cragg, G.M., Newman, D.J. 2005. Plants as source of anticancer agents. *J Ethnopharmacol.* 100: 72-79.
8. Dredge, K., Dalglish, A.G., Marriott, J.B. 2003. Angiogenesis inhibitors in cancer therapy. *Curr Opin Investig Drugs*, 4: 667-674.
9. Gupta, S., Zhang, D., Yi, J., Shao, J. 2004. Anticancer activities of Oldelandia diffusa. *J Herb Pharmacother.* 4: 21-33.
10. Gurib- Fakim, 2006. Review- Medicinal plants: Traditions of yesterday and drugs of tomorrow, *Mol. Asp. Med.* 27: 1-93.
11. Harbone J. B. Phytochemical methods 1998. London: Chapman and Hall; pp. 117-119.
12. Jamuna, S., Subramaniam, P., Karthika, K. 2014. Phytochemical analysis and evaluation of leaf and root parts of the medicinal herb, *Hypochaeris radicata* L. for in vitro antioxidant activities *Asian Pac J Trop Biomed.* 4(1): S359-S367.
13. Jayaprakasam, B., Zhang, Y., Seeram, N.P., Nair, M.G. 2003. Growth inhibition of human tumor cell lines by withanolides from *Withania somnifera* leaves. *Life Sci.* 74: 125-32.
14. Krishna, T.M., Shiva, D., Keerthi, D., Nandini, A., Aswaq, A., Reddy, T. 2013. In vitro evaluation of antioxidant properties of *Cucumis melo* L. extracts of leaves and fruit. *Int J Pharm Bio Sci.*, 4(1): 705-712.
15. Modha, J., Modha, N. 2007. Role of Ayurveda in the Management of Cancer. International Centre for Ayurveda Studies, Gujarat Ayurveda University Jamnagar, Gujarat, India.
16. Monks, A., Scudiero, D., Skehan, P., Shoemaker, R., Paull, K., Vistica, D. 1991. Feasibility of a high-flux anticancer drug screen using a diverse panel of cultured human tumor cell lines. *J Natl Cancer Inst.* 83: 57-66.
17. Mossman, T. 1983. Rapid colorimetric assay for cellular growth and survival: application to proliferation and cytotoxicity assays. *J Immunol Methods*, 65: 55-63.
18. Parinitha, M., Srinivasa, B.H., Shivanna, M.B. 2005. Medicinal plant wealth of local communities in some villages in Shimoga distinct of Karnataka. *India J Ethnopharmacol.* 98: 307-312.
19. Ramamurthy, V., Durgadevi, K. 2017. Studies on ethanolic extract of *Brassica oleracea* L. var. Italica (Broccoli) and its effect on Dalda induced in Albino Rats. *Global Journal for Research Analysis*, 6(3): 593 - 596.
20. Skehan, P., Storeng, R., Scudiero, D., Monks, A., Mc Mohan, J., Vistica, D. 1990. New colorimetric cytotoxic assay for anticancer-drug screening. *J Natl Cancer Inst.* 82: 1107-12.
21. Sofowara, A. 1993. Medicinal plants and traditional medicine in Africa. Spectrum Books Ltd., Ibadan: Nigeria, 289-300.
22. Trease, G.E., Evans, W.C. 1989. *Pharmacognosy*. Bailliere Tindall, London, Edn. 11: 45 - 50.