



FREQUENCY OF HYPOMAGNESAEMIA IN PRE TERM LABOUR

Dr. Ayesha Jamal Khattak*¹, Dr. Sana Fatima² and Dr. Ambreen Azad³

Pakistan.

*Corresponding Author: Dr. Ayesha Jamal Khattak
Pakistan.
<https://doi.org/10.17605/OSF.IO/FM7TV>

Article Received on 21/05/2020

Article Revised on 11/06/2020

Article Accepted on 01/07/2020

ABSTRACT

Objective: To determine the frequency of hypomagnesaemia in women with preterm labor. **Study Design:** Cross sectional study **Settings:** This study was carried out in department of obstetrics & Gynaecology Nishtar Hospital Multan. **Duration of study:** This study was conducted from November 2018 to 10th May 2019. **Subject and methods:** A total of 219 pregnant women with singleton pregnancy, preterm labor and parity 1 – 4 were included in the study. Women with history of diabetes, hypertension smoking and polyhydramnios amniotic fluid index (AFI)>25 cm on ultrasound were excluded. Venous blood was drawn from entire subjects to evaluate the serum magnesium level and sent immediately to laboratory for analysis venous blood samples were obtained by venipuncture and collected in lithium heparin specimen bottles. Data was collected for hypomagnesaemia. **Results:** age range in this study was from 20 to 40 years with mean age of 30.356±3.60 years, mean parity 1.936±0.96, mean gravida 2.936 ± 0.96, mean gestational age 32.000±2.25 weeks and mean BMI was 27.337 ± 1.67 kg/m². Majority of the patients were from 20 – 30 years (51.1%). History of preterm delivery was 17.4%. Majority of patients belong to middle economic status (66.2%). Hypomagnesaemia was seen in 37.9% patients. **Conclusion:** it is concluded that low serum magnesium (hypomagnesaemia) is associated with preterm onset of labor. Maternal hypomagnesaemia may be used as a predictor of preterm labor:

KEYWORDS: Pregnancy, preterm labor, hypomagnesaemia.

INTRODUCTION

Preterm labor is the onset of labor before 37 completed weeks of gestation.^[1] One of the important causes of preterm delivery is the occurrence of spontaneous preterm labor.^[2,3] Although having varied etiology preterm labor may be due to changes in various biochemical functions starting at cellular level, signifying value of trace elements, especially magnesium. Magnesium plays a vital role in cellular function.^[4] Magnesium level in the serum range from 1.5 to 2.1 mg/dl.^[4] It is well established that serum magnesium levels fall throughout pregnancy. This falling level of magnesium plays a vital role in the onset of labor. This decreased level of magnesium causes reduced magnesium level in myometrium causing initiation of labor.^[5] The uterine relaxation effect caused by magnesium is because of the antagonism effect on calcium provoked uterine excitability.^[6] Hypomagnesaemia causes cervix to dilate facilitating passage of various microorganisms into cervix and finally uterus.^[7]

In a study conducted by Kehinde S showed that hypomagnesaemia was seen in 47% women presenting with preterm labour.^[8] Uptill now no study has been conducted in our population on role of magnesium in

preterm labour. This study is conducted to reduce morbidity and mortality resulting from prematurity associated with hypomagnesaemia.

MATERIAL AND METHODS

It was a cross sectional study conducted from November 2018 to May 2019 at obstetrics and Gynaecology department Nishtar Hospital Multan. Non-probability consecutive sampling technique was used. 219 women 20-40 years old having preterm labour and singleton pregnancy on ultrasound were included in the study after permission from ethical committee and research department. Women having history of diabetes, hypertension, cigarette smoking and polyhydramnios (amniotic fluid index AFI>25cm on ultrasound) were excluded from the study. Baseline demographic data of patients was collected. Informed consent was given by every patient venous blood sample was taken from each patient to determine serum magnesium level and was sent for analysis to laboratory immediately.

These samples were collected by a third year resident trainee in lithium heparin specimen bottles after obtaining by venipuncture. Data regarding hypomagnesaemia was collected and recorded on the proforma which was especially designed.

RESULTS

Age range in this study was from 20 to 40 years with mean age of 30.356 ± 3.60 years, mean parity 1.936 ± 0.96 , mean gravida 2.936 ± 0.96 , mean gestational age 32.000 ± 2.25 weeks and mean BMI was 27.337 ± 1.67 kg/m² as shown in table – I. Majority of the patients were from 20 – 30 years (51.1%) as shown in table – II. History of

preterm delivery was 17.4% as shown in table III. Hypomagnesaemia was seen in 37.9% patients as shown in table IV. Stratification of hypomagnesaemia with respect to age, parity and history of preterm delivery are shown in table V, VI, VII respectively.

Table-I: Mean \pm SD of age, parity, gravida, gestational age and BMI n=219.

| Demographics | | Mean \pm SD |
|--------------|-------------------------|-------------------|
| 1 | Age (years) | 30.356 \pm 3.60 |
| 2 | Parity | 1.936 \pm 0.96 |
| 3 | Gravida | 2.936 \pm 0.96 |
| 4 | Gestational Age (weeks) | 32.000 \pm 2.25 |
| 5 | BMI(Kg/m ²) | 27.337 \pm 1.67 |

Table- II: Frequency and percentage of patients according to age n=219.

| Age Groups (years) | No Patients of | %age |
|--------------------|----------------|-------|
| 20-30 | 112 | 51.1% |
| 31-40 | 107 | 48.9% |

Table- III: Frequency and percentage of patients according to history of preterm delivery n=219.

| Preterm Delivery | No Patients of | %age |
|------------------|----------------|-------|
| Yes | 38 | 17.4% |
| No | 181 | 82.6% |

Table- IV: Frequency and percentage of patients according to Hypomagnesaemia n=219.

| Hypomagnesaemia | No Patients of | %age |
|-----------------|----------------|-------|
| Yes | 83 | 37.9% |
| No | 136 | 62.1% |

Table- V: Stratification of Hypomagnesaemia with respect to age groups.

| Age Groups(years) | Hypomagnesaemia | | P value |
|-------------------|-----------------|------------|---------|
| | Yes | No | |
| 20-30 | 39(34.8%) | 73(65.2%) | 0.337 |
| 31-40 | 44(41.1%) | 63(58.9%) | |
| Total | 83(37.9%) | 136(62.1%) | |

Table- VI: Stratification of Hypomagnesaemia with respect to parity.

| Parity | Hypomagnesaemia | | P value |
|--------|-----------------|--------|---------|
| | Yes | Parity | |
| 1-2 | 60(37.5%) | 1-2 | 0.841 |
| 3-4 | 23(39%) | 3-4 | |
| Total | 83(37.9%) | Total | |

Table- VII: Stratification of Hypomagnesaemia with respect to history of preterm delivery.

| History of preterm delivery | Hypomagnesaemia | | P value |
|-----------------------------|-----------------|-------|---------|
| | Yes | | |
| Yes | 26(68.4%) | Yes | 0.000 |
| No | 57(31.5%) | No | |
| Total | 83(37.9%) | Total | |

DISCUSSION

Multiple studies have been conducted to determine relationship between preterm labour and low maternal serum magnesium levels. But no such study is carried out in Pakistan to establish this association between low serum magnesium levels and preterm labour. In our study, hypomagnesaemia was present in 37.9% patients.

Our study results match with results of study conducted by Kehinde S and her colleagues. In their study hypomagnesaemia was seen in 47% of patients presenting with preterm labour.^[9] In a study conducted at Bangladesh by Shahid AR and her colleagues frequency of hypomagnesaemia was 60% in 10 women presenting with preterm labour. Incidence of hypomagnesaemia

slightly less than 46% is seen in same number of patients in a study conducted by Shahid *et al.*^[10] This discrepancy may be because of difference in cut off points for magnesium (1.5 versus 1.9 mg/dl, respectively) in the two different studies. Main emphasis of our study is the role of level of serum magnesium in precipitating preterm labour. Previous reports and studies have documented a reduced level of serum magnesium in patients presenting with preterm labour.^[11,14] Study conducted by Pushpa and Jagdish, showed that in patients presenting with preterm labour serum magnesium level was low.^[15] A study conducted by Begum *et al* also documented that in women presenting with preterm labour there was a marked reduction in serum magnesium level (mean 1.77 ± 0.36). There are other studies and systemic reviews that failed to establish relationship between preterm labour and low serum magnesium levels.^[15,17] These studies suggested that use of magnesium whether oral or parenteral had no beneficial effect to delay preterm labour or prevent the occurrence of premature birth.^[15,18,20] and so it can't be routinely used as a tocolytic agent.^[20] But several studies revealed reduced serum magnesium levels in patients presenting with preterm labour showing that hypomagnesaemia is an important risk factor for causing preterm labour.

CONCLUSION

Hypomagnesaemia is a contributing factor for preterm labour. So, low maternal serum magnesium levels may be utilized for prediction of preterm labour.

REFERENCES

1. Sponge CY, Mercer BM, Dalton M, Kilpatrick S, Blackwell S, Sade G. Timing of indicated late-preterm and early-term birth. *Obstet Gynecol*, 2011 Aug; 118(2 Pt. 1): 323–33.
2. Anotayanonth S, Subhedar NV, Neilson JP, Harigopal S. Betamimetics for inhibiting preterm labour. *Cochrane Database Syst Rev*, 2004; 4: CD004352.
3. Vijay R, Prasad GS. Tocolytic with ritodrine: a comparative study in preterm labour. *Pak J Med Sci.*, 2006; 22: 64-69.
4. Sakaguchi Y, Fujii N, Shoji T. Hypomagnesemia as a significant predictor of cardiovascular and noncardiovascular mortality in patients undergoing hemodialysis. *Kidney Int.*, 2013; 85: 174–81.
5. Dario's ES, Seitz B, Watts SW. Smooth muscle pharmacology in the isolated virgin and pregnant rat uterus and cervix. *J Pharmacology Ext Ther*, 2012 Jun; 341(3): 587-96.
6. Swain R, Kaplan, Macules B. Magnesium for the next millennium. *South Med J.*, 1999; 92: 1040-7.
7. Hantoushzadeh S, Jafarabadi M, Khazardoust S. Serum magnesium levels, muscle cramps and preterm labour. *IntJ Gynaecol Obstet*, 2007; 98(2): 153-4.
8. Okunade KS, Oluwole AA, Adegbesan-Omilabu MA. A study on the association between low maternal serum magnesium level and preterm labour. *Adv Med*, 2014; 704875: 1-6.
9. Okunade KS, Oluwole AA, Adegbesan-Omilabu MA. A study on the association between low maternal serum magnesium level and preterm labour. *Adv Med*, 2014; 704875: 1-6.
10. Shahid AR, Hosni AU, Tahini HZ. Hypomagnesaemia in pregnancy: a predictor of preterm labour. *J Dhaka Med Coll*, 2010; 19(1): 51-57.
11. Kamal S, Shoran A, Kumar U, Shahid SK, "Serummagnesiumlevel in preterm labour," *Indian Journal of Pathology and Microbiology*, 2003; 46(2): 271–273.
12. R. B. Kurzal, "Serum magnesium level in pregnancy and in preterm labour," *The American Journal of Perinatology*, 1991; 8: 119–127.
13. D. Pushpo and W.M. A. Jagdish, "A study of serum magnesium level in preterm labour," *Journal of Obstetrics and Gynaecology of India*, 1991; 41: 269–273.
14. H. Begum, L. Shamsuddin, and S. Khatun, "Relationship of preterm labour with serum magnesium level," *Bangladesh Journal of Obstetrics & Gynaecology*, 2004; 19(1): 3–6.
15. R. Smolarczyk, J. Wójcicka-Jagodźńska, E. Romejko, P. Piekarski, K. Czajkowski, and J. Teliga, "Calcium-phosphorusmagnesiumhomeostasis in women with threatened pretermdelivery," *International Journal of Gynecology and Obstetrics*, 1997; 57(1): 43–48.
16. J. Wójcicka-Jagodźńska, E. Romejko, P. Piekarski, K. Czajkowski, R. Smolarczyk, and T. Lipiński, "Second trimestercalcium-phosphorus-magnesium homeostasis in women with threatened preterm delivery," *International Journal of Gynecology and Obstetrics*, 1998; 61(2): 121–125.
17. G. M. Arian, T. Panzitt, F. Gucer *et al.*, "Course of maternal serum magnesium levels in low-risk gestations and in preterm labor and delivery," *Fetal Diagnosis and Therapy*, 1999; 14(6): 332–336.
18. S. Han, C. A. Crothers, and V. Moore, "Magnesium maintenance therapy for preventing preterm birth after threatened preterm labour," *Cochrane Database of Systematic Reviews*, no. 7, Article ID CD000940, 2010.
19. Crothers CA, Hiller JE, Doyle LW. Magnesium sulphate for preventing preterm birth in threatened preterm labour. *Cochrane Database Sys Rev.*, 2002; 4: ID CD001060.
20. J. P. Elliot, "Magnesium sulfate as a tocolytic agent," *The American Journal of Obstetrics and Gynecology*, 1983; 147(3): 277–284.