



ANALYSIS OF EFFECT OF IRON DEFICIENCY ANEMIA ON HBA1C IN DIABETIC PATIENTS

Dr. Rohama Sardar*¹, Dr. Abdul Shakoor Shakir² and Dr. Baseerat Adnan³

Pakistan.

*Corresponding Author: Dr. Rohama Sardar
Pakistan.
<https://doi.org/10.17605/OSF.IO/TQMUB>

Article Received on 21/05/2020

Article Revised on 11/06/2020

Article Accepted on 01/07/2020

ABSTRACT

Introduction: Iron deficiency is one of the most prevalent forms of malnutrition. Globally, 50% of anemia is attributed to iron deficiency. **Aims and objectives:** The basic aim of the study is to analyze the effect of iron deficiency anemia on HbA1c in diabetic patients in Pakistan. **Material and methods:** This cross sectional study was conducted in Sheikh Zayed Hospital, Rahim Yar Khan during January 2019 to October 2019. This study include all diabetic patient who visited the OPD of the hospital. Socio-demographic characteristics were collected using structured questionnaires. An enrollment form containing past medical history pertaining to chronic diseases such as kidney disease, heart related problem, skin diseases, blood coagulation disorders and other medical complaints were completed for every individual. **Results:** In 120 cases of this study, mean HbA1c value was found to be $6.87 \pm 1.4\%$. The mean ferritin levels in male and female cases were 14.88 ± 8.62 and 8.92 ± 5.72 ng/ml, respectively. Also, mean FPG concentration was 99.66 ± 14.73 mg/dl in total cases. Hemoglobin levels in male and female cases were 9.54 ± 1.4 and 9.37 ± 1.33 g/dl, respectively. The mean MCV was 53.2 ± 8.16 fL and mean MCH was 17 ± 3.7 pg/cell for the all cases (both males and females). **Conclusion:** It is concluded that Iron deficiency anemia elevates HbA1c levels in diabetic individuals with controlled plasma glucose levels. The elevation is more in patients having plasma glucose levels between 100 to 126 mg/dl. Hence, before altering the treatment regimen for diabetes, iron deficiency anemia should be considered.

KEYWORDS: Iron deficiency, anemia, HbA1c, diabetes.

INTRODUCTION

Iron deficiency is one of the most prevalent forms of malnutrition. Globally, 50% of anemia is attributed to iron deficiency. Ferritin is the storage form of iron, and it reflects the iron status accurately. An earlier study showed that reduced iron stores have a link with increased glycosylation of hemoglobin A1C (HbA1c), leading to false high values of HbA1c in non-diabetic individuals.^[1] HbA1c is the most predominant fraction of HbA1, and it is formed by the glycosylation of terminal valine at the β -chain of hemoglobin. It reflects the patient's glycemic status over previous 3 months. HbA1c is widely used as a screening test for diabetes mellitus, and American Diabetes Association has recently endorsed HbA1c $\geq 6.5\%$ as a diagnostic criterion for diabetes mellitus. Its alteration in other conditions, such as hemolytic anemia, hemoglobinopathies, pregnancy, and vitamin B12 deficiency has been explained in a study conducted by Sinha *et al.* Although iron deficiency is the most common nutritional deficiency, reports of the clinical relevance of iron deficiency on HbA1c levels have been inconsistent.^[2]

HgbA to HbA1c conversion takes place during the entire life span of the red blood cell and the rate of this reaction is faster in diabetics because of the higher prevailing glucose concentration, resulting in a higher concentration of HbA1c.^[3] Red blood cells (RBC) are freely permeable to the plasma glucose molecules, and hemoglobin is practically exposed to the same glucose concentrations as plasma. Therefore, HbA1c level is directly proportional to average blood glucose concentration over the previous 4weeks to 3months or the average lifespan of the erythrocyte.^[4]

There are a number of methods available to estimate glycosylated hemoglobin like immunoturbidimetry, ion exchange high performance liquid chromatography (HPLC), boronate affinity, and enzymatic method. HbA1c level of $\geq 6.5\%$ is sufficiently sensitive and specific to identify individuals who are at risk for developing retinopathy and who should be diagnosed as diabetic.^[5]

MATERIAL AND METHODS

This cross sectional study was conducted in Sheikh Zayed Hospital Rahim Yar Khan 2019 to October 2019. This study was done with the approval of ethical committee of hospital. This study include all diabetic patient who visited the OPD of the hospital.

Data collection

Socio-demographic characteristics were collected using structured questionnaires. An Enrollment form containing past medical history pertaining to chronic diseases such as kidney disease, heart related problem, skin diseases, blood coagulation disorders and other medical complaints were completed for every individual. All patients were asked to provide a detailed history and were subjected to a physical examination. The levels of hemoglobin, MCH, hematocrit, MCV, MCHC, platelet count, total leucocyte count (TLC), and differential leucocyte count (DLC) were measured.

Biochemical analysis

Five milliliter of venous blood was collected by needle and syringe technique aseptically from each of the study

participants. CBC, serum ferritin and HbA1c tests were done following manufacturers procedure for running each test. CBC was done by using Cell dyn 1800 hematology analyzer, HbA1c by COBAS C 111 analyzer and Serum ferritin was determined.

Statistical analysis

All the data were collected and analysed using SPSS version of 20.0. All the values were expressed in mean and standard deviation.

RESULTS

In 120 cases of this study, mean HbA1c value was found to be $6.87 \pm 1.4\%$. The mean ferritin levels in male and female cases were 14.88 ± 8.62 and 8.92 ± 5.72 ng/ml, respectively. Also, mean FPG concentration was 99.66 ± 14.73 mg/dl in total cases. Hemoglobin levels in male and female cases were 9.54 ± 1.4 and 9.37 ± 1.33 g/dl, respectively. The mean MCV was 53.2 ± 8.16 fL and mean MCH was 17 ± 3.7 pg/cell for the all cases (both males and females).

Table 1: Baseline characteristics of the study subjects.

| | Female | | Male | | Total (n = 120) | |
|-------|-------------------|-------------------|-------------------|-----------------|-------------------|-------------------|
| | IDA | NA | IDA | NA | IDA | NA |
| Age | 54.65 ± 12.98 | 54.66 ± 12.02 | 56.6 ± 9.5 | 52.1 ± 8.08 | 55.46 ± 11.66 | 53.59 ± 10.59 |
| HbA1c | $7.02 \pm 1.58^*$ | 5.82 ± 0.53 | $6.67 \pm 1.06^*$ | 5.59 ± 0.85 | $6.87 \pm 1.4^*$ | 5.65 ± 0.69 |

HbA1c was significantly higher in females and in subjects above 50 years of age compared to males and subjects below 50 years of age.

Table 2: Mean HbA1c (%) genderwise in subjects stratified in two groups based on their age.

| | Female | | Male | | Total | |
|---------|-------------------|-----------------|-------------------|-----------------|-------------------|-----------------|
| | IDA | NA | IDA | NA | IDA | NA |
| Age <50 | 6.78 ± 1.88 | 5.59 ± 0.51 | $6.54 \pm 0.85^*$ | 5.48 ± 0.67 | $6.71 \pm 1.64^*$ | 5.53 ± 0.59 |
| Age >50 | $7.16 \pm 1.37^*$ | 5.91 ± 0.52 | $6.70 \pm 1.12^*$ | 5.69 ± 0.94 | $6.94 \pm 1.27^*$ | 5.77 ± 0.72 |

DISCUSSION

HbA1c has emerged as a marker of glycemic control, glycemic risk and predictor of diabetic complication and as screening tool for diagnosis of DM. Anemia may either increase or reduce the HbA1c values due to changes in the half-life of RBC.^[7] Numerous studies have been conducted on the effect of iron deficiency anemia on HbA1c in diabetic patient or non-diabetic patients and different results have been obtained but there is no clear explanation on mechanism how iron-deficiency affects HbA1c.^[8]

Iron deficiency anemia is one the most common anemia amongst the nutritional anemias in India. Initial studies conducted by Brooks *et al.*, Gram Hansen *et al.*, and Coban *et al.* showed effects of iron therapy on glycosylated hemoglobin and found a significant reduction in HbA1c levels after iron therapy in non-diabetic population.^[9] According to the explanation provided by Sluiter *et al.*, hemoglobin glycosylation is an irreversible process. Hence, HbA1 levels in erythrocyte

will be increased with cell age. In iron deficiency, red cell production decreases, consequently an increased average age of circulating red cells ultimately leads to elevated HbA1 levels.^[10] According to some workers, the changes in HbA1c levels were due to different laboratory methods used to analyze it. Goldstein *et al.* demonstrated that HbA1c measured by HPLC was increased two hours after a standard breakfast and incubating the red cell in 0.9% saline at 37°C for five hours eliminated this increment, which was explained by presence of labile HbA1c. This effect was eliminated by reagents used in newer enzymatic kits. Rai and Pattabiraman conducted a study to evaluate different methods used to analyze HbA1c and found no significant difference between them.^[11]

CONCLUSION

It is concluded that Iron deficiency anemia elevates HbA1c levels in diabetic individuals with controlled plasma glucose levels. The elevation is more in patients having plasma glucose levels between 100 to 126 mg/dl.

Hence, before altering the treatment regimen for diabetes, iron deficiency anemia should be considered.

REFERENCES

1. Rafat D, Rabbani TK, Ahmad J, Ansari MA. Influence of iron metabolism indices on HbA1c in non-diabetic pregnant women with and without iron-deficiency anemia: effect of iron supplementation. *Diabetes MetabSyndr*, 2012 Apr-Jun; 6(2): 102–5.
2. Trim O, Küçükerdoğan A, Günay U, Eralp O, Ercan I. Effects of iron deficiency anemia on hemoglobin A1C in type 1 diabetes mellitus. *Pediatr Int.*, 1999 Aug; 41(4): 357–62.
3. Mortensen HB, Christophersen C. Glucosylation of human haemoglobin A in red blood cells studied in vitro. Kinetics of the formation and dissociation of haemoglobin A1C. *ClinChimActa*, 1983 Nov; 134(3): 317–26.
4. Mortensen HB, Lund A, Christophersen C. Glucosylation of human haemoglobin A. Dynamic variation in HbA1c described by a bio kinetic model. *ClinChimActa*, 1984 Jan; 136(1): 75–81.
5. Rahbar S. An abnormal hemoglobin in red cells of diabetes. *ClinChimActa*, 1968 Oct; 22(2): 296–8.
6. Rahbar S, Blumenfeld O, Rainey HM. Studies of an unusual hemoglobin in patients with diabetes mellitus. *BiochemBiophys Res Commun*, 1969 Aug; 36(5): 838–43.
7. Trivially LA, Rainey HM, Lai HT. Hemoglobin components in patients with diabetes mellitus. *N Engl J Med.*, 1971 Feb; 284(7): 353–7.
8. Sluiter WJ, van Essen LH, Reitman WD, Doorknobs H. Glycosylated haemoglobin and iron deficiency. *Lancet*. 1980; 2:531–2.
9. Goldstein DE, Path SB, England JD, Hess RL, Da Costa J. Effects of acute changes in blood glucose on HbA1c. *Diabetes*, 1980 Aug; 29(8): 623–8.
10. Rai KB, Pattabiraman TN. Glycosylated haemoglobin levels in iron deficiency anemia. *Indian J Med Res*, 1986 Feb; 83: 234–6.
11. Hashimoto K, Noguchi S, Morimoto Y, Hamada S, Masada K, Imai S, et al. A1C but not serum glyated albumin is elevated in late pregnancy owing to iron deficiency. *Diabetes Care*, 2008 Oct; 31(10): 1945–8.