



AMIKACIN RESISTANCE PATTERN AND AEROBIC BACTERIAL PROFILE IN DIABETIC FOOT INFECTION PATIENTS

Mohammad Asaduzzaman^{*1}, Farha Matin Juliana², Sharmin Akhter¹, Jharna Akter¹, Nazmunnahar Ruma¹,
Nilufa Yesmin¹, Tasnuva Ferdous¹ and Sakib Hossen¹

¹Department of Biochemistry, Primeasia University, Dhaka, Bangladesh.

²Department of Biochemistry and Molecular Biology, Jahangirnagar University, Dhaka, Bangladesh.

*Corresponding Author: Mohammad Asaduzzaman

Department of Biochemistry, Primeasia University, Dhaka, Bangladesh.

Article Received on 21/07/2021

Article Revised on 11/08/2021

Article Accepted on 01/09/2021

ABSTRACT

Introduction: The goal of this study was to determine the frequency of bacterial isolates cultured from diabetic foot infections and to assess their amikacin resistance and susceptibility. **Methods:** A total of 412 diabetic foot lesions were included in this prospective analysis. The antibiotic susceptibility pattern of bacteria isolated from foot lesions was assessed using the Kirby-Bauer disk diffusion method. **Results:** The most commonly isolated Gram-positive bacteria were *Staphylococcus aureus*, followed by Enterococcus spp. and CoNS spp.. The most commonly isolated Gram-negative bacteria were *Klebsiella* spp., *Pseudomonas* spp., *Proteus* spp., *Escherichia coli*, *Enterobacter* spp., *Citrobacter* spp., *Citrobacter freundii*, and *Proteus vulgaris*. Amikacin was found to be 84.28% sensitive against the highest infection causing *Staphylococcus aureus* and followed by *Klebsiella* spp. (75.00%), *Pseudomonas* spp. (59.57%), *Proteus* spp. (92.85%), *Escherichia coli* (93.33%), *Enterobacter* spp. (100.00%) and *Enterococcus* spp. (91.30%). **Conclusion:** The present study confirmed the prevalence of amikacin drug resistant pathogens (17.96%) in diabetic foot ulcers. The diverse bacteria infecting the wound must be evaluated, as well as the antibiotic susceptibility patterns of the isolates from the infected lesion. This information is critical for selecting the right medications, eliminating resistance trends, and lowering healthcare costs.

KEYWORDS: Diabetic foot infection. Polymicrobial infections, Amikacin.

INTRODUCTION

Diabetes is a life-long condition. It affects many people all over the world. It is a significant public health issue.^[1] About one-fourth of people with diabetes will develop an ulcer throughout their lifetime, and as many as half of those ulcers turns into infected.^[2,3] In human beings with diabetes and foot ulcers, numerous elements, such as inappropriate antibiotic treatment, the chronic nature of the wound, and frequent clinic admission, can have an effect on the presence of multidrug-resistant microorganisms inside the ulcer.^[4] Moreover, the particular organisms identified in diabetic foot infections can differ not only from patient to patient and hospital to hospital but additionally from one a part of the country to some other.^[5]

Diabetic foot is a crippling condition. It offers long stretches of hospitalization. Health center prices may be very excessive and now and again not possible to undergo the charges, and the end result of an amputated limb. The ghostly limb adds insult to injury to the already battered psyche. No surprise, one of the most feared complications of diabetes is diabetic foot. The

classic triad of neuropathy, ischemia, and infection characterizes diabetic foot. The first priority should be to prevent diabetic foot. This can be accomplished by identifying high-risk individuals, such as those who have peripheral neuropathy, peripheral vascular disease, foot deformities, or callu.^[7]

If not treated promptly, infectious microorganisms are linked to amputation of the infected foot, as well as an increase in hospital stay, cost of management, morbidity, and mortality.^[7] Because most diabetic foot infections are true emergencies, antibiotic therapy should be initiated as soon as possible to improve the chances of saving the limb. Clinical presentation, gram-staining results, and knowledge of the organisms most commonly isolated from a specific infection should all be used to guide initial empirical therapy.^[8]

Diabetes-related foot ulcers are common and estimated to affect 15% of all diabetic individuals during their lifetime. Amputation is required in 15 to 20% of patients with such foot ulcers. Almost 85% of the amputations are preceded by diabetic foot ulcers.^[9,10,11] Peripheral

sensory neuropathy is the most important risk factor for the development of foot ulcers, followed by peripheral vascular disease. In diabetics, the proportion of neuropathic, neuroischemic, and purely ischemic lesions is 54, 34, and 10%, respectively.^[11] It is estimated that approximately 40,000 legs are amputated in India each year, with 75 percent of them being neuropathic with secondary infection, which is potentially preventable. Barefoot walking, illiteracy, low socioeconomic status, late presentation by patients, primary care physician ignorance about diabetic foot care, and belief in alternative systems of medicine all contribute to this high prevalence.^[12]

Infection in a diabetic foot is a limb threatening condition because the consequences of deep infection in a diabetic foot are more disastrous than elsewhere mainly because of certain anatomical peculiarities. The foot has several compartments, which are inter-communicating and the infection can spread from one into another, and lack of pain allows the patient to continue ambulation further facilitating the spread. The foot also has soft tissues, which cannot resist infection, like plantar aponeurosis, tendons, muscle sheaths, and fascia. A combination of neuropathy, ischemia, and hyperglycemia worsens the situation by reducing the defense mechanism.^[13]

The correct antibiotic selection based on the antibiograms of isolates from diabetic foot infections is critical for the proper management of these infections. As a result, the current study sought to assess the bacteriology of diabetic foot ulcers at Hospital Geral de Palmas in Tocantins, Brazil, in order to determine the relative frequencies of bacterial isolates cultured from foot infections and to assess the isolated bacteria's in vitro antibiotic resistance and susceptibility to an amikacin antibiotic.

MATERIALS AND METHODS

Materials

Study Design

This cross-sectional study was designed to assess the bacterial profile cultured from the wounds of diabetic foot patients, and to assess the functional pattern of the amikacin antibiotic on the cultured microorganisms.

Data Collection

Laboratory data from May 2020 to May 2021 were routinely collected from the microbiology department from Bangladesh Institute of health and sciences hospital (BIHS) Dhaka, Bangladesh. The total sample volumes were 412.

Methods

Swab sampling

Wound beds were prepared before specimen collection, where the wound immediate surface exudates and contaminants were cleansed off with moistened sterile gauze and sterile normal saline solution. Dressed wounds

were cleansed with non bacteriostatic sterile normal saline after removing the dressing. Aseptically the end of a sterile cotton-tipped applicator was rotated over 1 cm² area for 5 seconds with sufficient pressure to express fluid and bacteria to surface from within the wound tissue.^[14] The specimens were placed into sterile transport containers and sent to the microbiology laboratory for aerobic culturing within 20 minutes. Anaerobic culturing was not performed in this study.^[15,16]

Bacterial Culture

Cultures were processed following the standard procedure.^[15,16] Samples were inoculated on MacConkey agar (Oxoid), Chocolate agar (Oxoid) and Blood agar (Oxoid) media plate under class-II laminar airflow (NUVO SanajiMalzemelzeni, ImalatVcTicaret A.S, Turkey). The inoculum on the plate was streaked out for discrete colonies with a sterile wire loop sterilized by auto loop sterilizer (Germany) following standard procedures. The culture plates were incubated at 37°C by an incubator (Germany) for 48 hours and observed for the growth of bacteria through formation of colonies. All the bacteria were isolated and identified using morphological, microscopy (Japan) and biochemical tests like TSI (HiMedia), MIU (HiMedia) and Simmons Citrate (HiMedia) agar following standard procedures.

Antibiotic Susceptibility Assessment

The disc diffusion technique was used for antibacterial susceptibility testing of the isolates.^[17,18,19] using commercial antibiotics containing discs. We used the commercial antibiotic disc. Bacterial susceptibility was determined by the Kirby-Bauer disc diffusion method using antibiotic containing discs from Oxoid Ltd, UK. Amikacin antibiotic discs were used in this study. Interpretation of results were analysed using zone sizes. Zones of inhibition ≥ 21 mm will be considered sensitive, 16-20mm intermediate and < 15 mm resistant. Isolates were classified as either sensitive or resistant based on the definition of the Clinical and Laboratory Standard Institute.^[18]

Statistical Analysis

Data will be assessed using the free software GNU PSPP stable release 1.4.1/ September 5, 2020. It has a graphical user interface and conventional command-line interface. It is written in C and uses GNU Scientific Library for its mathematical routines. The name has "no official acronymic expansion".

RESULTS

A diabetic treatment center provided us with 412 participants for this research. They had all been diagnosed with diabetes and had diabetic feet. Males made up 247 (60.0%) of the subjects, with 165 females (40.0%) (Figure-1).




Figure-1: Sex distribution of the diabetic foot patients.

Among 412 diabetic foot subjects in the age group 10-20 years, males were 6 (1.5%) and females were 1 (0.2%). The age group > 20-30 years showed that males were 2

(0.5%) and females were 13 (3.2%). 27 (6.6%) males and 17 (4.1%) females were found in the age group > 30-40 years. The age group > 40-50 years showed that almost equal males and females were in this area. Males were 55 (13.3%) and females were 57 (13.8%). The highest female subjects were found in the > 40-50 year age group. The highest peak for diabetic foot subjects went to males, 84 (20.4%) in the age group category > 50-60 years and in this category females were 42 (10.2%). 48 males (11.7%) and 32 females (7.8%) were found in the age group > 60-70 years. The age group > 70-80 years was represented by 21 males (5.1%) and 3 females (0.7%). age groups > 80-90 years and > 90-100 years showed no female subjects. 1 male (0.2%) was found in the age group > 80-90 years and 2 males (0.5%) were found in the age group > 90-100 years (Figure-2).




Figure-2: Sex distribution in different age groups.

Table-1: Distribution of microorganisms among diabetic foot patients.

Microorganism	Number	Percent	Valid Percent	Cumulative Percent
<i>Acinetobacter</i> spp.	12	2.91	2.91	2.91
<i>Citrobacter freundii</i>	4	0.97	0.97	3.88
<i>Citrobacter</i> spp.	10	2.43	2.43	6.31
<i>CoNS</i> spp.	12	2.91	2.91	9.22
<i>Enterobacter</i> spp.	27	6.55	6.55	15.78
<i>Enterococcus</i> spp.	23	5.58	5.58	21.36
<i>Escherichia coli</i>	30	7.28	7.28	28.64
<i>Klebsiella</i> spp.	52	12.62	12.62	41.26
<i>Proteus</i> spp.	42	10.19	10.19	51.46
<i>Proteus vulgaris</i>	3	0.73	0.73	52.18
<i>Pseudomonas aeruginosa</i>	8	1.94	1.94	54.13
<i>Pseudomonas</i> spp.	47	11.41	11.41	65.53
<i>Serratia marcescens</i>	1	0.24	0.24	65.78
<i>Staphylococcus aureus</i>	141	34.22	34.22	100.00
Total	412	100.00	100.00	

Among 412 diabetic foot patients, 2.91% were infected with *Acinetobacter* spp., 0.97% were with *Citrobacter*

freundii. 2.43% were infected with *Citrobacter* spp., *CoNS* spp. were in 2.91% of patients. *Enterobacter* spp.

was responsible for 6.55% of infections. *Escherichia coli* caused 7.28% of infections. 12.62% infection identified for *Klebsiella* spp. and 10.19% for *Proteus* spp.. *Proteus vulgaris* infected a small percentage of the patients and that was 0.73%. *Pseudomonas aeruginosa* was dedicated

to 1.94% infection. Among all microorganisms, the lowest infection was found by *Serratia marcescens* and the infection rate was 0.24%. The highest number of infections were caused by *Staphylococcus aureus*. Here we found a 34.22% infection rate (Table-1).




Figure-3: Distribution of gram positive (+ ve) and gram negative (-ve) microorganisms among diabetic foot patients.

Among the all microorganisms,^[14] highest of organisms were gram negative,^[10] and lowest numbers were gram positive.^[4] *Acinetobacter* spp.^[12] *Citrobacter freundii*,^[4] *Citrobacter* spp.^[10] *Enterobacter* spp.^[27] *Escherichia coli*,^[30] *Klebsiella* spp.^[52] *Proteus* spp.^[42] *Proteus*

vulgaris.^[3] *Pseudomonas aeruginosa*.^[8] and *Pseudomonas* spp.^[47] were gram negative bacteria. *Klebsiella* spp., *Pseudomonas* spp., *Proteus* spp., *Escherichia coli*, *Enterobacter* spp., *Citrobacter* spp., *Citrobacter freundii*, *Proteus vulgaris*




Figure-4: Sex distribution of gram positive (+ ve) and gram negative (-ve) microorganisms among diabetic foot patients.

Figure-4 described sex distribution regarding specific microorganism. i) In respect to *Acinetobacter* spp. pattern was 3.24% for male and 2.42% for female, ii) 0.01% male was found for *Citrobacter freundii* iii) 2.02% of males and 3.03% of females were found infected with

Citrobacter spp. iv) 2.43% male was found for CoNS spp. and female was 3.64%. v) 7.29% male was found for *Enterobacter* spp. and female was found 5.45%. vi) 6.07% of males were infected with *Enterococcus* spp. and 9.09% female. vii) *Escherichia coli* found in 7.29%

male, female percentage was 7.27. viii) Male found 11.74% and 13.94% female in respect to *Klebsiella* spp. ix) *Proteus* spp. was found in 10.93% male and in 9.09% female. x) *Proteus vulgaris* was found in 0.81% of males and in 0.61% of females. xi) 1.62% of males were found with *Pseudomonas aureginosa* infection. 2.42% females

were found with *Pseudomonas aureginosa* infection. xii. No female was found with *Serratia marcescens* and 100.00% were males. xiii. *Staphylococcus aureus* was found as a big name in our experiment. 34.41% of males were infected by this organism and the female percentage was 33.94.




Figure-5: Amkacin's functional profile.

As a whole, we got a scenario that amikacin was resistant for 17.96% of diabetic foot patients and 82.04% of patients found okay (Figure-5).




Figure-6: Resistant and sensitivity pattern of amikacin against specific microorganism.

Figure-6 described resistant and sensitivity pattern against amikacin. i) In respect to *Acinetobacter* spp. pattern was 50-50, ii) 100.00% sensitivity was found for *Citrobacter freundii* iii) 80.00% sensitive and 20.00% resistant applied for *Citrobacter* spp. iv) 100.00% sensitivity was found for CoNS spp. v) 100.00% sensitivity was found for *Enterobacter* spp. vi) 8.70% resistant *Enterococcus* spp. was found rest 91.30% was sensitive. vii) *Escherichia coli* showed 6.67% resistance against amikacin, sensitivity percentage looked high (39.33%). viii) amikacin found 25% resistant and 75%

sensitive against *Klebsiella* spp. ix) *Proteus* spp. was found 7.14% resistant and 92.86% sensitive. x) Three persons were with found with *Proteus vulgaris* infection. 33.33% diabetic foot infection with this organism showed resistance against amikacin and 66.67% were fine with amikacin. xi) Eight persons were with found with *Pseudomonas aureginosa* infection. 37.50% diabetic foot infection with this organism showed resistance against amikacin and 63.50% were fine with amikacin. xii) Only one person was found with *Serratia marcescens* and showed 100.00% sensitivity. xiii)

Staphylococcus aureus was found as a big name in our experiment. 15.71% of this organism showed resistance against target antibiotic and 84.29% showed sensitive.

DISCUSSION

Foot infections in patients with diabetes are a common, complex, and costly problem.^[20] In the present study, we found that elderly patients (>60 years of age) constituted the majority of patients with foot infections. This may be explained by the fact that foot lesions occur commonly among patients with diabetes, particularly the elderly and those with sensory neuropathy.^[21] Previous studies have shown that the susceptibility to foot infections is greater in male patients than in female patients.^[22,23] However, in our study, we did not find differences between male and female patients, which may be because of the limited number of patients.

Diabetic foot ulcers are colonized by pathogenic bacteria that may predispose a susceptible patient to a lower extremity infection, defined as the invasion and multiplication of microorganisms in body tissues associated with tissue destruction or host inflammatory responses.^[24] In the present study, we found that the majority of lesions were located on the right toe and plantar region, and varied in duration from 1 day to more than 90 days. Additionally, recent lesions (1-30 days) were the most common. Our findings are in accordance with the results of Donoso *et al.* (2013).^[25]

This study is limited by the fact that cultures for anaerobic bacteria could not be performed. In diabetic foot infections, the role of anaerobic bacteria is particularly unclear; some studies have reported that anaerobic bacteria play a minor role.^[26,27] while other studies found a high incidence of anaerobic bacteria.^[23,28]

The most common pathogens isolated were Gram-positive cocci, such as *Staphylococcus aureus* and *Staphylococcus saprophyticus* (CONS) and Gram-negative rods, such as *Proteus* spp. and *Enterobacter* spp. Although the findings of our study are consistent with the results of previous studies showing that Gram-positive bacteria were predominant in diabetic foot infections.^[28] other studies have reported that Gram-negative bacteria were predominant in particular regions.^[29,30] These results suggest, in part, differences in the type and severity of infections.^[23,31] Aerobic Gram-negative bacteria (mainly Enterobacteriaceae and sometimes *Pseudomonas aeruginosa* or other Gram-negative species) are usually isolated in conjunction with Gram-positive cocci in patients with chronic or previously treated infections.^[29] which is consistent with our findings. Polymicrobial infections accounted for 70% of all infections. Although polymicrobial etiology has been implicated in diabetic foot infections.^[30] a previous study reported the predominance of monomicrobial infections.^[30] These discrepancies suggest differences in diabetic foot infections, with severe infections usually

having polymicrobial isolates and mild infections usually having monomicrobial isolates.^[31,34]

CONCLUSIONS

The present study report has some limitations because cultures for anaerobic bacteria could not be performed and sample size was small. However, it confirmed the high prevalence of multidrug-resistant pathogens in diabetic foot ulcers. Diabetic foot infections were predominantly due to Gram-positive bacteria, such as *Staphylococcus aureus*, or were polymicrobial infections. Many studies on the bacteriology of diabetic foot infections have reported results that vary and are often contradictory.^[28,29,31] In such cases, application of molecular techniques may lead to more accurate microbial characterizations and targeted antibiotic therapy. Therefore, it is necessary to evaluate the different microorganisms infecting the wound on a routine basis and to know the antibiotic susceptibility patterns of the isolates from the infected wound in patients with diabetic foot lesions. This knowledge is crucial for planning the treatment of these patients with the appropriate antibiotics, reducing resistance patterns, and minimizing healthcare costs. We hope the data presented on this article can assist the clinicians in determining the multidrug-resistant pathogens in diabetic foot ulcers. In this study amikacin was found resistant for 17.96% of diabetic foot patients and 82.04% of patients were found susceptible.

REFERENCES

1. Sivanmaliappan TS, Sevanan M. Antimicrobial susceptibility patterns of *Pseudomonas aeruginosa* from diabetes patients with foot ulcers. *Int J Microbiol*, 2011; 2011: 605195. (Accessed 2014 January 14). Available at <http://www.hindawi.com/journals/ijmicro/2011/605195/>.
2. Lipsky BA, Berendt AR, Deery HG, Embil JM, Joseph WS, Karchmer AW, et al. Diagnosis and treatment of diabetic foot infections. *Clin Infect Dis.*, 2004; 39: 885-910.
3. Lavery LA, Armstrong DG, Murdoch DP, Peters EJ, Lipsky BA. Validation of the Infectious Diseases Society of America's diabetic foot infection classification system. *Clin Infect Dis.*, 2007; 44: 562-565.
4. Kandemir O, AkbaySahin E, Millan A, Gen R. Risk factor for infection of the diabetic foot with multi-antibiotic resistant microorganisms. *J Infect*, 2007; 54: 439-445.
5. El-Tahawy AT. Bacteriology of diabetic foot infections. *Saudi Medical J*, 2000; 21: 344-347.
6. Pendsey SP. Understanding diabetic foot. *International journal of diabetes in developing countries*, 2010 Apr; 30(2): 75.
7. Hartemann-Heurtier A, Robert J, Jacqueminet S, Ha VG, Golmard JL, Jarlier V, et al. Diabetic foot ulcer and multidrug-resistant organisms: risk factors and impact. *Diabet Med*, 2004; 21: 710-715.

