



**SOME NEW HOST RECORDS FOR FOLIICOLOUS FUNGI FROM DISTRICT
BALRAMPUR U. P. INDIA**

Rajiv Ranjan* and Ajay Kumar

Postgraduate Department of Botany M.L.K (P.G.) College, Balrampur- 271 201 (U.P.) India.

***Corresponding Author: Rajiv Ranjan**

Postgraduate Department of Botany M.L.K (P.G.) College, Balrampur- 271 201 (U.P.) India.

Article Received on 07/02/2022

Article Revised on 28/02/2022

Article Accepted on 20/03/2022

ABSTRACT

Balrampur is a city and a municipal board in Balrampur district in the state of Uttar Pradesh, India. The authors surveyed diversified habitat of Balrampur for foliicolous fungi and collected 16 new hosts for Foliicolous fungi from Balrampur during July 2021 to January, 2022.

KEYWORDS: New Hosts, Foliicolous fungi, Balrampur, U.P.

INTRODUCTION

The leaves provide a very suitable habitat for the growth & development of fungal pathogen by providing ample surface area and nutrient supply. Such leaf inhabiting fungi are known as foliicolous and the invaded area of the leaf appears as leaf spot or leaf lesion. The weed and forest plants serve as reservoir of leaf spot pathogen which on getting opportunity may spread to agriculture & horticulture plants.

Balrampur is situated on the bank of river Rapti and is the district headquarters of Balrampur district. The creation of Balrampur District was done by G.D.No. 1428/1-5/97/172/85-R-5 Lucknow dated May 25, 1997 by the division of District Gonda. Siddharth Nagar, Shrawasti, Gonda District, are situated in the east-west and south sides respectively and Nepal State are situated in its northern side. One of the most popular Hindu worship place is situated in Tulsipur about 27 kilometers from the district headquarters. It is known as Devi Patan. The temple has the distinction of having included in 51 "Shaktipeethas" of Goddess Durga according to Hindu mythology. Balrampur city is in close vicinity of Shrawasti where Lord Gautam Buddha is considered to have displayed his supernatural powers in the spiritual transformation of Angulimala, a famous dacoit who wore a necklace (mala) of fingers (anguli). The area of the district is 336917 Hec. In which the agriculture irrigated area is 221432 Hec. In the north of the district is situated the Shivalics ranges of the Himalayas which is called Tarai Region. Keeping it in view, the authors surveyed most diversified habitat of Balrampur during July 2021 to January, 2022.

MATERIALS AND METHODS

The climatic condition favours the growth of various types of phanerogamic vegetation along with seasonal and annual crops and other plants. With a view to study the foliicolous fungi in their natural habitat, frequent collection trips will be arranged. The following articles would be required for collecting foliicolous fungi- collection containers, hand lens, pruning scissor or secateurs, light plant pressures, blotting paper, paper envelop, field note book etc.

Laboratory processing and preliminary examination.

Preparations

- (a) Photograph of both host and pathogen will be taken.
- (b) Scrap mount: - If the organisms are superficially attached with the host tissue scrap mounts are made by a sharp razor or scalpel.
- (c) Collodion Preparation: - A drop of collodion solution is applied to a colony on the leaf. The fungus gets embedded entirely and the dried film is peeled off readily from the host surface. Removal of collodion by acetone on a glass slide gives undisturbed preparation.
- (d) Squash preparation: - The fruiting body is mounted, cleared and examined. Then the preparation is tapped vigorously and reheated. In this way the fruiting body is broken and content is released.
- (e) Hand cut Section preparation: A hand cut section of infected tissue is made with sharp razor to study immersed or semi-immersed fungi. Section cutting for host parasite interaction / relation.

Staining and Mounting

For routine microscopic study in the lab temporary slides are made in different type of stains and mountants according to nature of fungal forms involved.

- (a) **Lacto- phenol cotton blue:** - The lacto-phenol mounting fluid is used for mounting-colored fungi. For locating cytoplasm, septa, guttules other structures and hyaline forms 0.05-0.01% cotton blue is added.
- (b) **Poly- vinyl Alcohol:** - Benson, 1969 is used in routine staining and mounting.
- (c) **Lacto-fuchsin:** - By this cell walls are stained more clearly, rapidly and with more suitable color specially for photography [Carmichael., 1955]. Slides prepared in mountants are sealed with wax or commercial good quality nail polish and are stored for further study.

Camera Lucida: - Drawings will be made of the distinctly different taxa of generic or species rank so as to show the morpho taxonomic features of vital importance.

The new host and their foliicolous fungal species are given below.

S.N.	Name of the Host & family	Name of the Foliicolousfungus
1.	<i>Litchi chinensis</i> Sonn. Litchi (Sapindaceae)	<i>Alternaria alternata</i> (Fr) Keissler
2.	<i>Hibiscus mutabilis</i> Linn. Gurhal (Malvaceae)	<i>Alternaria dianthi</i> Stev& Hall
3.	<i>Caladium bicolor</i> (Aiton)Vent. (Araceae)	<i>Alternaria sp.</i> Neergaard
4.	<i>Dracaena draco</i> (L.) L. (Asparagaceae)	<i>Alternaria dianthicola</i> Neergaard
5.	<i>Crinum latifolium</i> Linn. Sudarshan (Amaryllidaceae)	<i>Aspergillus sp.</i> Mich.ex Fr.
6.	<i>Abelmoschus esculentus</i> (L.) Moench Ladyfinger, Bhindi (Malvaceae)	<i>Cercosporaabelmoschi</i> Ell. &Ev.
7.	<i>Alstoniascholaris</i> R.Br. Black board tree, <i>Saptaparni</i> , Milk Wood Pine (Apocynaceae)	<i>Cercosporaalstoniae</i> Amy Rosemann
8.	<i>Cucurbita maxima</i> Duchesne Kaddu, Pumpkin (Cucurbitaceae)	<i>Cercosporacirtullina</i> (Cooke)
9.	<i>Amorphophallus companulatus</i> DecneSooran, Jimikand (Araceae)	<i>Cercosporachevalleri</i> Sacc.
10.	<i>Ficus religiosa</i> L. Peepal (Moraceae)	<i>Cercosporafici-religiosae</i> Chiddarwar
11.	<i>Cymbopogonjwarancusa</i> (Jones Schutt) Jwarancus (Poaceae)	<i>Cercosporascipicola</i> (Fuckel) Van Zinderen Bakker.
12.	<i>Livistona chinensis</i> R.Br. (Arecaceae)	<i>Curvulariafallax</i> B. Oedijn
13.	<i>Colocasia esculenta</i> L. (Schott) Arvi (Araceae)	<i>Cladosporium colocasiae</i> Sawada
14.	<i>Sida acuta</i> Burm. F. (Malvaceae)	<i>Cercospora sp.</i> Fres.
15.	<i>Mangifera indica</i> L. Aam (Anacardiaceae)	<i>Meliolamangiferae</i> Earle
16.	<i>Jasminum sambac</i> (L) Aition. Bella (Oleaceae)	<i>Meliola sp.</i> Fr.

The literature Ellis, 1971, 1976; Deighton, 1976, 1979; Braun, 1987, 1988 a, b, 1991; Braun *et al.*, 1992; Bagyanarayan and Braun, 1999; Bagyanarayan *et al.*, 1995; Vasudeva, 1963; Chupp, 1954; Bilgramiet *al.*, 1979, 1981, 1991; Jamaluddin *et al.*, 2004; Singh *et al.*, 2009; Mall, 2011a,b, Srivastava 2011; Parmar *et al.*,

Morpho taxonomic treatment. - Hitherto undescribed forms of foliar fungi will be executed with the help of present literature and expertise available at hand.

- New taxon will be described in English or Latin or both as and when required.
- Material (holotypes) will be deposited in recognized Herbaria for accession no.
- The Mycobank number of each new Taxa will be procured.

RESULTS AND DISCUSSION

The authors surveyed during July 2021 to January, 2022 in diversified habitats of Balrampur for the collection, study and documentation of the leaf spot micro fungi infecting variety of the angiosperems has resulted in abundant gathering of the fungal specimens. The holotype of collections for allotment of accession number from HCIO is in process. The authors collected 16 new hosts for foliicolous fungi from Balrampur.

2012; Behraet *al.*, 2012; Kumar and Mall, 2012, 2013, 2015; Mall, 2015 a, b, Rani *et al.*, 2015; Singh and Mall, 2015; Mall and Tripathi, 2016; Tripathi *et al.*, 2016; Singh *et al.*, 2020 reveals that the fungal taxa mentioned above are hitherto unexplored from Balrampur. Hence

are the new host records for Indian mycoflora from Balrampur.

SCOPE OF STUDY

The Foliicolous Fungi causes huge losses every year in different parts of the world. The fungal pathogens producing leaf spots infect a large variety of hosts including most of the crops, forests and other plants. The destruction caused by these enemies of leaves is a serious problem before us. The focus of this research is identification & documentation of foliicolous fungi which will assist in the discovery of new fungicides and ideas to overcome from the severity of these enemies of nature as well as in the protection of floral diversity from the infection of these pathogens and also in the conservation of valuable flora of the area.

ACKNOWLEDGEMENTS

The author is thankful to Professor J. P. Pandey Principal, M.L.K. (P.G.) College, Balrampur and Head of Department Botany for providing facilities.

REFERENCES

1. Bagyanarayana, G., Braun U. & Jagdeeswar P. (1995). Notes on Indian *Cercospora* and allied genera (I) Gyptog Bot, 5: 363-366.
2. Bagyanarayan, G. & Braun U. (1999). Phytopathogenic micromycetes from India. *Sydowia*, 51(I): 1-19.
3. Behera, S.K., A.K. Mishra, N. Sahu, A. Kumar, N. Singh, A. Kumar, O. Bajpai, L.B. Chaudhary, P.B. Khare & R. Tuli (2012). The study of microclimate in response to different plant community association in tropical moist deciduous forest from northern India. *Biodiversity and Conservation*, 21(5): 1159-1176.
4. Bilgrami, K.S., Jamalddin and Rizwi, M.A. (1979). *Fungi of India*, Part-I. Today and Tomorrow's Printers and Publishers. New Delhi, 467.
5. Bilgrami, K.S., Jamalddin and Rizwi, M.A. (1981). *Fungi of India*, Part-II. Today and Tomorrow's Printers and Publishers. New Delhi, 140.
6. Bilgrami, K.S., Jamalddin and Rizwi, M.A. (1979). *Fungi of India*. List and References. Today and Tomorrow's Printers and Publishers. New Delhi, 778.
7. Braun, U. (1987). A monograph of the Erysiphe's (Powdery mildews). *Beihefte zur Nova Hedwigia*, 89: 1-700.
8. Braun, U. (1988 a). Studies on *Ramularia* and allied genera (I). *Int. J. Mycol Lichenal*, 1: 3 (2/3).
9. Braun, U. (1988 b), Studies on *Ramularia* & allied genera (II). *Nova Hedwigia*, 50: 499-521.
10. Braun, U. (1991). Taxonomic problems of the *Ramularia/Cercospora* complex. *Studies in mycology*.
11. Braun, U. Bagyanarayana G. & Jagdeeswar P. (1992). Notes on Indian *Cercospora* and allied genera (II) *Int. J. Mycol. Lichenal*, 14(3): 361-374.
12. Carmichael J.W. (1955). Lacto- fuschin, A new medium for mounting fungi. *Mycologia*, 47: 611.
13. Chupp, C. (1954). A monograph of the fungus genus *Cercospora*. Ithaca, New York.
14. Deighton, F.C. (1967). Studies on *Cercospora* and allied genera-II, *Passalora, Cercosporidium* and some species of *Fusicladim* on *Euphorbia*. *Mycol. pap*, 112: 80.
15. Deighton, F.C. (1976). Studies on *Cercospora* and allied genera- VI, *Pseudocercospora* Spwg., *Pantospora* Cif. And *Cercoseptoria* Petr. *Mycol. pap.*, 140: 156 pp.
16. Deighton, F.C. (1979). Studies on *Cercospora* and allied genera- VII, New species and redispotion's. *Mycol. Pap*, 144: 1-56.
17. Ellis, M.B. (1971). *Dematiaceous Hyphomycetes*. CMI, Kew, U.K., 608.
18. Ellis, M.B. (1976). *More Dematiaceous Hyphomycetes*. CMI, Kew, U.K., 608.
19. Jamaluddin, Goswami, M.G. and Ojha, B.M (2004). *Fungi of India, 1989 – 2001*. Scientific Publishers (India), Jodhpur, 326.
20. Kamal, Verma, R.P. & Morgan-Jones, G. (1986). Notes on Hyphomycetes-LI *Kameshwaromyces*. A new foliicolous, sooty mould- like genus from Madhya Pradesh, India. *Mycotaxon*, 25: 247-250.
21. Kumar, Ajay and Mall, T. P. (2012). Foliicolous fungi from Bahraich U.P. (India). *Indian Journal of Bioinformatics and Biotechnology*, 1(3): 32-34.
22. Kumar, Ajay and Mall, T. P. (2013). *Cercospora alstoniae* sp. nov. on *Alstoniaschlolaris* from Bahraich (U.P.) India. *Indian Journal of Bioinformatics and Biotechnology*, 2(1): 1-3. Myco bank no. MB 805222.
23. Kumar, Ajay and Mall, T.P. (2015). Status of Susceptible Hosts for Foliicolous Fungi in North Central Tarai Forests of (U.P.) India. *Int. J. Curr. Res. Bioscl. Plant Biol*, 2(11): 27-30.
24. Mall, T.P. (2011 a). Foliicolous Fungi: Earth Living Treasure in North Central Tarai Forest of Uttar Pradesh (India). *Ind. Jour. Pl. Heath*, 3(1): 8-20.
25. Mall, T.P. (2011 b). *Cercospora oudhensis* Mall sp. nov. on threatened plant *Indopiptaenia oudhensis* from Shrawasti, U.P. India. *Trends in Biosciences*, 4(1): 132-133.
26. Mall, T. P. and Ajay Kumar (2015). Effect of Climate Change on Status of Foliicolous Fungi from Bahraich U.P. India. *Indian Journal of Pathology: Research and Practice*, 4(2): 109-112.
27. Mall, T.P. (2015). Diversity of Foliicolous Fungi from North Central Tarai Forests of (U.P.) India. *Agricultural Science Research Journal*, 5(12): 195-205.
28. Mall, T.P. (2016). *Diospyros cordifolia* Roxb. -An under exploited Potent Ethnomedicinal Feed- A Review. *World Journal of Pharmaceutical Research*, 5(8): 172-177.
29. Parmar, Arpita, Mall, T. P. and Singh, R. B. (2012). Population dynamics of mycorrhizal fungi in

- rhizosphere of pigeon pea (*Cajanus cajan*) (L.) Millsp. *Current Botany*, 3(5): 64-68.
30. Rani, S., Singh, R., Gupta, S., Dubey, S., and Rajdan, V.K. (2015). Identification of resistant sources and epidemiology of early blight (*Alternaria solani*) of tomato (*Lycopersicon esculentum*) in Jammu and Kashmir. *Indian Phytopath*, 68(1): 87-92.
 31. Singh, D. P. and Mall, T. P. (2009). Some additional records of foliicolous fungi from North Central Tarai Forests of U.P. *Environment Conservation Journal*, 10(1&2): 27-29.
 32. Singh, D.P. and Mall, T.P. (2015). Foliar Fungi from North Western Tarai Forests of Uttar Pradesh India. ASR Publications, 102, Pooja Colony-II PaviSadakpur, Loni Ghaziabad, 464-472.
 33. Singh, R., Verma, S.K., Yadav, S., Bhojak, P., Kumar, S. (2020). Morphology and phylogeny of *Pseudocercosporahamiltoniani*- A new species. comparable to *Sirosporium* from Uttarakhand, India. *Phytotaxa*, 458(4): 281–293.
 34. Srivastava, S.K. (2004). Floristic diversity in Uttar Pradesh- an overview. *Journal of Economic and Taxonomic Botany*, 28(2): 292–334.
 35. Srivastava, S.K. (2011). Plant diversity and conservation strategies of Uttar Pradesh. *Phytotaxonomy*, 11: 45–62.
 36. Tripathi, S.C., Kumar, Ajai and Mall, T.P. (2016). *Cercosporabahraichensis* sp. nov. on *Sorghum vulgare* -a great nutrimental potent ethnomedicinal plant. *Agricultural Science Research Journal*, 6(7): 162-165.
 37. Vasudeva, R.S. (1963). *Indian Cercosporae*, ICAR, New Delhi.