



A STUDY OF SERUM TESTOSTERONE LEVELS IN DIABETES MELLITUS TYPE 2 MALE PATIENTS

Ammara Quraishi¹, Vatsal Vyas², Nishita Karulkar³, Hariom Vaja⁴, Ajaykumar Raiyani⁵, Abhishek Vadher^{*6}

¹Dow International Medical College.

²Senior Medical Officer, Health and Care Foundation.

³Doctor, Mamta Maternity and General Hospital, Umbergaon.

⁴Medical Student, BJ Medical and Civil Hospital, Ahmedabad.

⁵Medical Officer, Civil Hospital, Ahmedabad.

⁶Intensivist, Texas Multispeciality Hospital.

***Corresponding Author: Abhishek Vadher**

Intensivist, Texas Multispeciality Hospital.

Article Received on 13/10/2022

Article Revised on 02/11/2022

Article Accepted on 23/11/2022

ABSTRACT

Introduction: There is a high prevalence of low serum testosterone in type 2 diabetes mellitus patients. Many studies have reported that there is a high prevalence of low serum testosterone in men with type 2 diabetes mellitus. The symptoms of low serum testosterone are loss of libido, erectile dysfunction, reduced muscle mass, low energy, increased adiposity. In this study, we tried to determine the level of serum testosterone in type 2 diabetes mellitus patients. **Methods:** A total of 241 patients were taken in the study. Out of this, 121 patients had type 2 diabetes mellitus and 120 were normal. For diagnosis of diabetes HbA1c level of ≥ 6.5 was taken. Serum testosterone levels were measured in all the participants. **Results:** The mean age of the participants was 46.95 ± 6.89 in diabetic patients and 45.86 ± 5.45 in the controls. Average height was 168 ± 5.3 cm in the diabetic patients and 170 ± 4.2 cm in the normal population. Average weight was 73 ± 6.5 kg in the diabetic patients and 69 ± 7.4 in the normal population. The mean serum testosterone levels in the diabetes population were 312 ± 14.6 ng/dl and in the control population was 678 ± 17.5 ng/dl. On applying chi square test, the p value was calculated to be 0.02. This shows the difference is statistically significant. **Conclusion:** Type 2 diabetes mellitus is associated with low levels of serum testosterone levels.

INTRODUCTION

It is estimated that 285 million people worldwide are affected by diabetes mellitus. By 2030, around 438 million people will be affected by diabetes mellitus. About 66% of this population is in low to middle income countries.^[1] As compared to western population, Asian population has a much higher tendency to develop diabetes. This occurs at a younger age, at lower degrees of obesity and a much higher rate given for the same amount of weight compared to Western population.^[2] Many studies have reported that there is a high prevalence of low serum testosterone in men with type 2 diabetes mellitus.^[3,4, and 5] Some studies have also showed a co-relation between reduced total testosterone and insulin resistance and then subsequent development of diabetes mellitus.^[6,7] The symptoms of low serum testosterone are loss of libido, erectile dysfunction, reduced muscle mass, low energy, increased adiposity.^[8,9] Total testosterone is largely determined by circulating sex hormone binding globulin. In normal men, 54% testosterone is bound to albumin and other proteins, 44% is bound to sex hormone binding globulin and 2% is in unbound state. Some studies believe that

low levels of serum testosterone are associated with changes in the levels of FSH and LH. There is an ongoing controversy about whether the low serum levels of serum testosterone are associated with hypogonadotropic hypogonadotrophism or not. The study by Ali et al showed that in patients with diabetic neuropathy, low serum testosterone levels were associated with low FSH and low LH. In this study, we tried to find a co-relation between levels of serum testosterone and type 2 diabetes mellitus.

MATERIALS AND METHODS

This is a retrospective case-control study with 121 patients and 120 controls. A total of 121 patients were taken in the study who consistently attended the diabetes clinic from May 2017 to May 2018 were taken in the study. For diagnosis of diabetes HbA1c level of ≥ 6.5 was taken. Patients with type 1 diabetes mellitus, renal involvement, liver disease and alcoholism were excluded from the study. 120 controls were taken. These were age and height and weight comparable to the study population. A thorough physical examination, vital examination including temperature, pulse, respiratory

rate, blood pressure etc of all the cases and controls was done. After history and physical examination, complete blood count, urine albumin, sugar, urine microscopy, random blood sugar on admission, serum urea, serum creatinine, D level, HDL level, VLDL cholesterol, LDL/HDL ratio were done. ECG was also done of all the patients to

rule out any sign of ischemia, arrhythmia, hypertrophy or any other change. Serum testosterone levels were measured in all the participants. Whole data was entered in Excel sheet and analysed by EPI INFO software. All calculations were done statistically. P value of <0.05 was considered to be statistically significant.

RESULTS

Table 1: Baseline Characteristics of the Population.

	Diabetic patients	Normal population
Age (yrs)	47±7.8	47±8.8
Height (cm)	170±6.7	172±7.8
Weight (kg)	75±6.8	73±8.9
Smoking :Never	94	90
Smoking : Former	57	42
Smoking: Current	68	48
Hypertension: Yes	38	16
Hypertension: No	145	154
Dyslipidaemia : Yes	25	21
Dyslipidaemia: No	158	159
Duration of Diabetes	5.9±2.68	
Mean HbA1c	7.8±1.0	4.8±0.6

As shown in the table above, the baseline characteristics of all the participants are showed in the table. The average age of the diabetic patients was 47±7.8. The average age in the normal population was 47±8.8. Average height was 170±6.7cm in the diabetic patients and 172±7.8cm in the normal population. Average weight was 75±6.8kgs in the diabetic patients and 73±8.9 in the normal population. 94 patients with diabetes never smoked while 90 participants in the normal population were non-smokers. The number of former smokers in the diabetic patients and normal pupation was 57 and 42 respectively. Current smokers are 68/183 in the diabetic patients and 48/180 in the normal population. 38 patients in the diabetic population were hypertensives and 145 were normotensives. 16 patients were hypertensive in the normal population and 154 were normotensive. 25 patients had dyslipidaemia and 158 did not have dyslipidaemia in the diabetic population. 21 participants had dyslipidaemia and 159 participants did not have dyslipidaemia in the normal population. Mean duration of diabetes was 5.9±2.68. Mean HbA1c was 7.8 ±1.0 in the diabetic population and 4.8±0.6 in the normal population.

Table 2: Serum testosterone levels in patients and controls.

	Type 2 diabetes	Control	P value
Serum testosterone levels (ng/dl)	312±14.6	678±17.5	0.02

Serum testosterone level in the diabetic population was 312±14.6 ng/dl. In the normal population, it was 678±17.5ng/dl. After applying student t test, the p value was calculated to be 0.02. This is less than 0.05 which

shows that the difference in the two group is significant and not due to chance.

DISCUSSION

Many studies have shown that about 25% of patients with type 2 diabetes mellitus have low serum testosterone levels. About 4% have subnormal testosterone concentrations with high FSH and H.^[10] Some studies have also shown that low serum testosterone is associated with diabetes related sexual dysfunction. A study from Australia showed that 43% of type 2 diabetes patients have total testosterone levels less than 10.^[11] A study from United Kingdom showed that 355 men with type 2 diabetes mellitus have total testosterone levels of less than 8 and 25% had symptoms of hypogonadism associated with 8-12 total testosterone.[12] 33.2% type 2 diabetes patients had hypogonadism in a study in Egypt.[13] A study from Brazil also showed that free testosterone and total testosterone levels were low in type 2 diabetes patients.[14] The Endocrine society also recommends measuring the levels of testosterone of patients with type 2 diabetes on a regular basis.[15,6] Many cross-sectional and longitudinal studies have showed that with the increase in age the level of total testosterone reduces in men.[17, 18,19] Many studies have also showed a co-relation between BMI and low serum testosterone levels in type 2 diabetes mellitus. However, these studies are controversial. Some studies have showed that the association between BMI and serum testosterone is significant.[20,21] On the contrary, there is also a study that has showed no co-relation between BMI and low testosterone.[22] There are many limitations of this study. It is a cross-sectional study and so we could not find the trend of the serum testosterone levels in the participants. From our study and conclusion, we can strongly say that an early universal screening program

can help in diagnosis of low serum testosterone levels and testosterone supplementation can be started accordingly. We recommend that all patients of type 2 diabetes mellitus undergo screening for serum testosterone. A hormonal baseline can also be established for comparison in future follow-ups.

CONCLUSION

Type 2 diabetes mellitus is associated with low levels of serum testosterone levels in our study population with a p value significance of 0.02.

REFERENCES

1. International Diabetes Federation. IDF Diabetes Atlas. Epidemiology and Mobility. International Diabetes Federation. [Last accessed on 2011 March 1]. Available from: <http://www.idf.org/> [Ref list].
2. Diabetes in Asia: epidemiology, risk factors, and pathophysiology. Chan JC, Malik V, Jia W, Kadowaki T, Yajnik CS, Yoon KH, Hu FB *JAMA*, 2009 May 27; 301(20): 2129-40.
3. Prevalence of androgen deficiency in men with erectile dysfunction. Köhler TS, Kim J, Feia K, Bodie J, Johnson N, Makhlouf A, Monga M *Urology*, 2008 Apr; 71(4): 693-7.
4. Comparison of serum free testosterone, luteinizing hormone and follicle stimulating hormone levels in diabetics and non-diabetics men- a case-control study. Rezvani MR, Saadatjou SA, Sorouri S, Hassanpour Fard M *J Res Health Sci*, 2012 Dec 13; 12(2): 98-100.
5. Hypogonadism, erectile dysfunction, and type 2 diabetes mellitus: what the clinician needs to know. Tamler R, Deveney T *Postgrad Med*, 2010 Nov; 122(6): 165-75.
6. Low testosterone levels are common and associated with insulin resistance in men with diabetes. Grossmann M, Thomas MC, Panagiotopoulos S, Sharpe K, Macisaac RJ, Clarke S, Zajac JD, Jerums G *J Clin Endocrinol Metab*, 2008 May; 93(5): 1834-40.
7. Testosterone, SHBG and risk of type 2 diabetes in the second evaluation of the Pizarra cohort study. Soriguer F, Rubio-Martín E, Fernández D, Valdés S, García-Escobar E, Martín-Núñez GM, Esteva I, Almaraz MC, Rojo-Martínez G *Eur J Clin Invest*, 2012 Jan; 42(1): 79-85.
8. A practical guide to male hypogonadism in the primary care setting. Dandona P, Rosenberg MT *Int J Clin Pract*, 2010 May; 64(6): 682-96.
9. Androgen replacement therapy improves psychological distress and health-related quality of life in late onset hypogonadism patients in Chinese population. Zhang XW, Liu ZH, Hu XW, Yuan YQ, Bai WJ, Wang XF, Shen H, Zhao YP *Chin Med J (Engl)*, 2012 Nov; 125(21): 3806-10.
10. Testosterone, SHBG and risk of type 2 diabetes in the second evaluation of the Pizarra cohort study.
11. Soriguer F, Rubio-Martín E, Fernández D, Valdés S, García-Escobar E, Martín-Núñez GM, Esteva I, Almaraz MC, Rojo-Martínez G *Eur J Clin Invest*, 2012 Jan; 42(1): 79-85.
12. Low testosterone levels are common and associated with insulin resistance in men with diabetes.
13. Clinical and biochemical assessment of hypogonadism in men with type 2 diabetes: correlations with bioavailable testosterone and visceral adiposity. Kapoor D, Aldred H, Clark S, Channer KS, Jones TH *Diabetes Care*, 2007 Apr; 30(4): 911-7.
14. Serum testosterone levels in diabetic men with and without erectile dysfunction. Ghazi S, Zohdy W, Elkhayat Y, Shamloul R *Andrologia*. 2012 Dec; 44(6): 373-80.
15. Diabetes mellitus is associated with subnormal serum levels of free testosterone in men. Rhoden EL, Ribeiro EP, Teloken C, Souto CA *BJU Int*, 2005 Oct; 96(6): 867-70.
16. Update: Hypogonadotropic hypogonadism in type 2 diabetes and obesity. Dandona P, Dhindsa S *J Clin Endocrinol Metab*, 2011 Sep; 96(9): 2643-51.
17. Testosterone therapy in adult men with androgen deficiency syndromes: an endocrine society clinical practice guideline. Bhasin S, Cunningham GR, Hayes FJ, Matsumoto AM, Snyder PJ, Swerdloff RS, Montori VM *J Clin Endocrinol Metab*, 2006 Jun; 91(6): 1995-2010.
18. Low testosterone levels are common and associated with insulin resistance in men with diabetes. Grossmann M, Thomas MC, Panagiotopoulos S, Sharpe K, Macisaac RJ, Clarke S, Zajac JD, Jerums G *J Clin Endocrinol Metab*, 2008 May; 93(5): 1834-40.
19. A practical guide to male hypogonadism in the primary care setting. Dandona P, Rosenberg MT *Int J Clin Pract*, 2010 May; 64(6): 682-96.
20. Hypogonadism and diabetes. Betancourt-Albrecht M, Cunningham GR *Int J Impot Res.*, 2003 Aug; 15 Suppl 4: S14-20.
21. The role of obesity and type 2 diabetes mellitus in the development of male obesity-associated secondary hypogonadism. Saboor Aftab SA, Kumar S, Barber TM *Clin Endocrinol (Oxf)*, 2013 Mar; 78(3): 330-7.
22. Sex hormone changes in morbidly obese patients after vertical banded gastroplasty. Bastounis EA, Karayiannakis AJ, Syrigos K, Zbar A, Makri GG, Alexiou D *Eur Surg Res.*, 1998; 30(1): 43-7.
23. Hypogonadotropic hypogonadism in erectile dysfunction associated with type 2 diabetes mellitus: a common defect? Tripathy D, Dhindsa S, Garg R, Khaishagi A, Syed T, Dandona P *Metab Syndr Relat Disord*, 2003 Mar; 1(1): 75-80.