



RECEIVER OPERATING CHARACTERISTIC (ROC) PLOTS: AN EVALUATION TOOL FOR CARDIO METABOLIC RISK ASSESSMENT IN DIABETICS

*¹Sakina S. Raghiv and ²Sabiha A. Vali

¹Visiting Faculty, Department of Home Science, LAD and SRP College for Women, Nagpur, India.

²Professor and Ex Head, Post Graduate Department of Home Science, Rashtrasant Tukadoji Maharaj Nagpur University, Nagpur, India.

*Corresponding Author: Dr. Sakina S. Raghiv

Visiting Faculty, Department of Home Science, LAD and SRP College for Women, Nagpur, India.

Article Received on 29/11/2021

Article Revised on 19/12/2021

Article Accepted on 09/01/2022

ABSTRACT

The study explored optimal anthropometric indicator and cut - off points for cardio metabolic risk (CMR) in 200 type 2 female diabetics with a mean age of 54.29±10.55. Receiver Operating Characteristic (ROC) was performed independently and statistical significance was obtained corresponding to AUC of each parameter. Cut off values as determined by Youden's index were 29.32 kg/m² (BMI), 96.50 cm (WC), 0.95 (WHR) and 0.60 (WHtR). The optimum cut off for WHtR remained close to the standard cut off value of 0.5 indicating WHtR as the best predictive value for evaluating metabolic risk. The AUC of BMI was lowest; 0.602 (p=0.088) while WHtR showed the highest AUC of 0.692 (p=0.001) thereby confirming its ability to be a better predictor of cardio metabolic risk factors followed by WC (AUC 0.668 p= 0.005). The study validates the usefulness of ROC as an evaluation tool for identifying diabetics at cardio metabolic risk.

KEYWORDS: Cardio Metabolic Risk, Youden's index, Area under curve (AUC).

INTRODUCTION

The world is in the phase of epidemiological transition where it is being observed that death due to non-communicable diseases exceeds death due to communicable diseases. Obesity is the most common nutritional pandemic in the developed world and is associated with co morbidities such as coronary artery disease, type 2 diabetes mellitus and hypertension adding to morbidity and mortality (Gutch M., et al.). The global obesity epidemic indicates no signs of abating, and is fueling an explosion in number of diabetics worldwide. Obesity and type 2 diabetes are of serious health concern. The term 'diabesity' has been coined to define obesity dependent diabetes.

Cardio metabolic risk (CMR) in type 2 diabetics could be additive or synergistic for CVD. In type 2 diabetes an increased cardiovascular risk often exists for several years before the onset of biochemical hyperglycemia. During this phase obesity and insulin resistance are often present, associated with hypertension and dyslipidemia, usually referred as metabolic syndrome. These risk factors may lead to the early development of CHD (Haffner S., et al.).

Prevalence of CAD was reported to be significantly higher in abdominal obese patients (WHtR_>0.55) than

in patients without abdominal obesity. WHtR may be a better marker of central obesity and may predict CAD more accurately than BMI and WC (Siavash M., et al.). The present study explored the predictive value of WHtR for Cardio Metabolic Risk (CMR) in 200 type 2 female diabetics. The screening potential of WHtR for adult cardio metabolic risk in type 2 diabetics was analyzed by applying Receiving Operating Curves (ROC) and compared to other indices such as BMI, WC and WHR.

MATERIALS AND METHOD

Participants

The study was conducted at Diabetes Care and Research Centre, Nagpur, Maharashtra with due approval of the ethical committee. 200 type 2 diabetic females above the age of 25 years with diabetic history of at least 1 year were included in the study. Type 1 diabetics, pregnant women, smokers, alcoholics and subjects with hepatic, renal, endocrine disorders and those on lipid lowering agents were excluded from the study. Informed written consent was obtained from every participant.

Measures

Anthropometry

A predesigned questionnaire was used to note the anthropometric measurements (height, weight, waist circumference and hip circumference of the subjects.

Body Mass Index (BMI), Waist Hip Ratio (WHR) and Waist Height Ratio (WHtR) were determined and recorded. The threshold cut off values adopted for anthropometric parameters for all the female subjects were BMI \Rightarrow 22.9 kg/m², WC \Rightarrow 80 cm, WHR \Rightarrow 0.85 and WHtR \Rightarrow 0.50

Biochemical Parameters

All the biochemical tests were performed at the laboratory of the study center. Individual blood markers included Glycemic Profile (FBG, PMBG and HbA1c) and lipid profile (TC, LDL and HDL). Blood pressure was measured using an electronic blood pressure device while the subjects were seated following a five-minute relaxation period.

Metabolic syndrome (MS) was diagnosed according to the National Cholesterol Education Program (NCEP ATP-III) criteria. MS was recognized as having at least two of the following risk factors along with diabetes FBG (>110 mg/dl): WC (>80cm), high triglycerides (TG) (>150 mg/dl), low high-density lipoprotein cholesterol (HDL-C) (<50), blood pressure (>120/80 mmHg)

Statistical Analysis

SPSS version 18.0 was used for all the statistical analysis. Data on anthropometric measurements and cardio metabolic risk factors was compiled means and standard deviation were calculated. Pearson's correlation coefficient was applied to the data and interpreted.

All variables were converted to direct variables i.e., 1= no risk and 2= risk based on standard reference values for female diabetic subjects. A new variable 'risk' was created based on the five indicators of cardio metabolic risk factors wherein, if three or more risk factors were observed, it was recorded as 'risk'. The association of BMI, WC, WHR and WHtR with this new variable was estimated using Receiver Operating Characteristic (ROC) analysis. The cut off values for each index was determined by applying Youden's index.

RESULTS AND DISCUSSION

Anthropometric Measurements

Base line data of diabetic subjects is presented in **Table 1**. The study subjects showed a mean height of 153 ± 6.35 , mean weight of 65.85 ± 11.96 and mean BMI was 28.05 ± 4.60 . **Gupta K., et al.**, reported on the prevalence of obesity in diabetics with a higher percentage of subjects in the obese grade I category.

The mean WC measurement of the subjects was 99.99 ± 10.73 . WHR measures the ratio of WC to HC unlike BMI which calculates the ratio of weight to height. The mean WHR was also observed to be higher (0.98 ± 0.074) than the cut off value. In this study, in addition to conventional indices of central obesity, WHtR was included as a screening tool. A high mean value of 0.65 ± 0.06 as compared to the standard cut off

value of 0.5 was observed. **Ashwell M., and Gibson S.**, stated that compared to BMI, WHtR was more closely associated with CVD risk factors among both men and women

Biochemical Parameters

The mean levels of glycemic parameters of the subjects are presented in **Table 1**.

The mean FBG value was 147.68 ± 47.79 . Observation from **Figure 1** reveals a high FBG in 80% of subjects. PMBG of the subjects reflect a similar trend as observed in case of FBG. The mean PMBG of the subjects is 226.55 ± 69.91 with more than 90.5% showing values above 180mg/dl. The data on HbA1C shows a very high mean value of 8.11 ± 1.64 . A high percentage of subjects (86%) showed HbA1C values much higher than the normal value of <6.5% suggested for diabetics indicating uncontrolled diabetes. **Gupta K., et al.**, have also reported a similar trend in the mean fasting and post meal blood glucose levels falling above the recommendations for diabetics.

Obesity indices vs. Glycemic profile

Table 2 shows the distribution of diabetic subjects based on obesity indices vs. glycemic profile.

Observations of the study reveal that irrespective of normal or elevated BMI, WC, WHR and WHtR of all the diabetic subjects showed elevated values of FBG, PMBG and HbA1c.

Results of Pearson's correlation coefficient (**Table 3**) shows negative but highly significant relationship of BMI to FBG ($r=0.200$, $p=0.000$) PMBG ($r=0.207$, $p=0.000$) and significant to HbA1c ($r=0.116$, $p=0.020$). Very low negative correlation is observed between WC and glycemic parameters. The association is observed to be non-significant with respect to HbA1c. However, FBG and PMBG show a negative and highly significant relationship to WC ($r = -0.133$, $p=0.008$ and $r = -0.150$, $p=0.003$ respectively). Insignificant relationship of WHR to FBG, PPBG and HbA1c is observed. Observations from the table reveal a very low negative relationship of WHtR to all the three glycemic parameters viz., FBG ($r = 0.187$) PMBG ($r = -0.186$) and HbA1c ($r = -0.113$). However the relationship is observed to be highly significant between FBG, PMBG ($p=0.000$) and significant in HbA1c ($p=0.024$) to WHtR.

Cardio Metabolic Risk

Figure 2 shows the distribution of subjects based on the cardio metabolic risk as assessed by high values of WC, BP, TG, HDL and FBG. Central obesity and high FBG values were observed in 96% of subjects. Hypertension and high TG was seen in 69% and 62% subjects respectively. More than 50% of subjects had low HDL levels. Each of the conditions have an associated effect, but clustering together they become synergistic, leading

to CVD risk. Based on this risk assessment 79% of the diabetic subjects were at risk of CVD (**Figure 3**)

Receiver Operating Characteristic (Roc) Curve Analysis

ROC was performed for all the indices independently to determine the validity in discriminating risk and non-risk group for CVD. The point on the ROC, which represented the largest sum of sensitivity and specificity, was chosen to obtain the optimal cut-off point for each of these four measurements in predicting cardio metabolic risk. The validity assessment of each parameter was carried out in terms of area under curve (AUC), sensitivity and specificity. The statistical significance was obtained corresponding to AUC of each parameter.

Body Mass Index (BMI)

Figure 4 provides the ROC plot for BMI of females. The standardized data on BMI for females was subjected to ROC analysis, which resulted into an AUC of 0.602 [95% CI: 0.487 – 0.718] with a corresponding p-value of 0.088, indicating insignificantly higher area as compared to 0.5. The optimum sensitivity and specificity were 0.42 and 0.78 respectively for a cut-off BMI value of 29.32 kg/m².

Waist Circumference (WC)

Figure 5 provides the ROC plot for waist circumference of females. The standardized waist circumference data for females was used for ROC analysis which resulted into an AUC of 0.668 [95% CI: 0.553 – 0.783] with a corresponding p-value of 0.005, indicating significantly higher AUC as compared to null value of 0.5. The optimum sensitivity and specificity obtained were 0.653 and 0.667 respectively for a cut-off level of 96.50 cm.

WAIST HIP RATIO (WHR)

ROC plot was obtained for standardized waist hip ratio for females as shown in **Figure 6**. The AUC was 0.603 [95% CI: 0.485 – 0.722] with a p-value of 0.084, indicating insignificantly different AUC as compared to null value of 0.5. The optimum sensitivity and specificity obtained were 0.717 and 0.481 respectively for a cut-off point 0.955

Waist Height Ratio (WHtR)

ROC plot was obtained for standardized waist height ratio for females as shown in **Figure 7**. The AUC was 0.692 [95% CI: 0.581 – 0.802] with a p-value of 0.001, indicating significantly higher AUC as compared to null value of 0.5. The optimum sensitivity and specificity obtained were 0.809 and 0.591 respectively for a cut-off point 0.60

Table 4 presents the AUC for obesity indices. BMI and WHR show the lowest and almost similar AUC. WHtR shows the highest AUC thereby confirming its ability to be a better predictor of cardio metabolic risk factors followed by WC.

Park Y. S., and Kim J. S., also reported that WHtR has the best predictive value for evaluating the metabolic risk factors compared to BMI or WC. **Jayawardana R., et al.**, reported that in all adults, males and females the AUC of WHtR was significantly higher than that of BMI, WC and WHR in diabetes mellitus, pre diabetes, hypertension, metabolic syndrome and hypercholesterolemia. **Hong X., et al.**, also reported from their data on ROC curve analysis that AUC of WHtR was the highest in both males and females. They concluded that WHtR might serve as a simple but most effective index of hyperglycemia. **Schneider H.J., et al.**, reported that short subjects were at a higher risk and had a 30% higher prevalence of metabolic syndrome than tall subjects when grouped by WC and not by WHtR, and recommended that WHtR rather than WC should be included in the definition of metabolic syndrome.

Table 1: Baseline Characteristics of subjects.

Variables	Mean (n=200)	Std. Deviation	Std. Error Mean
Height (cm)	153.06	6.35	0.44
Weight (kg)	65.85	11.96	0.84
BMI (kg/m ²)	28.05	4.60	0.32
WC (cm)	99.99	10.73	0.75
HC (cm)	102.26	10.69	0.756
WHR	0.98	0.07	0.00
WHtR	0.65	0.06	0.00
FBG (mg/dl)	147.68	47.79	3.38
PMBG (mg/dl)	226.55	69.91	4.94
HbA1c (%)	8.110	1.64	0.11




Figure 1: Distribution of subjects (%) based on Glycemic Profile.

Table 2: Association of Glycemic Profile with Obesity Indices.

Obesity Indices		Percentage distribution	FBG	PMBG	HbA1c
			Mean	Mean	Mean
BMI	Normal (n=25)	12.5	170± 70.96	251.04± 95.71	8.3± 1.84
	Elevated (n= 175)	87.5	144.28± 41.75	221.48± 63.92	8.08± 1.62
WC	Normal (n=11)	5.5	160.82± 79.91	230.64± 84.26	8.52± 2.21
	Elevated (n=189)	94.5	146.72± 44.53	224.86± 68.36	8.09± 1.62
WHR	Normal (n=13)	6.5	133.92± 34.63	209.69± 66.89	7.62± 1.43
	Elevated (n=187)	93.5	148.44± 47.62	226.25± 69.29	8.14± 1.66
WHtR	Normal (n=2)	1	138± 38.18	243.5± 61.51	8.65± 0.49
	Elevated (n=198)	99	147.59± 47.10	224.99± 69.28	8.1± 1.65

Table 3: Correlation of Obesity Predictors with Glycemic Profile of Subjects.

Obesity Indicators		FBG	PMBG	HbA1c
BMI	r	-0.200	-0.207	-0.116
	p value	0.000	0.000	0.020
	p summary	HS	HS	S
WC	r	-0.133	-0.150	-0.065
	p value	0.008	0.003	0.196
	p summary	HS	HS	NS
WHR	r	0.093	0.075	0.082
	p value	0.065	0.132	0.101
	p summary	NS	NS	NS
WHtR	r	-0.187	-0.186	-0.113
	p value	0.000	0.000	0.024
	p summary	HS	HS	S




Figure 2: Distribution of Subjects Based on Presence of Cardio Metabolic Risk Factors (%)




Figure 3: Distribution of Subjects Based on Presence or Absence of Risk




Figure 4: ROC Plot for Body Mass Index.




Figure 5: ROC Plot for Waist Circumference.




Figure 6: ROC Plot for WHR.




Figure 7: ROC Plot for WHtR.

Table 4: AUC of Obesity Indices.

S. No.	Parameter	AUC	p-value
1	BMI	0.602	0.088
2	WC	0.668	0.005
3	WHR	0.603	0.084
4	WHtR	0.692	0.001

CONCLUSION

The worldwide prevalence of metabolic syndrome is increasing. This increasing trend has been observed in Asian countries. The most effective screening measure must be practical as well as effective. BMI requires measures of weight and height, WC measures central obesity alone and has the complexity of gender differences. Waist hip ratio does not take into account height in defining obesity and hence may under or over evaluate the cardio metabolic risk in tall and short individuals. WHtR requires measures of WC along with height. Uncontrolled diabetes was a significant observation of the study. In spite of being under oral hypoglycemic treatment, the study subjects reflected extremely high FBG, PMBG and HbA1c levels exceeding the standard cut off criteria. Statistical interpretation using Receiver Operating Characteristics (ROC) showed that the Area under Curve (AUC) for WHtR was higher than AUC for BMI and WHR thereby confirming its ability to be a better predictor of cardio metabolic risk factors followed by WC. A large percentage of diabetic subjects in the present study are under the risk of cardio vascular diseases as assessed by different screening tools for obesity and cardio metabolic risks and therefore reflect a gloomy situation that needs immediate intervention.

ACKNOWLEDGEMENT

The authors express their gratitude to Dr. Sunil Gupta, Diabetologist and Director of Sunil's Diabetes Care and Research Centre, Nagpur for granting permission to conduct this study at his research centre.

REFERENCES

1. Ashwell M., Gibson S., Waist to height ratio is a simple and effective obesity screening tool for cardiovascular risk factors: Analysis of data from the British National Diet and Nutrition Survey of adults aged 19-64 years, 2009; 2(2): 97-103
2. Gupta K., Vali S., Gupta S., Measures of Body composition amongst people with obese vs. non obese type 2 diabetics *Research Journal*, 2016; 12(2): 51-58.
3. Gutch M., Kumar S., Abhinav G., Gupta K., Razi M. S., Irisin: A clue for metabolic disorders. *Journal of Obesity and Metabolic Research*, 2016; (3)1: 41-45.
4. Haffner S., Stern M., Hazuda H., Mitchell B., Patterson J., Cardiovascular risk factors in confirmed prediabetic individuals. *JAMA*, 1990; 263: 2893-2898.
5. Hong X., Li J., Liang Y., Wang Z., Xu F., The relationship between body mass index, waist circumference, waist-to-hip ratio, waist-to-height ratio and hyperglycemia: a three-year follow-up study in Nanjing. *Chinese Journal of Epidemiology*, 2009; 30(3): 209-213.
6. Jayawardana R., Ranasinghe P., Sheriff M., Matthews D., Katulanda P., Waist to height Ratio: A Better Anthropometric Marker of Diabetes and Cardio Metabolic Risks in South Asian Adults. *Diabetes Research and Clinical Practice*, 2013; 99(3): 292-294.
7. National Institutes of Health 2001 *The third report of the National Cholesterol Education Program Expert Panel on Detection, Evaluation, and Treatment of High Blood Cholesterol in Adults (Adult Treatment*

Panel III). NIH Publication 01–3670. Bethesda, MD: National Institutes of Health .

8. Park Y., Kim J., Association between Waist to Height Ratio and Metabolic Risk Factors in Korean Adults with Normal Body Mass Index (BMI) and Waist Circumference. *The Tohoku Journal of Experimental Medicine*. 2012; 228(1): 1-8.
9. Schneider H., Klotsche J., Silber S., Stalla G., Wittchen H., Measuring Abdominal Obesity: Effects of Height on Distribution of Cardio Metabolic Risk Factors Using Waist Circumference and waist-to-height ratio. *Diabetes Care*, 2011; 34-37.
10. Siavash M., Sadeghi M., Salarifar F., Amini M., Shojaee M., Comparison of Body Mass Index and Waist Height Ratio in Predicting Definite Coronary Artery Disease. *Annals of Nutrition and Metabolism*, 2014; 53(3): 62-63.