

**NOVEL SYNTHESIS AND STUDY OF IMPACT OF SUBSTITUTED 1,3-THIAZINE AND ITS NANOPARTICLES ON PHYTOTIC GROWTH OF SOME FLOWERING PLANTS**

**Chhaya D. Badnakhe\***

Department of Chemistry, Dr.Manorama and Prof.H.S.Pundkar, Arts, Commerce and Science College, Balapur, Dist. Akola.

\*Corresponding Author: Chhaya D. Badnakhe

Department of Chemistry, Dr.Manorama and Prof.H.S.Pundkar, Arts, Commerce and Science College, Balapur, Dist. Akola.

Article Received on 04/08/2022

Article Revised on 25/08/2022

Article Accepted on 15/09/2022

**ABSTRACT**

The synthesis, spectral analysis and biological activities of 4-phenyl-2-hydroxy-chlorosubstituted-2-imino-1,3-thiazines have been carried out. In this case 4-(2'-hydroxy-3',5'-dichlorophenyl)-6-(4''-nitrophenyl)-2-imino-3,6-dihydro-1,3- thiazine (A) has been screened. The compound A was synthesized from 2'-hydroxy-3,5-dichlorophenyl-4-(4''-nitrophenyl)chalcone (a) by the action of thiourea, The compound (a) was synthesized from 2'-hydroxy-3',5'-dichloroacetophenone by the action of p-nitrobenzaldehyde in ethanol and 40% NaOH. The nanoparticles of the compounds A have been prepared by using ultrasonic technique. The titled compound and its nanoparticles were screened for their growth promoting activity on some flowering plants viz.. *Crysanthemum coronarium*, *Dahalia pinnata*, *Verbena officinalis*, *Iberis amara*.

**KEYWORDS:** Chalcone, thiazine, thiourea, growth promoting activities.

**INTRODUCTION**

Thiazine is a six membered ring system, which contains two hetero atoms [N and S] placed in a heterocyclic ring at 1, 3 positions. Many workers have synthesized different 1,3-thiazines. The researchers have reported the synthesis of several thiazines<sup>[1-6]</sup> and also their potent biological activities such as blood platelet aggregation inhibitors<sup>[7]</sup>, antibacterial<sup>[8-9]</sup> antiallergic<sup>[10]</sup>, anticholesterenic<sup>[11]</sup> and antifungal.<sup>[12]</sup> Moreover thiazine nucleus is a pharmacophore of cephalosporin that occupy a very important place in the field, of antibiotics and drug chemistry. Chalcones and their analogues having  $\alpha,\beta$ -unsaturated carbonyl system are very versatile substrates for the evolution of various reactions and physiologically active compounds. The reaction of thiourea with  $\alpha,\beta$ -unsaturated ketones also results in the formation of 1,3-thiazines. The chlorosubstituted thiazines with amino group at position 2 in the ring exhibit promising biological activities.<sup>[13-16]</sup>

In the present study, the chlorosubstituted 1,3-thiazine (A) has been prepared along with their nanoparticles and

screened them for their growth promoting activity on some flowering plants viz. *Crysanthemum coronarium*, *Dahalia pinnata*, *Verbena officinalis*, *Iberis amara*.

**EXPERIMENTAL**

All the glassware's used in the present work were of pyrex quality. Melting points were determined in hot paraffin bath and are uncorrected. The purity of compounds was monitored on silica gel coated TLC plate. IR spectra were recorded on Perkin-Elmer spectrophotometer in KBr pellets,  $H^1$  NMR spectra on spectrophotometer in  $CDCl_3$  with TMS as internal standard. UV spectra were recorded in nujol medium. The analytical data of the titled compounds was highly satisfactory. All the chemicals used were of analytical grade. All the solvents used were purified by standard methods. Physical characterisation data of all the compounds is given in Table 1.

**Table 1: Characterisation data of newly synthesized compounds.**

Compounds	Molecular formula	M.P. in °C	% of yield	% of element			
				C	H	N	S
	$C_8H_6O_2Cl_2$	54	80	47.90/48	2.95/3		
a	$C_{15}H_9O_4NCl_2$	250	70	53.10/53.25	2.40/2.66	3.98/4.18	
A	$C_{16}H_{11}O_3N_3Cl_2S$	120	70	48.50/48.60	2.35/2.53	10.40/10.63	8.00/8.10

**2'-Hydroxy 3',5'-dichloroacetophenone**

2'-Hydroxy-5-chloroacetophenone (3g) was dissolved in acetic acid (5 ml), and mixed with sodium acetate (3g). To this reaction mixture chlorine in acetic acid reagent (40 ml; 7.5 w/v) was added dropwise with stirring. The temperature of the reaction mixture was maintained below 20°C. The mixture was allowed to stand for 30 minutes and then poured into water. A pale yellow solid thus obtained was filtered, dried and crystallized from ethanol to yield the compound.

**Preparation of 2'-hydroxy-3,5-dichlorophenyl-4-(4''-nitrophenyl)-chalcone (a)**

2'-Hydroxy-3',5'-dichloroacetophenone (0.1 mol) was dissolved in ethanol (50 ml) and p-nitrobenzaldehyde (0.1 mol) was added gradually to the solution and the mixture was heated to boiling. Then aqueous sodium hydroxide solution [40%; 40 ml] was added dropwise with constant stirring. The mixture was stirred mechanically at room temperature for about half an hour and kept for overnight. It was then acidified by hydrochloric acid (10%) solution. The solid product thus separated, was filtered, and washed with sodium bicarbonate (10%) followed by water. Finally it was crystallized from ethanol acetic acid mixture to get the compound (a).

**Preparation of 4-(2'-hydroxy-3',5'-dichlorophenyl)-6-(4''-nitrophenyl)-2-imino-3,6-dihydro-1,3-thiazine (A)**

2'-Hydroxy-3,5-dichlorophenyl-4-(4''-nitrophenyl)-chalcone (a) (0.01 mol) and thiourea (0.02 mol) were dissolved in ethanol (30 ml). To this aqueous KOH solution (0.02 mol) was added. The reaction mixture was refluxed for three hours, cooled and diluted with water then acidified with 1:1 HCl. The product thus obtained was crystallized from ethanol to get the compound (A).

The newly synthesized compound was characterised on the basis of elemental analysis, molecular determination, UV, IR, NMR. spectral data.

**The UV, IR, and NMR spectral data-Compound (A)****UV: Spectrum No. 1**



The UV-Vis spectrum of the compound A reported in dioxane showed  $\lambda_{\max}$  value 495 nm corresponding to  $n \rightarrow \pi^*$  transition.

**IR (KBr):-Spectrum No. 2**

3365.34  $\text{cm}^{-1}$  (-OH phenolic), 2925.2  $\text{cm}^{-1}$  (aliphatic -C-H stretching), 3068.24  $\text{cm}^{-1}$  (aromatic -C-H stretching), 3017.30  $\text{cm}^{-1}$  (-N-H stretching), 1648.7  $\text{cm}^{-1}$  (-C=N stretching), 1342  $\text{cm}^{-1}$  [(C-N) (C-NO<sub>2</sub>) stretching], 738.13  $\text{cm}^{-1}$  (C-Cl stretching in aliphatic), 1177.7  $\text{cm}^{-1}$  (C-Cl stretching in aromatic).

**PMR:-Spectrum No. 3**

$\delta$  1.2 (s, 1H, -C-H);  $\delta$  2.7 (s, 1H, =N-H);  $\delta$  3.6 (s, 1H, =N-H);  $\delta$  3.7 (s, 1H, =C-H);  $\delta$  7.6 to 8.1 (m, 6H, Ar-H);  $\delta$  12.6 (s, 1H, O-H).

**Spectrum No. 01.****Spectrum No. 02.**



The plants from group B were sprayed with the solution of test compounds at weekly intervals. The field experiments were conducted to compare the treated plants of group B with untreated plants of controlled group A. In this context, the observations were recorded on 7, 14, 21, 28, 35, 42, 45, 56, 63, 70, 77, 84, 91 days after sowing corresponding to early vegetative, late vegetative, flowering, pod filing and pod maturation,

with special reference to number of leaves and height of shoots.


The results of field's experiments are tabulated in the tables 2,3 and 4.

Table (2): Activity of the test compounds A

**Table No. (02): 4-(2'-Hydroxy-3',5'-dichlorophenyl)-6-(4''-nitrophenyl)-2-imino-3,6-dihydro-1,3-thiazine (A).**

Periodicity of Observations [in days]	<i>Crysanthemum coronarium</i>				<i>Dahalia pinnata</i>				<i>Verbena officinalis</i>				<i>Iberis amara</i>			
	Shoot height		No. of leaves		Shoot height		No. of leaves		Shoot height		No. of leaves		Shoot height		No. of leaves	
	C	T	C	T	C	T	C	T	C	T	C	T	C	T	C	T
7	1.0	1.0	1	1	2.5	1.5	2	2	4.4	4	2	2	2	2	2	3
14	1.2	1.2	2	2	7.5	7	2	2	10	8	2	2	2.1	2.5	2	3
21	1.3	1.4	7	10	8	12	2	4	15	11	3	5	2.3	2.8	3	4
28	1.5	1.6	9	11	9	19	3	6	16	18	4	6	2.5	2.7	4	5
35	1.6	1.8	10	12	11	26	4	7	18	19	5	9	2.8	3.0	5	7
42	1.8	2.0	12	15	17	42	5	8	19	20	7	12	3.0	3.4	6	8
49	2.3	3.5	14	18	25	48	6	8	20	21	8	14	3.5	3.9	8	9
56	3.6	4.0	16	22	28	52	7	9	23	25	10	17	3.8	4.5	10	12
63	5.5	6.7	18	24	31	55	8	10	25	30	12	18	4.2	5.0	12	14
70	7	12	20	28	34	60	9	11	27	32	14	20	4.6	5.4	14	16
77	14	19	22	30	36	63	10	13	28	35	16	23	5.5	7.0	16	18
84	20	24	24	32	38	65	11	15	29	36	18	25	7.2	14	20	24
91	24	30	26	36	40	68	12	17	36	38	20	27	8.2	17	25	29

**Impact of the compound 4-(2'-Hydroxy-3',5'-dichlorophenyl)-6-(4''-nitrophenyl)-2-imino-3,6-dihydro-1,3-thiazine (A) on phytotic growth of *Crysanthemum coronarium***



Impact of the compound 4-(2'-Hydroxy-3',5'-dichlorophenyl)-6-(4''-nitrophenyl)-2-imino-3,6-dihydro-1,3-thiazine (A) on phytotoxic growth of *Dahalia pinnata*

Control



Treated



15 days



30 days



45 days



90 days

Impact of the compound 4-(2'-Hydroxy-3',5'-dichlorophenyl)-6-(4''-nitrophenyl)-2-imino-3,6-dihydro-1,3-thiazine (A) on phytotoxic growth of *Verbena officinalis*

Control

Treated



15 days



30 days




45 days



90 days

Impact of the compound 4-(2'-Hydroxy-3',5'-dichlorophenyl)-6-(4''-nitrophenyl)-2-imino-3,6-dihydro-1,3-thiazine (A) on phytotic growth of *Iberis amara*



### RESULT AND DISCUSSION

The titled compound and its nanoparticles were screened for their growth promoting activity on test flowering plants viz, *Crysanthemum coronarium*, *Dahlia pinnata*, *Verbena officinalis*, *Iberis amara*.

When a comparison of morphological characters was made between those of treated and control group plants, it was interesting to note that all the treated plants exhibited significant shoot growth and considerable increase in the number of leaves as compared to those of untreated ones.

### ACKNOWLEDGEMENTS

The authors are thankful to the Principal, Dr.D.H. Pundkar, Dr.Manorama & Prof.H.S.Pundkar, Arts, Commerce & Science College, Balapur, Dr. B.B. Wankhade, Principal, Malkapur Vidnyan Mahavidyalaya, Malkapur for providing necessary facilities to carry out the research work.

### REFERENCES

- Swarnkar P.K., Kriplani B., Gupta G.N., Oijha K.G., synthesis and antimicrobial activity of some new phenothiazine derivatives, *E.J. Chem.*, Jan. 2007; 4(1): 14-20.

2. Kakade B.S., "Synthesis in heterocyclic compounds (Role of DMSO as a solvent," Ph.D. Thesis, Nagpur University, 1981.
3. Chincholkar M.M., and Ramekar M.A. *J. Ind. Chem. Soc.*, 1994; 71(4): 199.
4. S.P. Rathod, A.P. Charjan and P.R. Rajput, *Rasayan J. Chem.*, 2010; 3(2): 363-367.
5. Dabholkar V.V. and F.Y. Ansari, *Indian J. Chem.*, Nov. 2008; 47 B.: 1759-1761.
6. D.H. Morey and S.N. Patil *ori., J. Chem.*, Mar., 2002.
7. C. Brown and R.N. Davidson, *Adv. Heterocycl. Chem.*, 1985; 38: 135.
8. P. Descacq, A. Nubrich, M. Capdepuy and G. Devanuz, *Eur, J. Med. Chem.*, 1990; 25: 285.
9. M.I. Younes, H.H. Abbas, and S.A.M. Metwally, *Arch, Pharma*, 1987; 230.
10. D.T. Witiar, M.E. Wolff and R.C. Covestri, "In Berger", *S. Medicinal Chemistry*", part III, Edi. Willey, Newyork, 1981; 603.
11. M. Crewzet and F. Helene, *Eurpat App. Ep.*, 121, 489, *Chem. Abstr.*, 102, 1985, 787244, *Chem., Abstr.*, 1978; 89: 108943 m..
12. F.M. Dean, K.A. Thakur and C.H. Gill, "Naturally occuring oxygen ring compounds", *J. Indian Chem. Soc.*, 1983; 60: 668.
13. P. Valenti, A. Bisi, A. Rampa, F. Belluti, S. Gobbi, A. Zampiron, M. Carrara, *Biosy, Med. Chem.*, 2002; 239.
14. Y.Q. Shi, J. Fukai, H Sakagami, W.J. Chang, P.Q. Yang, F.P. Wang, T. Nomura, *J. Nat. Prod.*, 2001; 64: 181.
15. G.J. Reddy, D. Latha, K.S. Rao, *Heterocycl. Commun.*, 2004; 10: 279.
16. T. Ghosh, S. Saba, C. Bandyopadhya, *Synthesis*, 2005; 11: 1845.