

**VALIDATED UV-SPECTROPHOTOMETRIC METHOD FOR SIMULTANEOUS  
ESTIMATION OF BENIDIPINE HYDROCHLORIDE AND TELMISARTAN IN BULK  
AND PHARMACEUTICAL DOSAGE FORM**

**M. M. Eswarudu\*, P. Srinivasa Babu, N. Sudhakar Reddy, G. Ouchitya, B. Vinay Kumar, S. Bhupathi Naidu,  
A. Prasanthi**

Department of Pharmaceutical Analysis, Vignan Pharmacy College, Vadlamudi, 522213, Andhra Pradesh, India.

\*Corresponding Author: M. M. Eswarudu

Department of Pharmaceutical Analysis, Vignan Pharmacy College, Vadlamudi, 522213, Andhra Pradesh, India.

Article Received on 11/07/2022

Article Revised on 01/08/2022

Article Accepted on 21/08/2022

**ABSTRACT**

A simple, economical, accurate, precise and less time-consuming UV Spectrophotometric method has been developed and validated for simultaneous estimation of Benidipine Hydrochloride and Telmisartan in bulk and pharmaceutical dosage form. In this method Benidipine HCl and Telmisartan are exhibits maximum absorbance ( $\lambda$  max) at 237 nm and 296 nm with methanol as the solvent. The method was validated as per the International Conference on Harmonization (ICHQ2R1) guidelines. Drugs followed the linearity in the concentration range of 1-5  $\mu$ g/mL and 10-50  $\mu$ g/mL with correlation coefficient ( $r^2$ ) of 0.9992 for Benidipine HCl and 0.9996 Telmisartan respectively. The validity of the proposed method was assessed by applying the standard addition technique where the percentage recovery of the added standard was found to be 99.63 and 100.31 for Benidipine HCl and Telmisartan. The limit of detection and quantification were calculated and found to be 0.1354 $\mu$ g/mL and 0.4103 $\mu$ g/mL and 1.6823  $\mu$ g/mL and 5.0980  $\mu$ g/mL for Benidipine HCl and Telmisartan respectively. The proposed method is recommended for routine analysis of Benidipine Hydrochloride and Telmisartan in bulk and pharmaceutical dosage forms in regular quality control testing laboratories.

**KEYWORDS:** Benidipine Hydrochloride, Telmisartan, UV Spectrophotometry, Beer's Law, Validation.

**INTRODUCTION**

Hypertension (HT) is a very common disorder, particularly for past middle age. It is not a disease in itself, but is an important risk factor for cardiovascular mortality and morbidity. For improved treatment of hypertension, Telmisartan and Benidipine HCl is the newer combination in market, this combination was developed to improve medication for Stage II hypertension.

Benidipine Hydrochloride (BEN HCl) is a dihydropyridine type of Calcium channel blockers used for the treatment of hypertension and angina pectoris. Chemically it is 5-O-[(3R)-1-benzylpiperidin-3-yl]3-O-methyl(4R)-2,6-dimethyl-4-(3-nitrophenyl)-1,4-dihydropyridine-3,5-dicarboxylate; It is very soluble in formic acid, soluble in methanol, slightly soluble in ethanol and practically insoluble in water. It acts by inhibiting trans membrane  $Ca^{2+}$  influx through the voltage dependent channels of smooth muscles in vascular walls.<sup>[1]</sup>

Telmisartan (TEL) belongs to Angiotensin receptor blockers with chemical name 2-[4-[[4-methyl-6-(1-

methylbenzimidazol-2-yl)-2-propylbenzimidazol-1-yl]methyl] phenyl] benzoic acid. It is soluble in strong base & methanol, and sparingly soluble in strong acid (except HCL). It interferes with the binding of angiotensin II to the angiotensin II AT<sub>1</sub>-receptor by binding reversibly and selectively to the receptors in vascular smooth muscle and the adrenal gland. Angiotensin II is a vasoconstrictor, which also stimulates the synthesis and release of an aldosterone blockage of its effects results in decrease in systemic vascular resistance.<sup>[2]</sup>

Both combination of Telmisartan and Benidipine drugs are used for the treatment of hypertension effectively. They work by relaxing the blood vessels and making the heart more efficient at pumping blood throughout the body. Literature survey revealed that the reported methods like UV and stability indicating RP-HPLC methods development and validation for estimation of Telmisartan and Benidipine HCl in bulk and pharmaceutical dosage form individually and in fixed dose combination.

The literature survey revealed that some UV methods.<sup>[3-7]</sup> and RP-HPLC methods.<sup>[8-9]</sup> are reported for the

determination of BEN HCl and some UV methods.<sup>[10-14]</sup> and RP-HPLC methods.<sup>[15-22]</sup> are reported for the determination of TEL individually and in both combination UV method.<sup>[23]</sup> and HPLC methods.<sup>[24-27]</sup> Present study involves the development and validation of a new UV Spectrophotometric method for the

determination of BEN HCl and TEL in bulk and in its pharmaceutical dosage form with good accuracy and precision. The analytical method was validated according to ICHQ2R1 validation parameters.<sup>[28-29]</sup> Chemical structure of Benidipine HCl and Telmisartan are shown in Figure 1.




Figure 1: Chemical structure of (a) Benidipine Hydrochloride (b)Telmisartan.

## MATERIAL AND METHODS

### Chemicals and Reagents

Benidipine Hydrochloride ( $\geq 99\%$ ), was bought from Yarrow Chem Products, Mumbai, India. and Telmisartan ( $\geq 98\%$ ) standard drugs (API) were obtained as gift samples from Dr. Reddy's Laboratories, Hyderabad, India. HPLC grade Methanol was purchased from Thermo Fisher Scientific India Pvt. limited, Mumbai, India. Water purified by the Milli-Q water purification system was used in the study. The rest of the chemicals and reagents were procured from standard commercial supplier.

### Equipment

Shimadzu (UV-1780) Double beam UV-Visible spectrophotometer with 1cm matched quartz cells was used for the measurement of absorbance. Shimadzu-AX-200 electronic balance was used for weighing the samples. Citizen-Ultrasonicator and Class 'A' volumetric glassware's were used.

### Selection of Solvent for Analysis

In the present study the UV spectra of BEN HCl and TEL were obtained from different solutions (Methanol, Acetonitrile, Distilled water, Formic acid, Dimethyl sulfoxide, Combination of Methanol and Water (9:1) Methanol and Formic Acid (5:5)) were studied. The two drugs were freely soluble in Methanol, Acetonitrile, Formic acid, Dimethyl sulfoxide, Combination of Methanol and Water (9:1), Methanol and Formic Acid (5:5). At the end of these studies, Methanol was chosen as solvent for studied drugs.

### Selection of detection wavelength for simultaneous estimation

In the present study the drug solutions of BEN HCl (5  $\mu\text{g/mL}$ ) and TEL (50  $\mu\text{g/mL}$ ) were prepared and scanned

over a range of 200-400 nm. It was observed that the drugs showed maximum absorbance at 237 and 354 nm for BEN HCl and 296 nm for TEL. The overlay spectra of BEN HCl and TEL (5  $\mu\text{g/mL}$  and 50  $\mu\text{g/mL}$ ) is shown in Figure 2. which was chosen as the detection wavelength (Isosbestic wavelength 245 nm) for the determination of Benidipine HCl and Telmisartan.

### Preparation of standard and Calibration curve solution of analytes

Primary stock solutions of BEN HCl and TEL were prepared separately by dissolving accurately weighed 10 mg of each pure drug samples in methanol in 100 mL volumetric flasks and made the volume up to the mark using same solvent to produce final concentration of 100  $\mu\text{g/mL}$  each of BEN HCl and TEL respectively. Appropriate aliquots (0.1, 0.2, 0.3, 0.4, 0.5 mL and 1, 2, 3, 4, 5 mL) of prepared standard solutions of BEN HCl and TEL we transferred into series of 10 mL volumetric flasks and diluted and made up to the mark with methanol to obtain final concentration of 1-5  $\mu\text{g/ml}$  of BEN HCl and 10-50  $\mu\text{g/ml}$  of TEL respectively. The above solutions were scanned over the range of 200 nm to 400 nm against reagent blank. The absorbances of each solution were measured at 245 nm against methanol as blank. A calibration curve was prepared by plotting absorbance versus concentration.




Figure 2: Overlay Spectra of Benidipine HCl and Telmisartan (5 µg/mL & 50 µg/mL).

### Estimation of Benidipine HCl and Telmisartan in Tablets

For the analysis of dosage form, twenty tablets of Benidin T (Benidipine HCl and Telmisartan -4 mg and 40 mg) were ground to fine powder and mixed thoroughly. A quantity of powder equivalent to 10 mg of each of the drug was transferred to 100 mL volumetric flask and dissolved in about 20 mL methanol sonicated for 10 min and volume was made up to mark with the same solvent. The insoluble excipients were separated by filtration through Whatman filter paper. After suitable dilution, absorbance of the prepared sample solutions (5 µg/mL and 50 µg/mL) of Benidipine HCl and

Telmisartan was recorded against the reagent blank at 245 nm.

### RESULTS AND DISCUSSION

Method validation was performed by following the International Conference on Harmonization (ICHQ2R1) guidelines.

**Linearity:** Fresh aliquots were prepared from standard stock solution ranging from 1-5 µg/mL and 10-50 µg/mL of Benidipine HCl and Telmisartan. The absorbance values of each concentration were recorded at 245 nm. For this study methanol was used as a blank. Results of Linearity study were shown in Table 1 and Figure 3.

Table 1: Linearity results of Benidipine HCl and Telmisartan.

S. No.	Benidipine HCl		Telmisartan	
	Concentration (µg/mL)	Absorbance	Concentration (µg/mL)	Absorbance
1	1	0.213	10	0.223
2	2	0.446	20	0.421
3	3	0.639	30	0.623
4	4	0.852	40	0.832
5	5	1.045	50	1.021




Figure 3: Linearity curves of Benidipine HCl and Telmisartan.

**Accuracy:** Accuracy of the developed method was confirmed by performing recovery studies at three different concentration ranges, each one in triplicate. From the recovery studies it was clear that the method

remains very accurate for quantitative estimation of tablet as the statistical results were within the acceptance range. Results of accuracy study were shown in Table 2

**Table 2: Accuracy data of Benidipine HCl and Telmisartan.**

S.No.	Benidipine HCl				Telmisartan			
	Amount added ( $\mu\text{g/mL}$ )	Amount found ( $\mu\text{g/mL}$ )	% Recovery	mean% recovery	Amount added ( $\mu\text{g/mL}$ )	Amount found ( $\mu\text{g/mL}$ )	% Recovery	Mean% recovery
1	1	0.97	97	97.66	10	10	100	99.8
2	1	0.96	96		10	9.97	99.5	
3	1	1	100		10	10	100	
4	3	2.97	99.5	99.26	30	30.02	100.1	100.1
5	3	2.94	98		30	30	100	
6	3	3.01	100.3		30	30.07	100.4	
7	5	5	100	99.33	50	50.02	100.09	100.3
8	5	4.98	99.6		50	50.04	100.1	
9	5	4.92	98.4		50	50.09	100.3	

**Precision:** The intraday and interday precision were executed by analyzing six independent analyses of 3  $\mu\text{g/mL}$  and 30  $\mu\text{g/mL}$  of Benidipine HCl and Telmisartan. The standard deviations, relative standard

deviation was calculated and results of the study are acceptable and can be considered to be very reasonable the results are summarized in Table 3 and Table 4

**Table 3: Intra-day precision data of proposed method.**

S.No.	Conc. ( $\mu\text{g/mL}$ )	Benidipine HCl			Conc. ( $\mu\text{g/mL}$ )	Telmisartan		
		FN	AN	EN		FN	AN	EN
1	3	0.639	0.638	0.637	30	0.623	0.622	0.621
2	3	0.638	0.638	0.635	30	0.622	0.622	0.623
3	3	0.639	0.637	0.636	30	0.623	0.621	0.621
4	3	0.638	0.638	0.637	30	0.621	0.622	0.622
5	3	0.639	0.636	0.637	30	0.623	0.623	0.621
6	3	0.637	0.637	0.636	30	0.622	0.621	0.622
Mean		0.638	0.637	0.636		0.622	0.621	0.621
SD		0.0008	0.0008	0.0008		0.0008	0.0007	0.0008
%RSD		0.127	0.128	0.128		0.131	0.121	0.131

**Table 4: Inter-day precision (Reproducibility) data of proposed method.**

S.No.	Conc. ( $\mu\text{g/mL}$ )	Benidipine HCl			Conc. ( $\mu\text{g/mL}$ )	Telmisartan		
		Day 1	Day 2	Day 3		Day 1	Day 2	Day 3
1	3	0.636	0.630	0.634	30	0.619	0.620	0.617
2	3	0.635	0.628	0.633	30	0.618	0.620	0.615
3	3	0.634	0.629	0.632	30	0.620	0.619	0.614
4	3	0.636	0.630	0.630	30	0.619	0.620	0.616
5	3	0.634	0.629	0.632	30	0.619	0.618	0.618
6	3	0.635	0.628	0.629		0.618	0.619	0.617
Mean		0.635	0.629	0.631		0.618	0.619	0.616
SD		0.0008	0.0008	0.0018		0.0007	0.0008	0.0014
%RSD		0.128	0.142	0.294		0.121	0.131	0.238

**Limit of Detection and Limit of Quantification:** The limit of detection and limit of quantification of Benidipine and Telmisartan by proposed method were determined using calibration curve.

LOQ and LOD were calculated as

$$\text{LOD} = 3.3 \times \text{S.D/S}$$

$$\text{LOQ} = 10 \times \text{S.D/S}$$

Where S is the slope of the calibration curve and SD is the standard deviation of response of least concentration of calibration curve in three replicates. Results of the study were shown in Table 5.

**Table 5: LOD and LOQ of Benidipine HCl and Telmisartan.**

S.No.	Drugs	LOD ( $\mu\text{g/mL}$ )	LOQ ( $\mu\text{g/mL}$ )
1	Benidipine HCl	0.1354	0.4103
2	Telmisartan	1.6823	5.0980

**Robustness:** Robustness of the method was determined by carrying out the analysis at three different wavelengths ( $\pm 2$  nm). The respective absorbance was noted and the result was indicated by % RSD. Results of the study were shown in Table 6.

**Table 6: Robustness studies data of the Proposed Method.**

S. No.	Conc. (µg/mL)	Benidipine HCl			Conc. (µg/mL)	Telmisartan		
		243 nm	245 nm	247 nm		243 nm	245 nm	247 nm
1	3	0.628	0.634	0.641	30	0.609	0.619	0.628
2	3	0.625	0.633	0.640	30	0.608	0.618	0.626
3	3	0.626	0.632	0.642	30	0.607	0.620	0.627
4	3	0.627	0.630	0.641	30	0.609	0.619	0.625
5	3	0.626	0.632	0.643	30	0.608	0.619	0.627
6	3	0.624	0.629	0.641	30	0.606	0.618	0.624
Mean		0.626	0.631	0.641		0.607	0.618	0.626
SD		0.0014	0.0018	0.0010		0.0011	0.0007	0.0014
%RSD		0.2259	0.2947	0.1610		0.1923	0.1216	0.2350

**Application of the proposed method to tablet dosage form:**

The proposed methods were applied to the quantification of Benidipine HCl and Telmisartan in tablet dosage forms. The results shown in Table 8,

suggest that the method is suitable for the determination of Benidipine HCl and Telmisartan with good accuracy and precision. The excipients in the dosage forms do not interfere in the assay procedure.

**Table 8: Assay results of Benidipine HCl and Telmisartan in tablet dosage form**

Formulation	Label Claim (mg/tablet)	Amount Found (mg/tablet)	% Assay
Benidin T	Benidipine- 4	3.93	98.2
	Telmisartan- 40	39.8	99.5

**CONCLUSION**

The developed method was found to be precise as the %RSD values for intra-day and inter-day were found to be less than 2%. Good recoveries of the drugs were obtained at each added concentration, which indicates that the method was accurate. The LOD and LOQ were found to be in microgram level, which indicates the sensitivity of the method. The method was also found to be robust as indicated by the %RSD values which are less than 2%. The results of assay shows that the amount of drugs was in good agreement with the label claim of the formulation as indicated by % assay. The proposed method also can be used for the routine quality control analysis of Benidipine HCl and Telmisartan in bulk and pharmaceutical formulations.

**ACKNOWLEDGEMENTS**

The authors are thankful to Management of Vignan Pharmacy College for providing necessary facilities for completion this research work and also thankful to Dr. Reddy's Lab Pvt. Limited, Hyderabad, for providing the gift sample of Telmisartan.

**REFERENCES**

- Japanese Pharmacopoeia, JP XVII, 475-476. Roja P, Eswarudu M.M, Ravi sankar P, Srinivasa Babu P. An Updated Review on Analytical Methods for Estimation of Azelnidipine and Telmisartan. Asian Journal of Pharmaceutical Research and development, 2022; 10(2): 59-76.
- Sohanabanu Malek, Laxman Prajapati, Amit Joshi, Mohammadali Kharodiya. Spectrophotometric Method for the Determination of Benidipine Hydrochloride in Pharmaceutical Formulation. Tropical Journal of Pharmaceutical and Life Sciences, 2018; 5(1): 01-07.
- Manish Kumar, Ajay Kumar Shukla, Ram Singh Bishnoi, C. P. Jain. Development of UV Spectrophotometric Method for the Determination of Benidipine Hydrochloride by using Quality by Design (QbD) Approach. International Journal of Applied Pharmaceutics, 2018; 10(4): 92-97.
- Ayca Karasakal. First Order Derivative Spectrophotometric Method for the Determination of Benidipine Hydrochloride Pharmaceutical Preparations and Forced Degradation Study. Optics and Spectroscopy, 2015; 118(6): 1002-1006.
- A. Karasakal G. Dogan; Y. Yalcin Gurkan. Molecules of Benidipine: experimental and theoretical investigation. Bulgarian Chemical Communications, 2017; 49(1): 133-139.
- Singhvi and S. C. Chaturvedi. Spectrophotometric methods for estimation of Benidipine Hydrochloride from Tablets, Indian Journal of Pharmaceutical Sciences, 1998; Mar-Apr: 113-114.
- Esen Bellur Atici, Bekir Karliga. Identification, Synthesis and characterization of process related impurities of Benidipine hydrochloride, Stress-testing/ Stability studies and HPLC/UPLC method validations. Journal of Pharmaceutical Analysis, 2015; 256-268.
- Nurgul Karadas, Senem Sanli, Mehmet Gumustas, Sibel A. Ozkan. Voltametric and RP-LC assay for determination of Benidipine HCL. Journal of Pharmaceutical and Biomedical Analysis, 2012; 116-125.
- Kishanta Kumar Pradhan, Uma Shankar Mishra, Aurobindo Sahoo, Kanhu Charana Sahu, Debananda Mishra, Ranjit Dash. Method development and validation of telmisartan in bulk and pharmaceutical dosage forms by UV spectrophotometric method. International Journal of Research in Pharmaceutical Sciences, 2011; 2(4): 526-530.
- Niranjan D. Chivate, Siddharth M. Patil, Jagdish K.

- Saboji, Anuradha N. Chivate. Development of UV Spectrophotometric method for estimation and validation of telmisartan as a pure API. *Journal of Pharmacy Research*, 2012; 5(6): 3331-3333.
10. Manish Kumar; Chinmoy Kumar; Shailendra Bhatt; A. Pandurangan; Vichitra Kaushik; Anuj Malik, Vipin Saini. Dissolution Method Development and Validation for Tablet Dosage form of Telmisartan Using Spectrophotometric Method. *Journal of Chemical and Pharmaceutical Research* .2018; 10(5): 148-156.
  11. Komal Patel, Komal Dhudasia, Amit Patel, Jayant Dave, Chaganbhai Patel. Stress degradation studies on Telmisartan and development of a validated method by UV spectrophotometry in bulk and pharmaceutical dosage forms. *Pharmaceutical Methods*, 2011; 2(4): 253-259.
  12. Ajit Pandey, H. Savarkar, Mukesh Singh, Dr. P Kashyap, Priyanka Ghosh. UV-Spectrophotometric Method for estimation of Telmisartan in Bulk and Tablet Dosage Form. *International Journal of Chem Tech Research*, 2011; 3(2): 657-660.
  13. M. V. Basaveswara Rao, A. V. D. Nagendra Kumar, M. Sivanadh, G. Venkata Rao. Validated RP-HPLC Method For the Estimation of Telmisartan in Tablet Formulation. *Bulletin of Pharmaceutical research*, 2012; 2(2): 50-55.
  14. Sujana K, Gowri Sankar D, Bala Souris O, Swathi Rani G. Stability Indicating RP-HPLC method for the determination of Telmisartan in Pure and Pharmaceutical Formulation. *International Journal of Pharmacy and Pharmaceutical Sciences*, 2011; 3(2): 164-167.
  15. V. Kabra, V. Agrahari, P. Trivedi. Development and Validation of a Reverse Phase Liquid Chromatographic Method for Quantitative Estimation of Telmisartan in Human Plasma. *International Conference on Biomedical Engineering*, 2009; 1297-1300.
  16. M. S. Palled, P.M.N. Rajesh, M. Chatter, A.R. Bhat. RP-HPLC Determination of Telmisartan in Tablet Dosage Forms, *Indian Journal of Pharmaceutical Sciences*, 2005; 108-110.
  17. Sharma Shaina, Soni Varinder, Rahar Sandeep, Bhatia Nitish. Development and Validation of Method for the Estimation of Telmisartan as API in Tablet Dosage form and Prepared Spherical Agglomerates by RP-HPLC, 2016; 4(1): 63-79.
  18. Phani Kishore Ch., Bhanu Prakash Reddy V, Dhanashri M Kale. Development and validation of stability indicating HPLC method for the estimation of Telmisartan related substances in tablets formulation. *International Journal of Pharmacy Research and Sciences*, 2010; 1(4): 493-501.
  19. Bhavna. A. Patel, Aanand Kumari. D. Captain. Development and Validation of RP-HPLC Method for Estimation of Telmisartan in Bulk and Formulation Using Fluorescence Detector. *Journal of Biomedical and Pharmaceutical Research*, 2014; 3(2): 44-48.
  20. Bhadoriya Upendra, Dhaked Hemant, Danodia Abhinandan Kumar. RP-HPLC Method Development and Validation for Estimation of Telmisartan in Bulk and Tablet Dosage Form. *International Journal of Drug Regulatory Affairs*, 2013; 1(2): 61-64.
  21. Khyati Patel, Darshil Shah, Dr. Dilip Maheshwari. Dual Wavelength Spectrophotometric Method for Estimation of Benidipine Hydrochloride and Telmisartan in Pharmaceutical Dosage Form. *World Journal of Pharmaceutical Research*, 2018; 7(5): 1494-1505.
  22. Vyas Apexaben Arvindbhai; Pragnesh Patani; Seju Patel. Stability Indicating HPLC method Development and Validation for the simultaneous Estimation of Benidipine HCL and Telmisartan in its Pharmaceutical Dosage form. *Pharma Science Monitor*, 2018; 9(2): 143-163.
  23. Majan Naim; Aejaz Ahmed; Khan GJ. Stability Indicating RP-HPLC Method Development and Validation for simultaneous Estimation of Telmisartan and Benidipine Hydrochloride in Pharmaceutical Dosage Form. *Asian Journal of Pharmaceutical and Clinical Research*, 2018; 11(5): 342-350.
  24. Varsha Chandgude, G. K. Dyade, R. B. Jadhav. Stability indicating Validated RP-HPLC method development for simultaneous estimation of Benidipine HCl and Telmisartan from Pharmaceutical Dosage form. *International Journal of Pharmaceutical Sciences and Research*, 2019; 8(11): 767-773.
  25. Eswarudu M.M, Venu Priya K, Chinna Eswaraiah M, Dipankar and B, Raja B. Validated RP-HPLC Method for Simultaneous estimation of Rosuvastatin and Telmisartan in bulk and pharmaceutical dosage form. *PHARMANEST, An International Journal of Advances in Pharmaceutical Sciences*, 2015; 6(1): 2648-2653.
  26. Q2A: Text on Validation of Analytical Procedures. In *International Conference on Harmonization. Federal Register*, 1997; 60(40): 11260-11262.
  27. Q2B: Text on; Validation of Analytical Procedures. In *International Conference on Harmonization. Federal Register*, 1997; 62(96): 27463-27467.