



EVALUATION OF CATALASE, PARAOXONASE1 ACTIVITIES IN TYPE 2 DIABETES AND THEIR ROLE IN GLYCEMIC CONTROL

Dr. SK. Deepthi^{1*}, Dr. J. N. Naidu², Dr. G. Amar Raghu Narayan³

¹Assistant Professor, Department of Biochemistry, Narayana Medical College.

²Professor, Department of Biochemistry, Narayana Medical College, Nellore.

³Professor, Department of Plastic Surgery, Narayana Medical College and Hospital, Nellore.

Article Received on 01/06/2015

Article Revised on 22/06/2015

Article Accepted on 13/07/2015

*Correspondence for

Author

Dr. SK. Deepthi

Assistant Professor,
Department of
Biochemistry, Narayana
Medical College, Nellore

ABSTRACT

Patients with type II diabetes mellitus (NIDDM) are more prone to ischemic heart disease (IHD). Free radical damage is mostly responsible for causing all complications in diabetic patients. Conflicting reports are available regarding the antioxidant status in patients of NIDDM complicated with IHD. This study was undertaken to investigate the oxidative status in patients of NIDDM and to assess their relation with plasma glucose, catalase and paraoxanase activity.

KEY WORDS: type II diabetes mellitus (NIDDM), paraoxanase1I (PON1), catalase (CAT)

INTRODUCTION

Diabetes mellitus is a chronic disorder resulting from a number of factors in which an absolute or relative deficiency of insulin or its function occurs. It is projected that by the year 2025, India alone would have 57 million diabetics mainly of type 2 DM constituting 90% of the diabetic population.^[1,2,3]

Oxygen free radicals contribute to the development of exacerbation of many mankind's most common illness, including heart attacks. The global burden of IHD is highlighted by the world health reports which estimates 58.6% of all deaths in 2014 were attributed to cardiovascular disease.^[4]

Catalase is a common enzyme found in nearly all living organisms exposed to oxygen. It catalyzes the decomposition of hydrogen peroxide to water and oxygen.^[5] It is a very

important enzyme in protecting the cell from oxidative damage by reactive oxygen species(ROS).

PON1 (paraoxonase-1) is an HDL (high density lipoprotein) – associated enzymes capable of hydrolysing diverse substrates from OP(organ phosphate) toxins to oxidized phospholipids. As such it has been linked with both the prevention of OP poisoning and inhibition of atherosclerosis initiated by oxidatively modified LDL9low density lipoprotein).^[6]

Paraoxanase is a glycoprotein synthesized in the liver and secreted into the blood, where it associates with the HDL and has been implicated in the detoxification of organophosphate and possibly in the prevention of LDL lipid peroxidation .^[7,8] by its paraoxonase activity and by preventing homocysteinylation of APO B¹⁰⁰.

Low serum paraoxonase activity has been associated with increased susceptibility to atherosclerosis which could be due to reduced capacity to detoxify lipid peroxides in diabetes. The aim of the study was to evaluate the catalase, PON1 activity in type 2 DM to know how they vary and how they are related to each other in diabetes.

MATERIAL METHOD

The study was conducted over a period of 6 months. The study includes 80 subjects recruited in medicine department in Narayana Medical college and Hospital out of 80, 50 were patients with type 2 DM and 30 were normal healthy subjects matched to the age and sex.

5ml blood samples were collected after 12 hours of fasting for estimation of fasting blood glucose and 2 ml blood is collected in a plain tube for serum paroxonase estimation. This procedure is carried out both in case and controls.

FBS was estimated by glucose oxidase peroxidase method.^[9]

Serum paraoxonase was estimated by spectrophotometric method^[10] estimation of catalase activity in RBC by hemolysate method.^[11]

RESULTS

The data analysis was done using SPSS software. The results were expressed as mean \pm standard deviation and range values.

The p value of 0.005 was considered for statistical significance.

The mean and standard deviation of all parameters of the study were calculated in patients and controls.

Table I shows the comparison of fasting blood sugar (FBS) in cases and controls.

Parameters	Cases	Controls	P value
FBS	160 ± 80	90.1 ± 14	0.001

Table II shows the comparison of paroxonase 1 (PON1) in cases and controls.

Parameters	Cases	Controls	P value
PON1	36.8 ± 11.2	58.6 ± 7.4	0.0001

Table III shows the comparison of catalase in cases and controls.

Parameters	Cases	Controls	P value
CAT	63 ± 19.7	165 ± 54	0.0001

Table IV shows CAT /PON1 ratio in cases and controls.

Parameters	Cases	Controls	P value
CAT / PON1`	2.72 ± 0.7	1.5 ± 0.36	<0.01

DISCUSSION

Hyperglycemia a hall mark of diabetic condition depletes natural antioxidants and facilitates the production of reactive oxygen species (ROS) which has the ability to react with all biological molecules like lipids, proteins, carbohydrates, DNA etc and exert cytotoxic effects on cellular components.^[12]

High levels of glucose are associated with non enzymatic glycation of both extra and intracellular proteins. The occurrence of free radical induced lipid peroxidation causes considerable changes in cell membrane.^[13] peroxidation of lipid membrane has been related to the pathogenesis of many degenerative diseases, such as atherosclerosis oxidative damage to DNA, ageing and DM^[14] in our study the mean values of FBS in cases was 160 ± 80 controls 90.1 ± 14 it was evident that the serum glucose levels were increased in type 2 DM patients when compared to healthy controls. P value 0.001 which was significant. Similarly mean values of PON1 activity in cases and controls were 36.8 ± 11.2 and 58.6 ± 7.4 respectively. It was evident that PON1 activity was decreased in cases when compared to controls and was significant (pvalue 0.0001)

Catalase is an antioxidant enzyme. It decomposes hydrogen peroxide. Which is a reactive oxygen species.

The activity of antioxidant enzymes in the course of diabetes depends on the glycemic control. In diabetes this antioxidant enzyme will be decreased because of poor glycemic control. The results of our study show that there was decrease in the activity of catalase in NIDDM.

Reduced CAT/ PON1 ratio can be used to assess poor glycemic control measurement of PON1 and CAT assess the oxidative stress in diabetes. Hence by using the above parameters we can access the oxidative stress and glycemic control in type 2 DM.

REFERENCES

1. H.Surekha Rani, G.Madhavi. Risk factors for coronary heart disease in type 2 diabetes. Indian Journal of clinical bicohemsitry 2005; 2012; 75-80.
2. King H, Aubert R.E., Herman W.H Global burden of diabetes, 1995 – 2025, Prevalence, numerical estimates and projection Diabetic care, 1998; 21: 1414 -1431.
3. Ramachandran A. Snehalatha C, Vishwanathan, Burden of type 2 diabetes and its complication. The Indian scenario Carr. Sci, 2002; 83: 1471-76.
4. Cardiovascular Health Research Initiative global burden of cardiovascular disease. South Asian Journal of preventive Cardiology, 2003; 7(2): 69-71.
5. Chelikani P, Fita I, Loewan PC, “Diversity of structures and properties among catalases”. Cell.Mol.Life Sci. January 2004; 61(2): 192-208. Doi:10.1007/S00018-003-3206-5(<http://dx.doi.org>).
6. Deakin SP, James RW. Genetic and environmental factors modulating serum concentration and activities fo the antioxidant enzyme paraoxonase-1 clin Sci (lond) 2004-Nov; 107(5): 435-47.
7. Dr.SK.Deepthi, Dr.G.Amar Raghu Narayan. Paroxanase acitivity and its concentration in type 2 diabetes mellitus. Int J Pharm Bio Sci 2012 Oct: 3(4).
8. Abott CA, Mackness serum paraoxonase activity, concentration and phenoty pe distribution in diabetes mellitus and its relationship to serum lipids and lipoproteins. Arterioscler Thromb. Vas.Biol 1995 Nov; 15(11): 1812-1818.
9. David B Sacks M.B, Ch.B.Determination of glucose in body fluids / 776. Carld. A Burtls Edward R. Ash wood, M.B, teitz text book of clinical biochemistry, 3rd ed, 1998; 778-780 philadelphia.
10. Sarkar P.D, Shivaprakash. Association between paraoxonase activity and lipid levels in patients with premature coronary artery disease. Clini Cachimica Acta. 2006; 373: 77-81.

11. Aebi.H.Catalase; In methods in enzymatic analysis. Ed-Bergmeyer HU. Vol 3 academic press, New York 276-286.
12. Dincer.Y.Akçay, T, Aldemir.Z and Likova. H. Effects of oxidative stress on glutathione pathway in red blood cells from patients with insulin dependent diabetes mellitus met. 2002; 51: 1360-1362.
13. Agarwal.S, Banerjee. S Effects of oxygen on ferrous sulphate induced lipids peroxidation in liposomal membrane. Ind.J.Biochem and Biophysics. 1995; 21: 331-334.
14. Chatterjee S.N, Agarwal.S and Amit kumar membrane lipid peroxidation and its pathological consequence. Ind J of Biochem and Biophysics, 1998; 25: 31.