

**ANTIMALARIAL ACTIVITY OF *PROSOPIS CINERARIA* (L) DRUCE AGAINST  
CHLOROQUINE SENSITIVE *PLASMODIUM FALCIPARUM* 3D7 STRAIN****P.V.V. Satish<sup>1</sup>, K. Somaiah<sup>2</sup>, P. Brahman<sup>3</sup>, N. Sree Rekha<sup>4</sup> and Dr. K. Sunita<sup>5\*</sup>**<sup>1,3,4,5</sup>Department of Zoology and Aquaculture, Acharya Nagarjuna University, Nagarjunanagar 522510, Guntur, Andhra Pradesh, India.<sup>2</sup>Department of Environmental Sciences, Acharya Nagarjuna University, Nagarjunanagar 522510, Guntur, Andhra Pradesh, India.**\*Correspondence for Author: Dr. K. Sunita**

Department of Zoology and Aquaculture, Acharya Nagarjuna University, Nagarjunanagar 522510, Guntur, Andhra Pradesh, India.

Article Received on 11/09/2015

Article Revised on 01/11/2015

Article Accepted on 21/11/2015

**ABSTRACT**

The present study aimed to evaluate the antiplasmodial activity of medicinal plant *Prosopis cineraria* against chloroquine (CQ)-sensitive *Plasmodium falciparum* 3D7 strain and cytotoxicity against THP-1 cell line. The plant *Prosopis cineraria* were collected from Acharya Nagarjuna University Campus, Nagarjunanagar, Guntur district, Andhra Pradesh, India. Crude extracts from dried leaves, stem, flowers and roots of *P. cineraria* were prepared through soxhlet extraction using methanol, chloroform, hexane, ethyl acetate and aqueous sequentially. These extracts were tested *in vitro* against laboratory adopted *P. falciparum* 3D7 strain. The crude extracts were also tested for their cytotoxicity against THP-1 cell line. The phytochemical screenings were also conducted with standard methods. The ethyl acetate extract of leaf, stem, flower and root; chloroform extract of root showed  $IC_{50}$  values from 5-50  $\mu$ g/mL with good antimalarial activity and were significant at  $P < 0.05$  and  $< 0.01$ . The chloroform extracts of leaf, stem, flower; the aqueous extracts of stem, flower and root showed  $IC_{50}$  values ranging from 50 to 100  $\mu$ g/mL which shown activity. Out of all the extracts, the ethyl acetate extract of flower ( $IC_{50} = 27.33 \mu$ g/mL) has shown excellent antimalarial activity. The methanol extracts of leaf, stem, flower and root; hexane extract of leaves, stem, flower and root; aqueous extract of leaves were inactive. All the extracts were non toxic to THP-1 cells. The phytochemical screening has revealed the presence of alkaloids, triterpenes, flavonoids, tannins, coumarins, carbohydrates, phenols, saponins, phlobatamins and steroids. This is the first report of antimalarial activity of *P. cineraria* and it is concluded that the ethyl acetate extract of flower is potent for the development of antimalarial drugs.

**KEYWORDS:** *Prosopis cineraria*, antiplasmodial activity,  $IC_{50}$ , cytotoxic activity, selectivity index.**INTRODUCTION**

Malaria is a curable, preventable and oldest recorded disease can be found even in ancient Indian medical literature like Charaka Samhita. The name malaria was originated from Italian words "mal" and "aria" which means bad air.<sup>[1]</sup> Malaria is very risky parasitic disease caused by protozoan parasites *Plasmodium vivax*, *Plasmodium falciparum*, *Plasmodium malariae* and *Plasmodium ovale* and the parasite has transmitted from human to human by the bite of infective *Anopheles* mosquito.<sup>[2]</sup> Each year 300 to 500 million new cases are being diagnosed and nearly 1.5 million people died; mainstream of deaths reported from Sub Saharan African countries, the majority of them were children under 5 years and pregnant women.<sup>[3]</sup> Malaria has an enormous impact on child health in malaria endemic countries and contributes to illness, respiratory infection, diarrhoeal disease and malnutrition.<sup>[4]</sup>

The prevalence of malaria increased in 1980s and 1990s as the parasites developed resistance to the most frequently used antimalarials and the vectors became resistance to insecticides.<sup>[5]</sup> The first effective drug is chloroquine and its resistance was reported in 1957, consequently distributed all over the world and reported from India in 1976.<sup>[6,7]</sup> Now artemisinin and its derivatives are used as first line treatment according to World Health Organization Proceedings of Malaria Treatment. Unfortunately artemisinin-resistant strains has been reported from Thai- Cambodia in 2009 and hasten the need for new antimalarials.<sup>[8]</sup>

Historically and traditionally plant parts have always been used as an important source in the medicine against malaria. Most effective antimalarial drugs such as chloroquine, quinine and artemisinin are derived from plants. The first effective malarial drug quinine was extracted from Cinchona tree; based on this structure chloroquine and primaquine were synthesized. The other

effective drug artemisinin was extracted from Chinese herbal tree *Artemisia annua* in 1972.<sup>[9]</sup> Artemisinin and its derivatives are now recommended by World Health Organization (WHO) worldwide, in combination with other drugs such as lumefantrine, amodiaquine, mefloquine, sulphadoxine-pyrimethamine (SP) as the first line treatment of malaria.<sup>[10]</sup> This fact has encouraged the continuing search for new natural product-derived antimalarial drugs. Several plants are used in traditional medicine for the treatment of malaria and fever in malaria endemic areas.<sup>[11]</sup>

The present plant *Prosopis cineraria* (L) Druce is a species of flowering tree in the pea family belonging to the family Mimosoideae and is a large shrub that grows up to 10 m height with prickly branches, prickles curved and compressed. It is indigenous to dry portions of Western Asia and the Indian subcontinent, including Afghanistan, Iran, India, Oman, Pakistan, Saudi Arabia, the United Arab Emirates, and Yemen. It is distributed all over the India particularly in the parts of Rajasthan, Gujarat, Haryana, Uttar Pradesh, Tamilnadu and Andhra Pradesh. It is adopted as a state tree of Rajasthan (India). A large and well known example of the species is the Tree of Life in Bahrain – approximately 400 years old and growing in a desert devoid of any obvious sources of water. This is also the national tree of the United Arab Emirates.<sup>[12,13]</sup> The present study evaluated the antiplasmodial activity of methanol (MeOH), chloroform (CHCl<sub>3</sub>), hexane (He), ethyl acetate (EtOAc) and aqueous (Aq) extracts from leaves, stem, roots and flowers of *Prosopis cineraria* (L) Druce.

## MATERIALS AND METHODS

### Plant Collection

Fresh samples of leaves, stem, roots and flowers from *Prosopis cineraria* (L) Druce were collected from ANU campus, Nagarjunanagar of Guntur district, Andhra Pradesh, India. The plant *P. cineraria* were deposited in the Department of Botany, Acharya Nagarjuna University and voucher specimen was deposited in the department. All the collected plant parts were washed thrice with tap water and twice with distilled water to remove the adhering salts and other associated animals. The authentication of the plant species were done by Prof. K. Khasim, Department of Botany, Acharya Nagarjuna University, Guntur, Andhra Pradesh, India.

### Extract preparation

Shade-dried plant samples were subjected for in 90% different organic solvents methanol, chloroform, hexane, ethyl acetate and water at 50-60°C in a Soxhlet apparatus. After complete extraction, the filtrates were concentrated separately by rotary vacuum evaporation (>45°C) and then freeze dried (-80°C) to obtain solid residue. The extraction percentage was calculated by using the following formula:

$$\text{Percentage of extraction} = \frac{\text{weight of the extract (g)}}{\text{weight of the plant material (g)}} \times 100$$

The extracts of plant were screened for the presence of phytochemical constituents by following the method of Sofowora (1982) and Kepam (1986).<sup>[14,15]</sup> The plant extracts were dissolved in dimethyl sulphoxide and filtered through millipore sterile filters (mesh 0.20 µm, Sartorius Stedim Biotech GmbH, Germany). The filtrate was used for testing at different concentrations of 200, 100, 50, 25 and 12.5 µg/mL.<sup>[16]</sup>

### Parasite cultivation

The antiplasmodial activity of plant extracts was screened against CQ-sensitive *P. falciparum* 3D7 strain obtained from ongoing cultures in the laboratory. They were cultured according to the method of Trager and Jensen (1976).<sup>[17]</sup> in candle jar dissector. *P. falciparum* were cultivated in human O<sup>Rh+</sup> red blood cells using RPMI 1640 medium (Sigma Laboratories Private Limited, Mumbai, India) supplemented with O<sup>Rh+</sup> serum (10%), 5% sodium bicarbonate and 50 µg/mL of gentamycin sulfate. Hematocrits were adjusted at 2% and cultures of parasite were used when they exhibited 2% parasitemia.<sup>[18]</sup>

### In vitro antimalarial screening

The *P. falciparum* malaria parasite culture suspension of 3D7 (synchronized with 5% sorbitol to ring stage) was seeded (200 µL/well) in 96 well tissue culture plates. Plant extracts were added in µL/well to get different concentrations of extract (200, 100, 50, 25, 12.5 µg/mL). Chloroquine treated parasites were kept as positive controls and DMSO treated parasites were kept as negative control. The parasites were cultured for 30 h in candle jar desiccator. The cultures were incubated at 37°C for 48 hours in an atmosphere of 2% O<sub>2</sub>, 5% CO<sub>2</sub> and 93% N<sub>2</sub>. 18 h before termination of the assay [<sup>3</sup>H] Hypoxanthine (0.5 µCi/well) was added to each well of 96 well plate. The effects of extracts in the cultures were evaluated by the measurement of [<sup>3</sup>H] Hypoxanthine incorporation into the parasite nucleic acids.<sup>[19]</sup> Each treatment has four replicates; at end of experiment one set of the pRBC cells collected from wells and smears were prepared. These smears were fixed in methanol and air dried. The smears were stained with Acridine Orange (AO) stain. Stained smears were observed under UV illumination microscope (Carl Zeiss - Germany) for confirmation of [<sup>3</sup>H] Hypoxanthine assay data, remaining other three replicates were used for [<sup>3</sup>H] Hypoxanthine assay. The experiment was terminated and the cultures were frozen and stored in -20°C. The parasites were harvested on glass filter papers using NUNC Cell Harvester and CPM counts were recorded in gamma scintillation counter. Control readings were considered as 100% parasite growth and calculated the parasite inhibition in plant extract treated wells. The parasite inhibition was calculated as follows:

$$\% \text{ Inhibition} = \frac{\text{Average CPM of Control} - \text{Average CPM of plant extracts}}{\text{Average CPM of Control}}$$

The IC<sub>50</sub> values were determined by plotting concentration of extract on X-axis and percentage of inhibition on Y-axis with dose-response curves using Minitab 11. 12. 32. Bit software.

#### Cytotoxicity of extracts on THP-1 monocyte cells

The assays were carrying out using 96-well flat-bottom tissue-culture plates. Cytotoxic properties of active plant extracts were assessed by functional assay<sup>[20]</sup> using THP-1 cells. The cells were cultured in RPMI-1640 medium which contained 10% fetal bovine serum, 0.21% sodium bicarbonate (Sigma), and 100 µg/mL penicillin and 50µg/mL gentamicin (complete medium). Briefly, cells (0.2×10<sup>6</sup> cells/200µL/well) were seeded into 96-well flat-bottom tissue-culture plates in complete medium. Drug solutions (200, 100, 50, 25 and 12.5µg/mL) were added after 24 h of seeding and incubated for 48 hours in a humidified atmosphere at 37°C and 5% CO<sub>2</sub>. DMSO as negative inhibitor ellipticine as a positive inhibitor was added to each well. At end of experiment ten micro liters of a stock solution of 3-(4,5- dimethylthiazol-2-yl)-2,5-diphenyltetrazolium bromide (MTT; 5µg/ml in 1x phosphate-buffered saline) was added to each well, gently mixed, and incubated for another 4 h. After

spinning the plate was centrifuged at 1,500 RPM for 5 min, the supernatant was discarded, subsequently added 100µL of DMSO (stopping agent). After formation of formazon was read on a micro titer plate reader (Versa max tunable multiwell plate reader) at 570 nm and the percentage of cell viability calculated using the following formula.<sup>[21]</sup> The selectivity index of *in vitro* toxicity was calculated for each extract as the IC<sub>50</sub> for THP-1 cells/ IC<sub>50</sub> for *P. falciparum*.

$$\% \text{ Cell viability} = \frac{\text{Mean absorbance in test wells}}{\text{Mean absorbance in control wells}} \times 100$$

The IC<sub>50</sub> values were determinates by plotting concentration of extract on X- axis and percentage of cell viability on Y- axis with dose response curves using Minitab 11. 12. 32. Bit software.

#### RESULTS

The Phytochemical studies revealed that the methanol, chloroform, hexane, ethyl acetate and aqueous extracts of leaf, stem, flower and root of *P. cineraria* have variety of Phytochemical constituents namely alkaloids, triterpenes, flavonoids, tannins, coumarins, carbohydrates, phenols, saponins, phlobatamins and steroids represented in Table 1.

**Table 1: Preliminary phytochemical evaluation of *Prosopis cineraria* in different extracts of leaves, stem, flowers and roots.**

Tested compounds <sup>2</sup>	Leaves					Bark					Roots					Flowers				
	Me	Ch	He	EtAc	Aqs	Me	Ch	He	EtAc	Aqs	Me	Ch	He	EtAc	Aqs	Me	Ch	He	EtAc	Aqs
Alkaloids	+	+	+	+	-	+	+	-	+	-	+	+	+	-	-	+	-	-	+	-
Coumarins	-	-	-	-	+	-	-	-	+	-	-	-	-	+	+	+	-	-	-	-
Carbohydrates	-	-	-	+	+	-	-	-	-	-	-	-	-	-	-	-	-	-	+	+
Phenols	+	-	+	-	-	+	-	-	-	-	-	-	+	+	-	+	-	+	-	-
Saponins	-	-	-	+	+	+	-	-	+	-	+	-	-	+	-	-	-	-	-	+
Tannins	+	-	+	+	-	+	-	+	+	-	+	-	+	-	-	-	-	-	-	-
Flavanoids	-	-	-	-	+	-	-	-	-	+	-	-	-	-	+	-	-	-	-	+
Terpenoids	-	-	-	+	-	+	+	-	-	-	+	-	-	+	-	-	+	-	-	-
Phlobatamins	+	-	-	+	-	-	-	-	-	+	-	-	-	-	-	-	+	-	-	-
Steroids	+	-	+	-	-	+	+	+	-	-	-	+	+	+	-	-	-	-	-	-

Me= Methanol; Ch= Chloroform; He= Hexane; EtAc= Ethyl Acetate; Aqs= Aqueous

+ present, - absent.

In the present study, crude extracts of methanol, hexane, chloroform, ethyl acetate and aqueous from leaves, stem, flowers and roots of *P. cineraria* were evaluated for their antimalarial potencies. This is the first report of antiplasmodial activity of experimental plant *P. cineraria* against *P. falciparum*. The  $IC_{50}$  values of the tested plant extracts against *P. falciparum* are listed in Table 2. The *in vitro* antiplasmodial activity of biological active substances is categorized into four groups basing on  $IC_{50}$  value ( $<5\mu\text{g/mL}$ -very active,  $<50\mu\text{g/mL}$ -active,  $50-100\mu\text{g/mL}$ - weakly active,  $>100\mu\text{g/mL}$  inactive).<sup>[22]</sup> Based on this categorization, the  $IC_{50}$  values of the ethyl acetate, aqueous and chloroform extracts of leaves, stem, roots and flowers of *P. cineraria* showed a range of inhibitory concentrations against CQ-sensitive *P. falciparum* 3D7 strain. The ethyl acetate extract of leaves ( $30.67\mu\text{g/mL}$ ), stem ( $36.00\mu\text{g/mL}$ ), flowers ( $27.33\mu\text{g/mL}$ ) and roots ( $41.00\mu\text{g/mL}$ ); and chloroform extract of root ( $40.33\mu\text{g/mL}$ ) showed good antimalarial activity and are significant at  $P<0.05$  and  $P<0.01$ . The aqueous extract of flowers ( $55.37\mu\text{g/mL}$ ) shown moderate antimalarial activity. Among these extracts, the

ethyl acetate extract of flowers showed excellent antimalarial activity ( $IC_{50}=27.33\mu\text{g/mL}$ ). The chloroform extracts of leaves, stem, flowers; the aqueous extracts of stem and roots showed  $IC_{50}$  values greater than  $50\mu\text{g/mL}$ . The methanol extracts of leaves, stem, flower and hexane extract of leaves and stem shown inactive antimalarial activity with  $IC_{50}$  values  $>100\mu\text{g/mL}$ . Moreover  $IC_{50}$  values of the aqueous extract of leaves, hexane extracts flowers and roots, methanol extract of roots were not determinate because of their unclear inhibition.


The *in vitro* cytotoxicity studies against THP-1 cell line were conducted for all the extracts. Out 20 extracts 19 showed  $IC_{50}$  value  $>200\mu\text{g/mL}$  whereas only chloroform extract of flower shown  $IC_{50}$  value  $54.64\mu\text{g/mL}$ . Finally, all the extracts showed  $IC_{50}$  values  $>200\mu\text{g/mL}$ . An extract is classified as non toxic when the  $IC_{50}$  value is  $>20\mu\text{g/mL}$ . Based on this, all the plant extracts are not harmful to *in vivo* studies. The selectivity indices indicate the low toxicity of tested extracts and safer for therapies (Table 2).

**Table 2: Antiplasmodial activity against CQ-sensitive *P. falciparum* 3D7 strain and cytotoxicity against THP-1 cell line of different crude extracts from *Prosopis cineraria*.**


Plant parts	Crude Extracts	% of Yield	$IC_{50}$ 3D7 strain ( $\mu\text{g/mL}$ )	$IC_{50}$ THP-1 cells ( $\mu\text{g/mL}$ )	SI
Leaves	Methanol	2.54	$145.33\pm11.37$ (117.08-173.58)	$>200$	$>1.37$
	Chloroform	10.82	$59.00\pm7.94$ (39.28-78.72)	$>200$	$>3.38$
	Hexane	7.08	$175.33\pm2.08$ (170.16-180.50)	$>200$	$>1.14$
	Ethyl Acetate	1.90	$30.67\pm1.15$ (27.80-33.54)	$>200$	$>6.52$
	Aqueous	4.42	$>200$	$>200$	$>1.00$
Stem	Methanol	3.84	$146.00\pm4.58$ (134.62-157.38)	$>200$	$>1.36$
	Chloroform	6.94	$67.67\pm2.52$ (61.42-73.92)	$>200$	$>2.95$
	Hexane	3.02	$>200$	$>200$	$>1.00$
	Ethyl Acetate	8.56	$36.00\pm1.00$ (33.52-38.48)	$>200$	$>5.55$
	Aqueous	4.68	$61.67\pm1.53$ (57.87-65.46)	$>200$	$>3.24$
Flower	Methanol	2.20	$>200$	$>200$	$>1.00$
	Chloroform	8.42	$76.67\pm3.06$ (69.08-84.26)	$54.67\pm4.16$ (44.32-65.01)	0.71
	Hexane	1.16	ND	$>200$	ND
	Ethyl Acetate	2.66	<b><math>27.33\pm4.93</math></b> (15.08-35.12)	$>200$	$>7.31$
	Aqueous	2.90	$55.37\pm3.79$ (46.26-65.07)	$>200$	$>3.61$
Root	Methanol	4.20	ND	$>200$	ND
	Chloroform	6.30	$40.33\pm1.53$ (36.54-44.13)	$>200$	$>4.95$
	Hexane	2.62	ND	$>200$	ND
	Ethyl Acetate	1.72	$41.00\pm2.65$ (34.45-47.57)	$>200$	$>4.87$
	Aqueous	4.42	$71.00\pm3.61$ (62.04-79.96)	$>200$	$>2.81$



Values are represented as mean±standard deviation, at 95% confidence intervals with upper and lower limits. ND- not determinate, SI- selectivity index,  $SI_{Plasmodium} = IC_{50} \text{ THP-1} / IC_{50} \text{ P. falciparum 3D7}$



Control



Flower extract treated

**Fig 1: Micrographs of synchronized ring stage *P.f3D7* culture treated with ethyl acetate flower extract (200 mg/μL) of *Prosopis cineraria* for 48 h showing inhibition of ring stages.**

## DISCUSSION

Malaria is still the most dangerous parasitic infectious disease which causes two million deaths every year. It is a great burden to developing nations, a number that could rise due to the increasing multi-drug resistance to all antimalarial drugs currently available.<sup>[23]</sup> There are several genetic polymorphisms identified in *P. falciparum* and *P. vivax* that can be providing reliable data about the prevalence of drug resistance. Amongst all, the *pfcr*, *pfmdr1*, *pfdhfr* and *pfdhps* associated with drug sensitivity, have great role in drug resistance mechanisms in parasites and is directly connected to treatment failure.<sup>[24]</sup>

From the past 20 years, many strains of *P. falciparum* have become resistant to chloroquine and other anti-malarial drugs. The development and spread of drug resistant strains of *P. falciparum* has limited effectiveness to the currently used malarial drugs. In view of this fact, the emergence and spread of parasites resisting to antimalarial drugs has caused an urgent need for novel effective alternative antimalarial drug compounds to be discovered and developed with minimal side effects.<sup>[25]</sup>

Plants have over the years, proved to be a good source of chemotherapeutic agents. Today, many of the drugs have been derived from plants resources such as quinine, chloroquine and artemisinin. Historically, medicinal plants have provided a source of inspiration for novel therapeutic drugs, as plant-derived medicines have made large contributions. According to WHO, now a days, 80% of the world's population rely on plants for their primary health care.<sup>[26]</sup> Plants are producing secondary metabolites for their defense, which play an important role of physiological activities in human body.<sup>[27]</sup> The medicinal value of plants is due to the substances that it contains, which produce a physiological action on the human body. Some examples of these plants are alkaloids, essential oils, tannins, resins and many others.<sup>[28]</sup> India had remarkable biodiversity and rich cultural traditions of plant use. Interestingly, today many of the pharmaceutical companies are utilizing such plant-based formulations in treatment of various diseases and disorders worldwide.<sup>[29]</sup>

The present investigation has evaluated the *in vitro* antiplasmodial activity of *Prosopis cineraria* with different extracts such as methanol, chloroform, hexane, ethyl acetate and aqueous forms of leaves, stem, roots and flowers. According to Rasoanaivo *et al.* (1992),<sup>[22]</sup> the *in vitro* antiplasmodial  $IC_{50}$  results classifies the biological active substances into four groups (<5μg/mL-very active, <50μg/mL-active, 50-100μg/mL-weakly active, >100μg/mL-inactive). Based on the hypothesis, out of the 20 extracts tested, five showed good ( $IC_{50}$  = 29-47.12μg/mL), six exhibited moderate ( $IC_{50}$  = 54-83.98μg/mL), while five displayed mild ( $IC_{50}$  = >100μg/mL) antiplasmodial activity as shown in Table 1. Remaining four extracts does not show any inhibition, due to this reason  $IC_{50}$  was not calculated. This is the first report of antiplasmodial activity for the experimental plant against *P. falciparum*. But the antimalarial activity of same genus *Prosopis juliflora* reported to have antimalarial activity against CQ-sensitive strain of *P. falciparum* (3D7). Ramajani reported the activity of the same plant against both CQ-resistant (K1) and CQ-sensitive (CY27) strains along with *P. berghii* with the dichloromethane extract and water-soluble extract. The dichloromethane extract of *P. juliflora* was significantly active under *in vitro* condition against both sensitive and resistant strains of *P. falciparum* and inhibited parasitemia of *P. berghei* infection in BALB/c mice when compared to untreated, implying direct parasitocidal activity. The water-soluble extract of *P. juliflora* was active *in vitro* against both sensitive and resistant *P. falciparum* strains, but no *in vivo* activity against *P. berghei*.<sup>[30]</sup>

The *P. cineraria* has been used in traditional medicine for treatment of leprosy, dysentery, bronchitis, asthma, leucoderma, muscular tremors and rheumatism. It has

been proved to possess anthelmintic, antibacterial, antifungal, antiviral and anticancer activities.<sup>[31]</sup> *Prosopis* spices have also been used in indigenous system of medicine as a folk medicine for various ailments. The bark is dry, acrid, bitter with sharp taste; antihelmintic tonic; cures leprosy, dysentery, asthma, leucoderma, piles, tremors of the muscle and wandering of the mind. The flowers are pounded mixed with sugar and used during pregnancy as safeguard against miscarriage. The ashes of the bark are rubbed over the skin to remove hair. The smoke of the leaves is good for eye troubles. Fresh leaf juice mixed with lemon juice is used for dyspepsia; extract of crushed pods is used for ear ache, tooth ache, pain relief from fractured bones. Aqueous extract of bark and leaves applied externally to treat skin disease disinfests wounds and promotes healing.<sup>[32]</sup>

The *in vitro* antiplasmodial activity of the *P. cineraria* ethyl acetate flower extract may be due to the presence of major chemical classes such as phenols and alkaloids. Hence alkaloids are strong antiplasmodial compounds. Except the alkaloids, the major chemical classes such as coumarins, phenols, polysaccharides and flavanoids also exerted strong antiplasmodial activities.<sup>[33]</sup> Bandaranayake (2002)<sup>[34]</sup> reported about the bioactive compounds and chemical constituents of mangrove plants.

Some of the traditional medicine involves the use of crude plant extracts which may contain an extensive diversity of molecules, often with indefinite biological effects. However, most of the available information regarding the medicinal potency of these plants is not provided with credible scientific data. For this reason, several researches have been reported the toxicity of medicinal plants.<sup>[35]</sup> In the present study, the *in vitro* cytotoxic effect against THP-1 cell lines to 20 extracts revealed that 19 showed  $IC_{50} > 200 \mu\text{g/mL}$  but chloroform extract of flower shown  $IC_{50}$  value of  $54.64 \mu\text{g/mL}$ . According to Falade et al., the cytotoxicity  $> 20$  considered as non toxic to animals and safer for further studies. Thus, all the plant extracts are not harmful and safer for therapies.<sup>[36]</sup>

Similarly, Bhagavan et al. reported the antimalarial activity of methanol, hexane, chloroform, acetone, and ethyl acetate extracts of *Citrus sinensis* (peel), *Leucas aspera*, *Phyllanthus acidus* (leaf), *Ocimum sanctum*, *Terminalia chebula* (seed) against CQ-sensitive (3D7) strain of *P. falciparum* and tested for their cytotoxicity on human laryngeal cancer cell line (HEp-2) and normal cell line (Vero).<sup>[37]</sup> Out of the 25 test extracts, they have concluded the leaf methanol and ethyl acetate extracts of *L. aspera*; ethyl acetate, methanol and acetone extracts of *P. acidus*; and seed acetone extract of *T. chebula* had good antiplasmodial activity ( $IC_{50}$  ranged from 4.76 -  $22.76 \mu\text{g/mL}$ ) with selectivity indices ranged from 2.04-9.97 for HEp-2 and  $> 2$  to  $< 12$  for Vero cells, respectively.

The mechanism of action might be due the inhibition of hemozoin biocrystallization by the alkaloids and inhibition of protein synthesis by triterpenoids (Samuel et al., 2012).<sup>[38]</sup> Additional *in vitro* and *in vivo* work aimed at understanding the mechanisms of action of the active plant extracts, isolating and characterizing the bioactive constituents is underway in our laboratories and will be reported in due course of time.

## CONCLUSION

In conclusion our shows that the ethyl acetate extract of flower of *Prosopis cineraria* exhibits good *in vitro* antiplasmodial activity against CQ-resistant strain of *P. falciparum*. Further evaluation of the extract may provide potential molecule for therapy of malaria.

## ACKNOWLEDGEMENT

The authors are thankful to the Co-ordinator, Department of Zoology and Aquaculture, Acharya Nagarjuna University for providing laboratory facilities.

## REFERENCES

1. Guna Rajan K, Balakrishna, Vijayan, Raja S. Evaluation of larvicidal activity of Pongamia pinnata extracts against three mosquito vectors. Asian Pac J Trop Biomed, 2013; 3(11): 853–858.
2. John MO, Christopher CK, Tom W, Collins O, Richard O, Zachary L, Daniel O, Jamie L, Stephen M, George A, Alloys S, John M, Sandra S, Richard D, Douglas JP. Parasitemia anemia and malarial anemia in infants and young children in a rural holoendemic Plasmodium falciparum transmission area. Am J Trop Med Hyg, 2006; 74(3): 376–385.
3. Tanner M, de Savigny D. Malaria eradication back on the table. Bull World Health Organ, 2008; 86: 82.
4. WHO. The World Health Report. Rolling Back malaria, Making a difference, 1999; 49–61.
5. Le Bars J, Durand R. The mechanisms of resistance to antimalarial drugs in Plasmodium falciparum. Fundam Clin Pharmacol, 2003; 17: 147–153.
6. Serena C, Paige ES, Jacques P, Sebastian E, Matthew B, Michael C, David C, Young TC, Mark EH, William A, Julia K, Karine GR. High content live cell imaging for the discovery of new antimalarial marine natural products. BMC Infect Dis, 2012; 12: 1–9.
7. Anupkumar R, Bhawna S, Sharma SK, Ghosh SK, Bhatt RM, Ashwani K, Mohanty SS, Pillai CR, Dash AP, Neena V. In vitro assessment of drug resistance in Plasmodium falciparum in five States of India. Indian J Med Res, 2012; 135: 494–499.
8. Noedl H, Youry Se, Schaecher K, Smith BL, Sochet D, Fukuda MM. Evidence of artemisinin-resistant malaria in Western Cambodia. N Engl J Med, 2008; 359: 2619–2620.
9. White NJ. Qinghaosu (Artemisinin): The price success. Science, 2008; 320: 2619–2620.
10. Mutabingwa TK. Artemisinin –based combination therapies (ACTs): best hope for malaria treatment

- but inaccessible to the needy! *Acta Trop*, 2005; 95: 305–315.
11. Tran QL, Tezuka Y, Ueda JY, Nguyen NT, Maruyama Y, Begum K, Kim HS, Wataya Y, Tran QK, Kadots S. In vitro antiplasmodial activity of antimalarial medicinal plants used in Vietnamese traditional medicine. *J. Ethnopharmacol*, 2003; 86: 249–252.
  12. Stella R, Narayanan N, Raj KB. Antitumour activity of *Prosopis cineraria* (L.) Druce against Ehrlich ascites carcinoma-induced mice. *Nat Pro Res*, 2011; 25(8): 857–862.
  13. Preeti K, Avatar SR, Mala A. Pharmacology, phytochemistry and therapeutic application of *Prosopis cineraria* (Linn): A review. *J of Plt Sci*, 2015; 3(1-1): 33-39.
  14. Sofowora A. Medicinal plants and traditional medicine in Africa. John Wiley and Sons, New York, 1982; 251.
  15. Kepam W. Qualitative organic analysis (Spectrochemical techniques). EdII. Mc Graw Hill, London, 1986; 40-58.
  16. Ouattara Y, Sanon S, Traore Y, Mahiou V, Azas N, Sawadogo L. Antimalarial activity of *Swartzia madagascariensis* Desv. (leguminosae), *Combretum glutinosum* Guill. and Perr. (Combretaceae) and *Tinospora bakis* Miers. (menispermaceae), Burkina Faso medicinal plants. *Afr J Tradit Complement Altern Med*, 2006; 3(1): 75–81.
  17. Trager W, Jensen JB. Human malaria parasites in continuous culture. *Science*, 1976; 193: 673–675.
  18. Trager W. The cultivation of *Plasmodium falciparum*: applications in basic and applied research in malaria. *Ann Trop Med Parasitol*, 1987; 81: 511–529.
  19. Simonsen HT, Jesper BN, Ulla WS, Ulf N, Pushpagadan P, Prabhakar J. In vitro screening of Indian medicinal plants for antiplasmodial activity. *J Ethnopharmacol*, 2001; 74: 195–204.
  20. Basim MA, Abdalla AA, Faris DM. In vitro inhibition of human leukemia THP-1 cells by *Origanum syriacum* L. and *Thymus vulgaris* L. extracts. *BMC Research Notes*, 2014; 7: 612–618.
  21. Khonkarn R, Okonogi S, Ampasavate C, Anuchapreeda S. Investigation of fruit peel extracts as sources for compounds with antioxidant and antiproliferative activities against human cell lines. *Food Chem Toxicol*, 2010; 48: 2122–2129.
  22. Rasoanaivo P, Ratsimamanga, Urverg S, Ramanitrhasimbola D, Rafatro H, Rakoto Ratsimamanga A. Criblage d'extraits de plantes de Madagascar pour recherche d'activité antipaludique et d'effet potentialisateur de la chloroquine. *J. Ethnopharmacol*, 1992; 64: 117–126.
  23. Ravikumar S, Samuel JI, Palavesam S, Murugesan G. In vitro antiplasmodial activity of ethanolic extracts of mangrove plants from South East coast of India against chloroquine-sensitive *Plasmodium falciparum*. *Parasitol Res*, 2011; 108: 873–878.
  24. Jovel IT, Rosa EM, Engels B, Rita P, Jackeline A, Gustavo F, Pedro EF, Maria IV, Irma GE, Anders B, Johan U 2011. Drug resistance associated genetic polymorphisms in *Plasmodium falciparum* and *Plasmodium vivax* collected in Honduras, Central America. *Malar J*, 2011; 10: 376.
  25. Pascual A, Philippe P, Françoise BV, Fabrice S, Denis M, Stéphane P, Pascal D, Didier B, Danièle M, Bernard F, Guillaume M, Nathalie B, Claude O, Eric D, Christophe R, Bruno P. Ex vivo activity of the ACT new components pyronaridine and piperaquine in comparison with conventional ACT drugs against isolates of *Plasmodium falciparum*. *Malar J*, 2012; 11: 45. doi: 10.1186/1475-2875-11-45.
  26. Okigbo RN, Ramesh P. Effects of plants and medicinal plant combinations as anti-infectives. *Af J Pharmacy and Pharmacol*, 2008; 2(7): 130–135.
  27. Vieira D, Padoani D, Soares J, Adriano J, Filho VC, de Souza MM, Couto AG 2012. Development of hydroethanolic extract of *Ipomoea pes-caprae* using factorial design followed by antinociceptive and anti-inflammatory evaluation. *Bra J of Pharmacog*, 2012; 23(1): 72–78. doi.org/10.1590/S0102-695X2012005000126.
  28. Lilybeth FO, Olga MN. Brine shrimp lethality assay of the ethanolic extracts of three selected species of medicinal plants from Iligan city, Philippines. *Int Res J Biological Sciences*, 2013; 2(11): 74–77.
  29. Johnson M, Babu A, Irudayaraj V. Antibacterial studies on in vitro derived calli of *Ocimum basilicum* L. *J Chem Pharm Res*, 2011; 2(2): 298–305.
  30. Ramazani A, Sardari S, Zakeri S, Vaziri B. In vitro antiplasmodial and phytochemical study of five *Artemisia* species from Iran and in vivo activity of two species. *Parasitol Res*, 2010; 107: 593–599.
  31. Malik A, Kalidhar SB. Phytochemical examination of *Prosopis cineraria* L. (Druce) leaves. *Indian J Pharm Sci.*, 2007; 69(4): 576–578.
  32. Akash G, Sanjeev KM. Review on *Prosopis cineraria*: A potential herb of Thar Desert. *Drug i n v en today*, 2015; 5: 60–65.
  33. Sherman PW, Billing J. Darwinian gastronomy: why we use spices. *Bioscience*, 1999; 49: 453–463.
  34. Bandaranayake WM. Bioactivities, bioactive compounds and chemical constituents of mangrove plants. *Wetlands Ecol Manag*, 2002; 10: 421–452.
  35. Lilybeth FO, Olga MN. Brine shrimp lethality assay of the ethanolic extracts of three selected species of medicinal plants from Iligan city, Philippines. *Int Res J Biological Sciences*, 2013; 2(11): 74–77.
  36. Flade MO, Akinboye DO, Gbotosho GO, Ajaiyeoba EO, Hapji TC, Abiodun OO, Oduola AM. In vitro and In vivo antimalarial activity of *Ficus thonningii* Blume (Moraceae) and *Lophira alata* Banks (Ochnaceae), Identified from the Ethnomedicine of the Nigerian Middle Belt. *J Parasitol Res.*, 2014; 1–6.



37. Bhagavan A, Rahuman AA, Kamaraj C, Kaushik NK, Mohanakrishna D, Sahal D. Antiplasmodial activity of botanical extracts against *Plasmodium falciparum*. *Parasitol Res*, 2011; 108(5): 1099-1109.
38. Samuel JI, Ravikumar S, Suganthi P. In vitro antiplasmodial effect of ethanolic extracts of coastal medicinal plants along Palak Strait against *Plasmodium falciparum*. *Asian Pac J Biomed*, 2012; 2(5): 364–367.