


**FORMULATION AND EVALUTION OF TOPICAL ANTIFUNGAL GEL CONTAINING
FLUCONAZOLE**
Gauri Ashok Phadtare*, Mukesh Tatyarao Mohite and Madhvi Anil Kuchekar

Dept. of Quality Assurance Techniques, Padmashree Dr. D.Y. Patil College of Pharmacy, Akurdi, Pune, 411044.

***Corresponding Author: Gauri Ashok Phadtare**

Dept. of Quality Assurance Techniques, Padmashree Dr. D.Y. Patil College of Pharmacy, Akurdi, Pune, 411044.

Article Received on 05/05/2016
Article Revised on 25/05/2016
Article Accepted on 15/06/2016
ABSTRACT

The present research has been undertaken with the aim to develop a topical gel formulation of Fluconazole. Fluconazole is an imidazole derivative and used for the treatment of local and systemic fungal infection. The oral use of Fluconazole is not much recommended as it has many side effects. Commercially Fluconazole topical gel preparation are not available in the market, thus this formulation is made for better patient compliance and to reduce the dose of drug and to avoid the side effects like liver damage and kidney damage. The gel was formulated by changing the polymer ratio. Formulation were evaluated for % yield, spreadability, extrudability, washability and viscosity in-vitro drug release study, skin irritation study, stability testing and finally it was concluded that formulation F3 was the best formulation among these formulation.

KEYWORDS: Fluconazole, Carbopol 934p, HPMC.

INTRODUCTION

Fungal infection of skin is now-a-days one of the common dermatological problem. The physicians have a wide choice for treatment from solid dosage to semisolid dosage form and to liquid dosage formulation. Among the topical formulation clear transparent gels have widely accepted in both cosmetics and pharmaceuticals.^[1] Topical treatment of dermatological disease as well as skin care, a wide variety of vehicle ranging from solids to semisolids and liquids preparations is available to clinicians and patients. Within the major group of semisolid preparations, the use of transparent gels has expanded both in cosmetics and in pharmaceutical preparation.^[2] For many decades treatment of an acute disease or a chronic illness has been mostly accomplished by delivery of drugs to patients using various pharmaceutical dosage forms, including tablets, capsules, pills, suppositories, cream, gel, ointments, liquids, aerosols and injectables, as drug carriers. Delivery of drugs to the skin is an effective and targeted therapy for local dermatological disorders. This route of drug delivery has gained popularity because it avoids first pass effects, gastrointestinal irritation, and metabolic degradation associated with oral administration. Due to the first pass effect only 25-45% of the orally administered dose reaches the blood circulation. In order to bypass these disadvantages the gel formulations have been proposed as topical application. Gels are defined as "semisolid system in which a liquid phase is constrained within a polymeric matrix in which a high degree of

physical and chemical cross-linking introduced". Fluconazole is a synthetic antifungal agent of the imidazole class; it works by slowing the growth of fungi that cause infection. It is used to treat fungal infection. Triazole drug targets the fungal-specific synthesis of membrane lipids. Fluconazole inserts preferentially into fungal membranes and disrupts their function. 5-fluorocytosine targets fungal specific DNA replication.^[3] Hydroxypropyl methylcellulose (HPMC), Carbopol 934p, has been used as hydrophilic polymers topically in gel drug delivery system.^[4]

MATERIALS AND METHODS^[5,6]
Material

Fluconazole, HPMC, carbopol934, trimethanolamine, glycerine, methyl paraben, propyl paraben, water.

Method

Polymer (like Carbopol 934p or HPMC) and purified water were taken in a beaker and allowed to soak for 24 hrs. To this required amount of drug (2 gm) was dispersed in water and then Carbopol 934p or HPMC was then neutralized with sufficient quantity of Triethanolamine. Glycerine as moistening agent, methyl paraben and Propyl paraben as preservatives were added slowly with continuous gently stirring until the homogenous gel was formed. Gel formulations of Fluconazole were prepared using different concentrations of carbopol934, HPMC.

(Table No.1) Optimized formulae of Fluconazole gel

Formulation code	Ingredients								
	Drug (gm)	Carbopol (gm)	HPMC (gm)	Water (ml)	Alcohol (ml)	Methyl Paraben (gm)	Propyl paraben (gm)	Glycerine (ml)	Triethanol amine (ml)
F1	2	1	-	60	4	0.1	0.05	10	4
F2	2	1	-	60	4	0.1	0.05	10	4
F3	2	0.5	0.75	60	4	0.1	0.05	10	4
F4	2	0.5	0.5	60	4	0.1	0.05	10	4
F5	2	0.75	0.5	60	4	0.1	0.05	10	4

Evaluation of Fluconazole gel^[7-24]

1. Percentage yield

The empty container was weighed in which the gel formulation was stored then again the container was weighed with gel formulation. Then subtracted the empty container weighed with the container with gel formulation then it gives the practical yield. Then the percentage yield was calculated by the formula.

$$\text{Percentage yield} = \frac{\text{Practical yield}}{\text{Theoretical yield}} \times 100$$

2. Drug content

Weighed 10 gm of each gel formulation were transferred in 250 ml of volumetric flask containing 20 ml of alcohol and stirred for 30 min. The volume was made up to 100 ml and filtered. 1 ml of above solution was further diluted to 10 ml with alcohol and again 1 ml of the above solution was further diluted to 10 ml with alcohol. The absorbance of the solution was measured spectrophotometrically at 260 nm. Drug content was calculated by the following formula.

$$\text{Drug content} = \frac{\text{Absorbance}}{\text{Slope}} \times \text{Dilution factor} \times \frac{1}{1000}$$

3. Determination of pH

Weighed 50 gm of each gel formulation were transferred in 10 ml of beaker and measured it by using the digital pH meter. pH of the topical gel formulation should be between 3 – 9 to treat the skin infections.

4. Spredability

The spredability of the gel formulation was determined, by measuring diameter of 1 gm gel between horizontal plates (20×20 cm²) after 1 minute. The standardized weight tied on the upper plate was 125 gm.

5. Extrudability

The gel formulations were filled into collapsible metal tube or aluminium collapsible tube. The tube were pressed to extrude the material and the extrudability of formulation was checked.

6. Viscosity Estimation

The viscosity of gel was determined by using a Brookfield viscometer DVII model with a T-Bar spindle in combination with a helipath stand.

a) Selection of spindle

Spindle T 95 was used for the measurement of viscosity of all the gels.

b) Sample container size

The viscosity was measured using 50 gm of gel filled in a 100ml beaker.

c) Spindle immersion

The T-bar spindle (T95) was lowered perpendicular in the centre taking care that spindle does not touch bottom of the jar.

d) Measurement of viscosity

The T-bar spindle (T95) was used for determining the viscosity of the gels. The factors like temperature, pressure and sample size etc. Which affect the viscosity was maintained during the process. The helipath T- bar spindle was moved up and down giving viscosities at number of points along the path. The torque reading was always greater than 10%. The average of three readings taken in one minute was noted as the viscosity of gels.

7. In vitro diffusion study

The abdominal skin of Albino mice, weighing 20 – 25 gm of 8 – 10 week old was shaved using hand razor and clean the skin with hot water cotton swab. 5 gm of gel was applied uniformly to skin. The skin was mounted between the compartments of the Frantz diffusion cell with stratum corneum facing the donor compartment. Reservoir compartment was filled with 100 ml phosphate buffer of pH 6.8. The study was carried out at 37 ± 1°C and speed was adjusted until the vortex touches the skin and it carried out for 4½ h. 5 ml of sample was withdrawn from reservoir compartment at 30 min interval and absorbance was measured spectrophotometrically at 260 nm. Each time the reservoir compartment was replenished with the 5 ml volume of phosphate buffer pH 6.8 solution to maintain constant volume.

RESULT AND DISCUSSION

(Table No.2) Percent yield of gel formulations

Formulation	percent yield
F1	99.59%
F2	98.34%
F3	97.44%
F4	99.81%
F5	98.76%

(Table No.3) Drug content of gel formulations

Formulation code	Drug Content
F1	94.41
F2	97.38
F3	98.24
F4	96.52
F5	95.07

(Table No.4) pH of gel formulations

Formulation	pH
F1	6.98
F2	7.01
F3	6.98
F4	6.5
F5	6.79

(Table No.5) Viscosity of gel formulations

Formulation	Viscosity(cp)
F1	8476
F2	4259
F3	4450
F4	4544
F5	6.79

(Table No. 6) Spredability of gel Formulations

Formulation	Spredability	
	R1	R2
F1	1.3	1.9
F2	2.1	2.9
F3	19	2.8
F4	1.7	2.3
F5	1.5	2.1

(Table No. 7) Extrudability of gel Formulations

Formulation	Extrudability
F1	+
F2	+++
F3	+++
F4	++
F5	++

Excellent (+++)

Good (++)

Average (+)

Poor (-)

(Table No.8) In Vitro diffusion chart

Time	% CDR				
	F1	F2	F3	F4	F5
0	0	0	0	0	0
30	12.95	16.87	14.88	14.03	13.67
60	39.51	44.39	41.39	40.76	40.05
90	47	48.49	47.95	47.02	46.91
120	56.59	56.01	57.18	56.24	55.74
150	62.84	61.28	63.59	62.31	61.89
180	71.84	72.69	73.26	72.13	71.84
210	80.17	79.37	82.15	81.68	80.86
240	87.27	86.16	89.07	88.19	87.98
270	95.98	94.09	97.03	96.83	96.03




Fig: In vitro diffusion for F3 Formulation.

CONCLUSION

Various formulation (F1, F2, F3, F4, F5) were developed by using suitable polymer (carbopol 934p and HPMC). Developed formulations of Fluconazole were evaluated for the physicochemical parameters such as percentage yield, drug content, pH, viscosity, spredability, extrudability, in vitro drug diffusion. Viscosity studies of various formulations revealed that formulation F3 was better compare to others. From among all the developed formulation F3 shows better drug diffusion, did good Rheological properties. pH of the F3 formulation is sufficient enough to treat the skin infections. Results indicated that the concentration of carbopol-934 and HPMC K4M significantly affects drug release and rheological properties of the gels. The viscosity of carbopol-934 gels was very high as compared to HPMC K4M gels but both gels showed decrease in drug release with increase in polymer concentration. Thus, gels can be successfully prepared using carbopol-934 and Hydroxypropyl methylcellulose as gelling agents in the

ratio 1:3(carbopol-934 and Hydroxypropyl methylcellulose) suitable for topical application Hence formulation F3 should be further developed for scale-up to industrial production.

REFERENCES

- Provost C.A review on Transparent oil-water gels. *Int J Cosmet Sci*, 1986; 8(7): 233-247.
- Gupta A, Mishra AK, Singh AK, Gupta V, Bansal P. Formulation and evaluation of topical gel of diclofenac sodium using different polymers. *Drug invention today*, 2010; 2(5): 250-253.
- Gerry Fink and the Fink lab. How antifungal drug kill fungi and cure disease. (Online). 2005 Feb 9 (cited 2005 Feb 9); Available from: URL:<http://www.medscape.com/viewprogram/2963-pn>.
- Golinkin HS. Process for fracturing well formulation using aqueous gels. US patent, US 4137182, 1979.
- Singh MP, Nagori BP, Shaw NR. Formulation Development & Evaluation of Topical Gel Formulations Using Different Gelling Agents and Its Comparison with Marketed Gel Formulation. *Int J Pharm Erud*, 2013; 3(3): 1-10.
- B. Niyaz Basha, Kalyani P., Divakar G. Formulation and evaluation of Gel containing Fluconazole-Antifungal. *Int J Drug Dev & Res*, 2011; 3(4): 109-12.
- Doaa AH, Dalia AE, Sally AA, Mohamed AE. Formulation and evaluation of fluconazole topical gel. *Int J Pharm Pharm Sci*, 2012; 4(5): 176-183.
- Sudipta DA, Arnab SA, Ananya BO. Design, development and evaluation of fluconazole topical gel. *Asian J Pharm and clin Res*, 2015; 8(2): 132-13.
- Ashni VE, Sukhdev SI, Rupinder KA, Upendra KJ. A Review: Topical Gels as Drug Delivery Systems. *Int J Pharm Sci Rev Res*, 2013; 23(2): 374-382.
- Prateek CH, Rajesh ST, Umesh RA. A Review on Topical Gel. *Int J Pharm & Bio Arc*, 2013; 4(4): 606-613.
- Loveleen PR, Tarun KG. Topical Gel: A Recent Approach for Novel Drug delivery. *Asian J Biomed and Pharm Sci*, 2013; 3(17): 1-5.
- Debjit B, Harish G, B. Pragati, S. Duraivel, K.P. Sampath Kumar, Recent Advances In Novel Topical Drug Delivery System, the pharma innovation, 2012; 1(9): 12-31.
- Roychowdhury S, Singh DH, Gupta R, Masih D, A review on pharmaceutical gel, *Int J Pharm Res Bio Sci*, 2012; 1(5): 21-36.
- Loveleenpreet k and Prabhjot k, Formulation and evaluation of topical gel of meloxicam, *Int J Res Pharm and Chem*, 2014; 4(3): 619-623.
- P.K. Lakshmi, Marka KK, Aishwarya S, Shyamala B, Formulation and evaluation of ibuprofen topical gel: a novel approach for penetration enhancement, *Int J applied pharma*, 2011; 3(3): 25-30.
- Joshi B, Gurpreet AC, Saini S, Singla V, Emulgel A comprehensive review on the recent advancse in topical Drug delivery, *Int Res J of Pharm*, 2011; 2(11): 66-70.
- Arun RR, Elwin J, Jyoti H, Sreerekha S, Formulation and evaluation of Ketoprofen solid dispersion incorporated topical gels, *Europ J Biomed and Pharm sci*, 2016; 3(1): 156-164.
- Swetha CH, Sellappan VP, Narayana RG, Nagarjuna R, Formulation and evaluation of Clarithromycin Topical Gel, *Int J Drug Dev & Res*, 2013; 5(4): 194-202.
- Ganesh M, Gouri D, Vijay G, Formulation and evalution of herbal gels, *Int J natural products and Res*, 2012; 3(4): 501-505.
- Marwa HS and Ghada FM, Evaluation of Topical Gel Bases Formulated with Various Essential Oils for Antibacterial Activity against Methicillin-Resistant Staphylococcus Aureus, *Trop J Pharm Res*, 2013; 12(6): 877-884.
- Aejaz A, Azmail K, Sanaullah S, Mohsin AA, Formulation and In vitro evaluation of aceclofenac solid dispersion incorporated gels. *Int J App Pharm*, 2010; 2(1): 7-12.
- Barhate SD, Potdar MB, Nerker P, Development of meloxicam sodium transdermal gel, *Int J Pharm Res and Dev*, 2010; 2(5): 1-7.
- Bazigha AK, Eman AF, Sahar FA, Heyam SS, Saeed KA, Development and evaluation of Ibuprofen transdermal gel formulations, *Trop J Pharm Res*, 2010; 9(4): 355-363.
- Jain S, Padsalg BD, Patel AK, Mokale V, Formulation, development and evaluation of Fluconazole gel in various polymer bases, *Asian J Pharm*, 2007; 1(6): 3-8.