

**ORAL SUBMUCOUS FIBROSIS- DURATION & FREQUENCY OF VARIOUS HABIT
FACTOR ARE CORRELATED WITH CLINICAL GRADING****Riya Agrawal***4th Year BDS Peoples Dental Academy Bhopal Madhya Pradesh.**Corresponding Author: Riya Agrawal**4th Year BDS Peoples Dental Academy Bhopal Madhya Pradesh.

Article Received on 22/10/2016

Article Revised on 12/11/2016

Article Accepted on 01/12/2016

ABSTRACT

Oral submucous fibrosis is a precancerous condition. It is now globally accepted as an Indian disease having highest malignant potential than any other oral precancerous lesion. **Aim:** To evaluate the effect of duration & frequency of various habit factor on the incidence & severity of OSMF. **Material and Method:** The study included 100 patients with OSMF attending to the department of oral medicine & radiology bhopal over a period of 3 month. A detailed questionnaire was filled with regarding patient's medical history along with the duration & frequency of gutkha, arecanut, tobacco, pan & smoking. **Result:** A total of 100 patients were studied of which 70 (70%) were males & 30 (30%) were females. Grade I OSMF was seen in 53% (53), Grade II OSMF in 47% (47). Gutkha & other arecanut product were most frequently consumed & showed significant risk in the severity of OSMF. Duration & frequency also significant. **Conclusion:** In this study, it was found that relative risk of OSMF increased with increase in the duration & frequency of the habit especially in younger age group.

KEYWORD: OSMF, Duration & Frequency, Gutkha, Tobacco, Arecanut, Clinical Grading.**I. INTRODUCTION**

Oral submucous fibrosis (OSMF) is a chronic, premalignant condition of the oral mucosa which was first described by Schwartz 1952.^[1] Pindborg and his associates defined the condition as "an insidious chronic disease affecting any part of the oral cavity and sometimes pharynx. although occasionally preceded by and / or associated with juxtaepithelial inflammatory reaction followed by fibroelastic changes in the lamina propria, with epithelial atrophy leading to stiffness of the oral mucosa and causing trismus and inability to eat."^[2] OSMF is well recognized potentially malignant condition in the oral cavity & the transformation rate is as high as 7.6% over a period of ten year have been reported from India.^[3] The age range of patients with OSMF is wide ranging between 20 and 40 years of age.^[4]

When OSMF was first described in 1952, it was classified as an idiopathic disorder.^[5] Later on various researchers put forward many hypothesis suggesting that OSMF is multifactorial origin with possible aetiological factors are areca nut, chilies, micronutrient deficiencies of iron, zinc and essential vitamins. Demonstration of various auto-antibodies and an association with specific HLA antigens has also been proposed.^[6] Areca nut is the main etiological factor in the causation of OSMF.^[7]

The use & dependence on areca nut in the form of pan-masala / Gutkha is rapidly increasing especially among youth in India. Many research has to be done in evidence

based dentistry, the role of duration and frequency of habits to the clinical staging of OSMF.^[8] The main purpose of this study was to correlate these variables of the habit to the clinical grading of OSMF.

II. MATERIAL AND METHOD

This study included 100 patients with osmf attending the department of oral medicine & radiology people dental academy, peoples college of dental science, peoples hospital peoples university Bhopal M.P. over a period of 6 month. A detailed questionnaire was filled with detailed regarding patients name, age, sex, medical history along with the duration & frequency of gutkha, arecanut, tobacco, pan & smoking.

INCLUSION CRITERIA

1. Subjects with definitive habit of gutkha, arecanut, tobacco in any form.
2. Subjects with history of at least one packets of chewable tobacco, gutkha, pan, per day at least 6 month.

EXCLUSION CRITERIA

The subject who were suffering from any other disease or taking any treatment for osmf or presented with any other oral mucosal lesion were excluded from the study.

The cases were divided into three stages according to the severity of the disease described by Nagesh & Bailoor (1993).^[9]

Grade I (mild) subjects had mild blanching, burning sensation on taking spicy food or hot beverages, no restriction in mouth opening, no tongue protrusion.

Grade II (moderate) subjects had moderate to severe blanching, mouth opening reduced, tongue protrusion reduced, burning sensation also in absence of stimuli, palpable bands felt, lymphadenopathy.

The collected data was sorted, tabulated & subjected to statistical analysis.

III. RESULTS

In the present study 100 OSMF subjects were screened, out of which 70% (70) were males and 30% (30) were females and out of which grade I OSMF was seen in 53%(53) & grade II OSMF in 47%(47) subjects.

The gender & the age of sample were found to be insignificant with no effect on the severity of osmf (Table 1) whereas duration and frequency of the habit

showed a significant variation of the clinical grading (Table 2).




FIGURE 1

TABLE 1

GENDER	GRADE I OSMF	GRADE II OSMF	TOTAL (OUT OF 100)
MALE	40 (57.1%)	30 (42.8%)	70
FEMALE	20 (66.6%)	10 (33.3%)	30
AGE RANGE			
15 to 24 years	14 (53.8%)	12 (46.15%)	26
25 to 34 years	6 (60%)	3 (30%)	10
35 to 44 years	33(52.3%)	30 (47.6%)	63

The etiological factors of pan and smoking showed non-significance for gradation of OSMF, whereas gutkha, arecanut & tobacco chewing showed significance(TABLE 2).

TABLE 2

ETIOLOGY FACTOR	GRADE I OSMF	GRADE II OSMF	TOTAL (OUT OF 100)
Gutkha	18 (47.3%)	20(52.6%)	38
Tobacco	9 (45%)	11 (55%)	20
Areacanut	10 (45.4%)	12 (54.4%)	22
Pan	8 (72.7%)	3 (27.2%)	11
Smoking	8 (88.8%)	1 (1.1%)	9

TABLE 3

DURATION GROUP	GRADE I OSMF	GRADE II OSMF	TOTAL (OUT OF 100)
Upto 3 year	12 (54.5%)	10 (45.4%)	22
4 to 7 year	25(53.1%)	22(46.8%)	47
8 to 10 year	10 (58.8%)	7(41.1%)	17
>10 year	6 (42.8%)	8 (57.1%)	14
FREQUENCY GROUP			
Upto 2 per day	10 (66.6%)	5(33.3%)	15
3 to 6 per day	15 (34.8%)	12 (44.4%)	27
7 to 10 per day	23 (53.4%)	20 (46.5%)	43
>10 per day	5 (33.3%)	10 (66.66%)	15

IV. DISSCUSION

Oral submucous fibrosis (OSMF) is considered as chronic and potentially malignant condition of the oral cavity that often leads to oral cancer.^[10] In our study out of 100 OSMF study population we observed 70% (70)

were males and 30% (30) were females, sex predilection conflicting earlier it was thought to be common in female.^[11] But at present study shows male to female ratio being 2.36:1, which was according to Chaturvedy et al in India.^[12] Our study shows that most of the OSMF

patients were between 35 - 44 years. Reporting an age between 20-39 years; while a study conducted among 1000 patients in Central India.^[13] Gutkha and other areca nut product users like tobacco when compared to pan and smoking users showed a significant occurrence of OSMF in the severity of the condition. The reason attributes to the fact that the gutkha and other arecanut product have more dry weight as compared to the other chewing product antioxidant capabilities of pan leaf which is known to be rich in beta-carotene its level decreases in all grades of osmf. Diet rich in beta carotene is used in the management of osmf to reduce disease severity. The effect of Smoking consumption alone have not been found in development of OSMF, but their addition to areca nut using can be a risk factor for OSMF.^[14]

In our study it was found that the patients who were taking the gutkha and other products more than 10 years developed mostly grade II of OSMF. It also found that the frequency of gutkha and other product consuming more than 10 times per day increase the severity of OSMF. Subject who were taking the gutkha and other products less than 10 times per day had grade I OSMF (TABLE 3). As the parameters of our study in the form of duration, frequency it had a significant correlation with the outcome of the severity of the disease. Ghutka is manufactured with the combination of arecanut, tobacco, lime, catechu and flavoring compound. Arecanut has high alkaloid arecoline and tobacco ingredients like nitrosamine, which are absorbed more in the patients who keep it for longer durations.^[15]

V. CONCLUSION

The present study revealed widespread habit of chewing gutkha and other arecanut by product to have shown a major role in occurrence and severity of OSMF. It also found that the relative risk of osmf rises with increase duration and frequency of daily consumption of commercially available arecanut & tobacco by product especially in younger age group. Prevention involve taking step to ban all these carcinogenic product from our society. These measure can play a significant role in the elimination of oral premalignant disease like OSMF & hence reduce the disease of oral cancer.

ACKNOWLEDGEMENT

We sincerely thanks to my parents, friends and my faculty, Department Of Oral Medicine and Radiology who supported during the study.

CONFLICT OF INTEREST

No conflict of interest involved so ever.

SOURCE OF FINDING

Nil.

REFERENCES

1. Angadi PV, Rekha K. Oral submucous fibrosis: a clinicopathologic review of 205 cases in Indians. *Oral and Maxillofacial Surgery*, 2011; 15(1): 15-9.

2. Pindborg JJ, Sirsat SM. Oral submucous fibrosis. *Oral Surg Oral Med Oral Pathol*. 1966; 22: 764-79.[PubMed].
3. Murti PR, Bhonsle RB, Pindborg JJ, Daftary DK, Gupta PC, Mehta FS. Malignant transformation rate in oral submucous fibrosis over a 17-year period. *Community Dent Oral Epidemiol*. 1985; 13(6): 340-341. doi: 10.1111/j.1600-0528.1985.tb00468.x. [PubMed] [Cross Ref].
4. Rajendran R, Sugathan C, Remani P, Ankathil R, Vijayakumar T. Cell mediated and humoral immune responses in oral submucous fibrosis. *Cancer*, 1986; 58(12): 2628-31.
5. Tilakaratne WM, Klinikowski MF, Saku T, Peters TJ, Warnakulasuriya S. Oral submucous fibrosis: Review on etiology and pathogenesis. *Oral Oncology*, 2006; 42: 561- 568.
6. Rajalalitha P & Valli S. Molecular pathogenesis of oral submucous fibrosis- a collagen metabolic disorder. *J Oral Pathol Med*, 2005; 34: 321-8.
7. Ranganathan K, Uma Devi M, Joshua E, Kirankumar K, Saraswathi TR. Oral submucous fibrosis: a case control study in Chennai South India. *J Oral Pathol Med*, 2004; 33: 274-7.
8. Reddy V, Wanjari P, Banda N, Reddy P. Oral Submucous Fibrosis: Correlation of Clinical Grading to various habit factor.
9. Gupta D, Gupta M, Golhar B. Oral submucous fibrosis; clinical study and management by physiofibrolysis. *J Indian Dent Assoc*, 1980; 52: 375-8.
10. Murti P, Gupta P, Bhonsle R, Daftary D, Mehta s. *Intern J Dent Clin*. 2011; 3: 21-24.
10. Li N, Jian X, Hu Y, Xu C, Yao Z, et al. Discovery of novel biomarkers in oral submucous fibrosis by microarray analysis. *Cancer Epidemiol Biomarkers Prev*, 2008; 17: 2249-5.
11. Kiran Kumar K, Saraswathi TR, Ranganathan K, Uma Devi M, Elizabeth J. Oral submucous fibrosis: a clinico-histopathological study in Chennai. *Indian J Dent Res*. 2007; 18: 106-11. doi: 10.4103/0970-9290.33785. [PubMed] [Cross Ref].
12. Chaturvedi V, Sharma A, Chakrabarti S. Salivary coagulopathy and humoral response in oral submucous fibrosis (OSMF). *Journal of the Indian Dental Association*, 1991; 62(3): 51-9.
13. Hazarey V, Erlewad D, Mundhe K, Ughade N. Oral submucous fibrosis: study of 1000 cases from central India. *J Oral Pathol Med*, 2007; 36: 12-7.
14. Shah N, Sharma P. Role of chewing and smoking habits in the etiology of oral submucous fibrosis (OSF): a case control study. *Journal of Oral Pathology & Medicine*, 1998; 27(10): 475-9.
15. Sinor P, Gupta P, Murti P, Bhonsle R, Daftary D, Mehta F, Pindborg J. A case control study of oral submucous fibrosis with special reference to the etiologic role of areca nut. *Journal of Oral Pathology & Medicine*, 1990; 19(2): 94-8.