

PHARMACOGNOSTICAL AND PHYTOCHEMICAL ANALYSIS OF *DESMODIUM TRIFLORUM* LEAVES

¹*Nitin Kumar and ²Saravanan K.

¹Research Scholar, Bhagwant Global University, Kotdwar, Uttarakhand, India.

²Professor, Bhagwant Global University, Kotdwar, Uttarakhand, India.



*Corresponding Author: Nitin Kumar

Research Scholar, Bhagwant Global University, Kotdwar, Uttarakhand, India.

Article Received on 12/01/2025

Article Revised on 02/02/2025

Article Accepted on 22/02/2025

ABSTRACT

Desmodium triflorum is mostly herb or shrub is belongs to the family Fabaceae. *Desmodium triflorum* contains alkaloids including *N, N*-dimethyl tryptophan methyl ester, dimethyltryptamine-*N*-oxide, hypaphorine (structurally related to plakohypaphorine), phenylethylamine, hordenine, tyramine, and trigonelline. *Desmodium triflorum* DC contains ursolic acid, vitexin, genistin, fucosterol. *Desmodium triflorum* plant given in eye diseases, spleen disorders, colic, diarrhea, menorrhagia, breast pain. Leaves used as galactogogue, good in diarrhea, dysentery and convulsion. Leaves are also used as antiseptic. Roots are used in cough, asthma, used as antiseptic and diuretic. In the present study, leaf powder of *Desmodium triflorum* has been investigated for isolation and characterization of bioactive compounds. The isolation and characterization of betasitosterol, stigmasterol were confirmed by spectroscopic methods.

KEYWORDS: *Desmodium triflorum*, Creeping tick trefoil, Steroids, betasisterol, stigmasterol.

1. INTRODUCTION

Desmodium triflorum is mostly herb or shrub is belongs to the family Fabaceae. It is known as tick-trefoil. It is used in the indigenous system of medicine. The other vernacular names are given below.

It grows in throughout eastern ghats and on plains upto 1600m with a wide range of habitats, optimum size in clay soil, especially by river bed and along road sides. These are mostly inconspicuous plants; few have bright or large flowers. Though some can become sizeable plants, most are herbs or small shrubs. Their fruit are loments, meaning each seed is dispersed individually enclosed in its segment. This makes them tenacious plants and some species are considered weeds in places.^[1,2,3]

2. MATERIALS AND METHODS

2.1 Plant Materials

Desmodium were leaves collected from Kotdwar in Uttarakhand and authentication of the above species was carried out Thirumala college of Pharmacy, Nizamabad, AP.

2.2 Standardization of *Desmodium triflorum* leaves

The leaf drugs of *Desmodium triflorum* was dried at room temperature, until they were free from the moisture and subjected to physical evaluation with different parameters. The parameters which were used for

evaluation are nature, odour, colour, taste, size, shape, width and length.

Finally leaves were subjected to size reduction by electric mixer grinder to get coarse powder and passed through sieve No. 40 to get uniform powder. Then the uniform powders were subjected to standardization with different parameters as per pharmacopoeias/literatures.

2.2.1 Quantitative Standards^[4]

2.2.1.1 Foreign Matter

Foreign organic matter is the material consisting of any or all of the following.

- (1) Parts of the organ or organs from which the drug is derived other than the parts named in the definition and description or for which the limit is prescribed in the individual monograph.
- (2) Any organs other than those named in the definition and description.
- (3) Matter not coming from the source plant and
- (4) Moulds, insects or other animal contamination.

2.2.1.2 Method

Weigh 100 to 500 g or the quantity specified in the individual monograph, of the original sample and spread it out in to a thin layer. Inspect the sample with unaided eye or with the use of a 6x lens and separate the foreign matter manually as completely as possible. Weigh and determine the percentage of foreign matter from the

weight of the drug taken. Use the maximum quantity of sample for coarse or bulky drugs.

2.2.2. Ethanol soluble extractive

Macerate 5 g of the air-dried drug, coarsely powdered, with 100 ml of ethanol of the specified strength in a closed flask for 24 hours, shaking frequently during the first 6 hours and allowing to stand for 18 hours. Thereafter, filter rapidly taking precautions against loss of ethanol evaporate 25 ml of the filtrate to dryness in tarred flat-bottomed shallow dish, dry at 105°C and weigh. Calculate the percentage of ethanol-soluble extractive with reference to the air-dried drug.

2.2.3. Water soluble extractive

Add 5 g to 50 ml of water at 80°C in a stoppered flask. Shake well and allow to stand for 10 minutes, cool, add 2 g of kieselguhr and filter. Transfer 5 ml of the filtrate to a tarred evaporating dish, 7.5 cm in diameter, evaporate the solvent on a water bath, continue drying for 30 minutes, finally dry in a steam oven for 2 hours and weigh the residue. Calculate the percentage of water-soluble extractive with reference to the air-dried drug.

2.2.4. Total ash

Unless otherwise stated in the individual monograph, weigh accurately 2 to 3 g of the air-dried crude drug in a tarred platinum or silica dish and incinerate at a temperature not exceeding 450°C until free from carbon, cool and weigh. If a carbon-free ash cannot be obtained in this way, exhaust the charred mass with hot water, collect the residue on an ashless filter paper, incinerate the residue and filter paper until the ash is white or nearly so, add the filtrate, evaporate to dryness and ignite at a temperature not exceeding 450°C Calculate the percentage of ash with reference to the air-dried drug.

2.2.5. Acid insoluble ash

Boil the ash obtained from the procedure mentioned in the total ash with 25 ml of 2M hydrochloric acid for 5 minutes, collect the insoluble matter in a Gooch crucible or on an ashless filter paper, wash with hot water, ignite, cool in a desiccators and weigh. Calculate the percentage of acid insoluble ash with reference to the air-dried drug.

2.2.6. Moisture content (by Loss on drying)

Weigh a glass-stoppered, shallow weighing bottle that has been dried under the same condition to be employed in the determination. Transfer to the bottle the quantity of the sample specified in the individual monograph, cover it and accurately weigh the bottle and the contents. Distribute the sample as evenly as practicable by gentle sidewise shaking to depth not exceeding 10mm. Place the loaded bottle in the drying chamber (oven or desiccators) as directed in the monograph, remove the stopper and leave it also in the chamber. Dry the sample to constant weight or for the specified time and at the temperature indicated in the monograph. After drying is completed, open the drying chamber, close the bottle promptly and allow it to cool to room temperature (where applicable)

in a desiccator before weighing. Weigh the bottle and the contents.

2.3 Extraction of *Desmodium triflorum* linn. leaves

The shade-dried leaves of *Desmodium triflorum* Linn, family Fabaceae was reduced to fine powder (# 40 size mesh) and around 300 gms of powder was subjected to successive hot continuous extraction (Soxhlet) with petroleum ether, and alcohol. Finally the drug was macerated with chloroform-water. Each time before extracting with the next solvent the powdered material was air dried in hot air oven below 50°C. After the effective extraction, the solvent were distilled off, the extract was then concentrated on water bath and the extract obtained with each solvents was weighed. Its percentage was calculated in terms of air-dried weight of plant material. The colour and consistency of the extracts was noted.

The obtained alcoholic and aqueous extracts were subjected to chemical investigation and pharmacological screening for its anticonvulsant activity by using MES test.

The shade dried powder around 500gm of *Desmodium triflorum* was subjected to continuous hot extraction with methanol. After the effective extraction, the extract was then concentrated on water bath and the extract obtained was weighed. About 100 gm of extract was taken for isolation.

2.4 Qualitative Chemical Investigation of Extract^[5,6]

Qualitative chemical tests were conducted for all the extracts of *Desmodium triflorum* to identify the various phytoconstituents. The various tests and reagents used are given below and observations are recorded in the table no.1 and table no.2

Tests for Carbohydrates

Molisch's test (General test): To 2-3 ml aqueous extract, added few drops of α -naphthol solution in alcohol, shaken and added concentrated H₂SO₄ from sides of the test tube was observed for violet ring at the junction of two liquids.

For Reducing Sugars

- Fehling's test: 1 ml Fehling's A and 1ml Fehling's B solutions was mixed and boiled for one minute. Added equal volume of test solution. Heated in boiling water bath for 5-10 min was observed for a yellow, then brick red precipitate.
- Benedict's test: Equal volume of Benedict's reagent and test solution in test tube were mixed. Heated in boiling water bath for 5 min. Solution appears green, yellow or red depending on amount of reducing sugar present in test solution.

Tests for Monosaccharides: Barfoed's test: Equal volume of Barfoed's reagent and test solution were

added. Heated for 1-2 min, in boiling water bath and cooled. Observed for red precipitate.

Tests for Hexose Sugars: *Cobalt-chloride test:* 3 ml of test solution was mixed with 2ml cobalt chloride, boiled and cooled. Added FeCl_3 drops on NaOH solution. Solution observed for greenish blue (glucose), purplish (Fructose) or upper layer greenish blue and lower layer purplish (Mixture of glucose and fructose).

Tests for Non-Reducing Sugars

- Test solution does not give response to Fehling's and Benedict's tests.
- Tannic acid test for starch: With 20% tannic acid, test solution was observed for precipitate.

Tests for Proteins

- Biuret test (General test): To 3 ml T.S added 4% NaOH and few drops of 1% C_4SO_4 solution observed for violet or pink colour.
- Million's test (for proteins): Mixed 3 ml T.S. with 5 ml Million's reagent, white precipitate. Precipitate warmed turns brick red or precipitate dissolves giving red colour was observed.
- Xanthoprotein test (For protein containing tyrosine or tryptophan): Mixed 3ml T.S. with 1 ml concentrated H_2SO_4 observed for white precipitate.
- Test for protein containing sulphur: Mixed 5 ml T.S. with 2 ml 40% NaOH and 2 drops 10% lead acetate solution. Solution was boiled it turned black or brownish due to PbS formation was observed.
- Precipitation test: The test solution gave white colloidal precipitate with following reagents:
 - Absolute alcohol
 - 5% HgCl_2 solution
 - 5% C_4SO_4 solution
 - 5% lead acetate
 - 5% ammonium sulphate

Tests for Steroid

- Salkowski Reaction: to 2 ml of extract, 2 ml chloroform and 2 ml concentrated H_2SO_4 was added. Shaked well, whether chloroform layer appeared red and acid layer showed greenish yellow fluorescence was observed.
- Liebermann-Burchard Reaction: Mixed 2ml extract with chloroform. Added 1-2 ml acetic anhydride and 2 drops concentration H_2SO_4 from the side of test tube observed for first red, the blue and finally green colour.
- Liebermann's reaction: Mixed 3 ml extract with 3 ml acetic anhydride. Heated and cooled. Added few drops concentrated H_2SO_4 observed for blue colour.

Tests for Amino Acids

- Ninhydrin test (General test):- 3 ml T.S. and 3 drops 5% Ninhydrin solution were heated in boiling water bath for 10 min. observed for purple or bluish colour.

- Test for Tyrosine: Heated 3 ml T.S. and 3 drops Million's reagent. Solution observed for dark red colour.
- Test for tryptophan: To 3 ml T.S. added few drops glyoxalic acid and concentrated H_2SO_4 observed for reddish violet ring at junction of the two layers.

d) Tests for Glycosides

Tests for Cardiac Glycosides

- Baljet's test:- A test solution observed for yellow to orange colour with sodium picrate.
- Legal's test (For cardenolides):- To aqueous or alcoholic test solution, added 1 ml pyridine and 1 ml sodium nitroprusside observed for pink to red colour.
- Test for deoxysugars (Kellar Killani test):- To 2 ml extract added glacial acetic acid, one drop of 5% FeCl_3 and concentrated H_2SO_4 observed for reddish brown colour at junction of the two liquid and upper layers bluish green.
- Liebermann's test (For bufadenolides):- Mixed 3 ml extract with 3 ml acetic anhydride. Heated and cooled. Added few drops concentrated H_2SO_4 observed for blue colour.

Tests for Saponin Glycosides

- Foam test: The drug extract or dry powder was shaken vigorously with water. Persistent foam was observed.
- Haemolytic test: Added test solution to one drop of blood placed on glass slide. Haemolytic zone whether appeared was observed.

Tests for Coumarin Glycosides

Test solution when made alkaline, observed for blue or green fluorescence.

Tests for Flavonoids

- Shinoda test: - To dried powder or extract, added 5 ml 95% ethanol, few drops concentrated HCl and 0.5 g magnesium turnings. Pink colour was observed.
- To small quantity of residue, added lead acetate solution observed for Yellow coloured precipitate.
- Addition of increasing amount of sodium hydroxide to the residue whether showed yellow colouration, which was decolorized after addition of acid was observed.
- Ferric chloride test: - Test solution, added few drops of ferric chloride solution observed for intense green colour.

e) Tests for Alkaloids

- Dragendroff's test: To 2-3 ml filtrate added few drops Dragendroff's reagent observed for orange brown precipitate.
- Mayer's test: - 2-3 ml filtrate with few drops Mayer's reagent observed for precipitate.
- Hager's test: - 2-3 ml filtrate with Hager's reagent observed for yellow precipitate.

- d) Wagner's test: - 2-3 ml filtrate with few drops of Wagner's reagent observed reddish brown precipitate.

Tests for Tannins and Phenolic Compounds

To 2-3 ml test solution, added few drops of whether showed following was observed.

- 5% FeCl₃ solution: - Deep blue-black coloured.
- Lead acetate solution: - White precipitate.
- Gelatin solution: - White precipitate.
- Bromine water: - Decoloration of bromine water.
- Acetic acid solution: - Red colour solution.
- Potassium dichromate: - Red precipitate.
- Dilute iodine solution: - Transient red colour.
- Dilute HNO₃: - Reddish to yellow colour.

2.5 Isolation of Phytoconstituents by using column chromatography

The methanolic extract of coarsely powdered (500gm) was prepared by using soxhlet apparatus. The crude extract was evaporated to dryness in a water bath to give dark brown mass. The methanolic extract around 100 gm was subjected to column chromatography on silica gel (60-120 mesh) using varying polarities, starting from Chloroform and Methanol to yield several fractions. The column was eluted firstly with pure chloroform and the methanolic extract was subjected to the column for isolation. The column eluted with chloroform and methanol (95:5, 90:10, 80:20 and 50:50). The elute from different solvents were concentrated to give dry residues. The obtained residues were further purified by rechromatogram by using pure solvent chloroform. The samples were confirmed by using TLC, by their melting point, IR NMR and MASS spectroscopy.

2.6 Spectroscopic Characterization

Different spectroscopic methods were used to elucidate the structure elucidation of isolated compounds. Among the spectroscopic techniques IR, ¹H NMR, ¹³C-NMR, MASS and GC-MS were carried out.

2.6.1 IR

The infrared spectra were recorded on PPERKIN ELMER SYSTEM ONE FTIR/ATR (Model: Spectrun one: FT-IR Spectrometer) Scan range 1-o cm⁻¹ with globar and mercury vapour lamp.

2.6.2 Mass

The mass spectra were recorded by using the GEOL GCMATE II GC-MS double focusing instrument, the maximum resolution 6000, and maximum calibrated mass: 1500 Daltons and source options are electron impact and chemical ionization.

2.6.3 NMR

The ¹H NMR and ¹³C NMR spectra were recorded by using Bruker AVANCE III 500 MHz (AV500 MHz) multinuclei solution NMR Spectrometer with 11.7 Tesla super conducting long hold magnet, activity shielded with standard bore (narrow bore: 5-5.4 cm) built in

cryoshims, 34 channel room temperature shims, amplifier power 300W, HP Work station with LCD TFT monitor and Windows XP based TOPSPIN-2 software with an additional software for processing. The IR, MASS and NMR spectral studies were carried out in Sophisticated Analytical Instrumental Facility (SAIF), Indian Institute of Technology Madras.

2.6.4 Melting points were measured by SRS Optimelt MPA 100 instrument.

3. RESULT AND DISCUSSION

In the present study, *Desmodium triflorum* leaves were collected and authentication of the drugs was carried out at Thirumala college of Pharmacy, Nizamabad, Telungana which was subjected to physical evaluation with different parameters which were prescribed in the pharmacopoeias/literatures, such as Foreign matters, Ethanol soluble extractive, Water soluble extractive, Total ash, Acid insoluble ash, Moisture content etc. the results of the standardized *Desmodium triflorum* leaves are given in the Table No. 1.

The standardized powdered leaf drugs of *Desmodium triflorum* was subjected to extraction by continuous or successive hot Soxhlet method according to increasing polarity order of the solvent, extracts were filtered, concentrated to dryness and weighed. The percentage was calculated in terms of air dried weight of the plant materials subjected to extraction by successive hot Soxhlet method. The resultant extracts were subjected to phytochemical investigation by the standard procedure prescribed in the text books. Methanolic extract was prepared and subjected to isolation of constituents.

The results of qualitative chemical investigation of various extracts of *Desmodium triflorum* have indicated the presence of the following compounds.

Pet ether extracts -	Steroids,	triterpenes,
Glycoside flavanoids.		
Chloroform extract	-Steroids,	glycoside,
flavanoids.		
Alcohol extract -	Carbohydrates, tannins and	
phenolic compounds.,		
Aqueous extract -	Carbohydrates, protein, amino	
acid and glycoside.		

The methanolic extract (of coarsely powdered 500gm) was prepared by using soxhlet apparatus. The crude extract was evaporated to dryness in a water bath to give dark brown mass. The methanolic extract around 80 gm was subjected to column chromatography on Silica Gel (230-400 Mesh) using varying polarities, starting from chloroform and methanol to yield several fractions. The column was first eluted with pure chloroform and methanolic extract was subjected to column for isolation.

The column eluted with Chloroform: Methanol (95:5, 90:10, 80:20 and 50:50). The elute from different solvents were concentrated to get residue, further

purified by rechromatogram with pure chloroform or methanol.

Three compounds were separated from methanolic extract of *Desmodium triflorum* fractions. The TLC shown two spots in the mixture were separated by using Chloroform: Methanol (97:3) solvent system. Further these compounds were confirmed by TLC method. The solvent system is Hexane:Acetone (3:1) by using 10% H₂SO₄ as detecting agent. The R_f value matches with the compounds beta-sitosterol 0.6 stigma sterol 0.58.

FT-IR

The resulted spectral graphs are presented in Fig 1. The IR spectral analysis revealed that a broad peak at 3395cm⁻¹ (S) indicating the presence of OH group. The peaks at 2924 & 2853cm⁻¹ indicating the presence of C=O and C-O stretching respectively. The peak at 1075cm⁻¹ indicating cycloalkane. Other peaks are 1736, 1645, 1461, 1408, 1167, 887, 721, 604 cm⁻¹. These data are compared with the literatures which are reported earlier.

MAS

The MS of this compound suggested that the molecular mass is 414 (Molecular formula C₂₉H₅₈O) having characteristic fragments observed at m/z 414, 390, 368, 346, 334, 310, 297, 279, 254, 212, 197, 169, 153, 136, 118, 108 and 93. The results are shown in Fig2.

¹H-NMR

The proton NMR spectrum 7 which are exhibited a broad triplet at 6.560, 6.551 and 6.541 and multiplet at 3.79, 3.73 and 3.70 corresponding to H-6 olefinic proton and H 30 proton respectively. Rest of the protons appeared in the high field region in between 2 to 3 ppm. The figure 3&4 shows the NMR spectra of the compound 2.

¹³C-NMR

The ¹³C- NMR spectrum Fig.5 shows the following peaks.

188.77, 165.44, 164.03, 158.48, 156.02, 135.25, 129.38, 116, 110.61, 105.15, 55.74, 40.41, 40.33, 40.24, 40.16, 40.07, 39.91, 39.74, 39.57 and 39.41.

All spectral data were compared with earlier reported studies.^[7, 8, 9, 10, 11, 12]

Sl.No.	Parameter	IP/BP as per literature	Observation
I.	Physical Test Colour Odour Taste	- Dark Green colour - Odourless - Slightly bitter	Green colour Odourless Slightly bitter
II.	Extractive value Aqueous Alcohol soluble	- Not less than 4.0% - Not less than 10.0%	4.28% 10.58%
III.	Moisture content (By Loss on Drying)	Not more than 3.9%	4.10%
IV.	Ash values Total Ash Acid Insoluble Ash -	Not more than 7.05% Not more than 2.72%	8.62% 4.5%

Table 2: Results of Qualitative Chemical Investigation of *Desmodium triflorum* leaves.

Sl. No.	Name of Test	Extracts			
		Pet. Ether	Chloroform	Alcohol	Aqueous
I.	Test for Carbohydrates				
	Molisch's test (General test)	-	-	-	+
	A) Test for reducing sugars				
	a) Fehling's	-	-	-	+
	b) Benedict's test	-	-	-	+
II	Test for Proteins				
	a) Biuret test (General test)	-	-	-	+
	b) Millions test for Proteins	-	-	-	-
	c) Xanthoprotein test	-	-	-	+
	d) Test for proteins containing sulphur	-	-	-	-
	e) Precipitation test	-	-	-	+
III	Test for Amino acids				
	a) Ninhydrin test (General test)	-	-	-	+
	b) Test for Tyrosine	-	-	-	-
	c) Test for Tryptophan	-	-	-	-
	d) Test for Cystein	-	-	-	+
	Test for Steroids				
	a) Salkowski reaction	+	-	+	-

	b) Liebermann-Burchard reaction	+	-	+	-
	c) Liebermann reaction	+	-	+	-
	Test for Triterpenoids				
	a) Salkowski reaction	+	-	+	-
	b) Liebermann-Burchard reaction	+	-	+	-
VI	Test for Glycosides				
	A) Test for cardiac glycosides				
	a) Baljet test	-	-	-	-
	b) Legals test (Test for Cardenolides)	-	-	-	-
	c) Test for deoxy sugars	-	-	-	-
	d) Liebermann's test	+	-	+	-
	B) Test for Anthraquinone glycosides				
	a) Borntrager's test	-	-	+	+
	b) Modified Borntrager's test	-	-	+	+
	C) Test for Saponin glycosides				
	a) Foam test	-	-	+	+
	b) Haemolysis test	-	-	-	-
	D) Test for Coumarin glycosides				
	a) Alkaline reagent test	-	-	-	-
	b) NaOH soaked paper test	-	-	-	-
VII	Test for Flavonoids				
	a) Ferric chloride test	+	+	+	+
	b) Shinoda test	+	+	+	+
	c) Alkaline reagent test	+	+	+	+
	d) Lead acetate test	+	+	+	+
VIII	Test for Alkaloids				
	a) Dagendroff's test	-	-	+	+
	b) Mayers test	-	-	+	+
	c) Hagers test	-	-	+	+
	d) Wagners test	-	-	-	-
	e) Murexide test for purine alkaloids	-	-	-	-
IX.	Test for tannins and phenolic compounds				
	a) 5% FeCl ₃ solution	-	-	+	+
	b) Lead acetate solution	+	+	+	+
	c) Gelatin solution	-	+	-	-
	d) Bromine water	-	+	-	-
	e) Acetic acid solution	-	-	-	-
	f) Dilute iodine solution	-	+	-	-
	g) Dilute HNO ₃	-	-	-	-
	h) Dilute potassium permanganate solution	-	-	-	-
X.	Test for Lipids	+	-	+	-

Note: '+' Present '-' Absent




Figure1: IR Spectra of compound1.






Figure 2: Mass spectra of compound 1.

Compound-2**Figure 3: HNMR Spectra of compound1.****Figure 4: Compound-1 1HNMR.**


Compound-1 ^{13}C NMR

Figure 5: CNMR Spectra of Compound.

CONCLUSION

The standardized leaf powder of *Desmodium triflorum* was subjected to extraction with various solvents, the resultant extracts were subjected to phytochemical investigation. As per phytochemical investigation, steroids flavanoids, triterpenes, carbohydrates glycosides amino acids and tannins were found to be present in the various extracts. Four chemical constituents were isolated from *Desmodium triflorum* are beta sitosterol, stigmasterol. Further confirmation was carried out by IR, MS, HNMR and CNMR spectroscopy.

NIST Library data. MS of this compound 2 suggested that the molecular mass is 414. The proton NMR spectrum which exhibited a broad triplet at 6.560, 6.551 and 6.541 and multiple at 3.79, 3.73 and 3.70 corresponding to H-6 olefinic proton and H 30 proton respectively. From the above information we concluded that the compound 1 is beta sitosterol. Another compound 2 was confirmed by TLC using n-Hexane: Acetone (3:1) having R_f value of 6.0 was confirmed as Compound 3 as stigma sterol.

MS of this compound 2 suggested that the molecular mass is 414. The proton NMR spectrum which exhibited a broad triplet at 6.560, 6.551 and 6.541 and multiple at 3.79, 3.73 and 3.70 corresponding to H-6 olefinic proton and H 30 proton respectively. From the above information that the compound 2 is beta-sitosterol (melting point 139) and other compound is stigmasterol (melting point 165).

ACKNOWLEDGEMENT

We are sincerely thankful to Bhagwant Global University for providing necessary facilities.

REFERENCES

1. <http://indianbiodiversity.org/species/show/229506>.
2. http://en.wikipedia.org/wiki/Gorana_triflora.
3. V S Agarwal, Drug Plants of India, Vol I, Kalyani Publishers, Page 328-9.
4. Indian Herbal Pharmacopoeia Indian drug manufacturers Association, 2002; 134-141.
5. Kokate CK., Practical pharmacognosy. 4th Ed. Dallaph Prakashan., 1994; 107-111.

6. Khandelwal KR., Practical pharmacognosy. Techniques and Experiments Nirali Prakashan., 13th Ed, 1995; 157-161.
7. Chaturvedula V S P. and Prakash I., Isolation of stigmasterol and beta sitosterol from the dichloromethane extract of *Rubus suavissimus.*, International Current Pharmacecutical Journal, 2012; 1: 239-242.
8. Kamboj A and Saluja A K. Isolation of stigmasterol and betasitosterol from petroleum ether extract of aerial parts of *Ageratum conyzoides*(Asteraceae). Inernational Journal of Pharmacy and Pharmaceutical Sciences, 2011; 3(1): 94-96.
9. Arjun Patra et al., Isolation and characterization of stigmast-5-en-3 β -ol (Betasitosterol) from the leaves of *Hygrophila spinosa* T. Anders. Inernational Journal of Pharma Science and Research, 2010; 1(2): 95-100.
10. Tripathi et al., Phytochemical studies for the roots of *Girardinia heterophylla*. Oriental Journal of Chemistry, 2013; 29(3): 1143-1148.
11. Saratha V., Iyyam Pillai S. and Subramanian S., Isolation and characterization of lupeol, A triterpenoid from *Calotropis gigantea* latex. International Journal of pharmaceutical Sciences Review and research. Article 010, 2011; 10(2): 54-57.
12. Kang W Y., Li G H., Hao X J. Two new triterpenes from *Neonaudea sesslifolia*. Acta Botanica Sinica, 2003; 45(8): 1003-1007.