ejpmr, 2020, 7(1), 506-509

SJIF Impact Factor 7.065

EUROPEAN JOURNAL OF PHARMACEUTICAL AND MEDICAL RESEARCH

www.ejpmr.com

Review Article

ISSN 2394-3211

EJPMR

SHIFT WORK AND ITS CONSEQUENCES: A REVIEW OF LONG-TERM HEALTH RISKS

Manju Dewan*

Assistant Professor in Zoology, DAV College, Sector 10, Chandigarh.

***Corresponding Author: Dr. Manju Dewan**

Assistant Professor in Zoology, DAV College, Sector 10, Chandigarh.

Email Id: manjudewan72@gmail.com

Article Received on 18/11/2019 **Article Revised on 08/12/2019** **Article Accepted on 01/01/2020**

ABSTRACT

Modern economies cannot function without shift workers, especially in sectors like healthcare, manufacturing, transportation and emergency services that depend on constant operations. While shift work is necessary to maintain productivity and meet societal needs, it also presents significant challenges for workers' health and well-being. The disruption of the body's natural circadian rhythm—the internal clock that regulates sleep-wake cycles—is a central concern associated with shift work. Irregular schedules, including night shifts and rotating shifts, interfere with this rhythm, resulting in wide range of physical, mental and emotional health issues. Numerous studies have demonstrated how shift work negatively impacts employees' health. Long-term sleep disturbance, elevated risk of metabolic disorders, cardiovascular diseases, mental health problems and even a higher occurrence of some cancer. The misalignment between the body's biological clock and the demands of the job contributes to various health challenges including sleep deprivation, fatigue, stress and compromised immune function. This review aims to synthesize existing research on the health issues related with shift work, offering a comprehensive overview of the psychological and physiological impacts. Additionally, the work environment, social isolation and lifestyle factors linked with shift work like unhealthy eating habits and lack of physical activity, further exacerbate these health risks.

KEYWORD:- Shift work, circadian rhythm, health issues, wellbeing.

INTRODUCTION

Presently shift work is an integral part of modern society, especially in industries such as healthcare, manufacturing, transportation and emergency services. Although shift work is necessary to sustain services and production, it has several drawbacks, chief among them being the health and welfare of employees. The disturbance of the circadian rhythm, the body's internal clock that controls sleep-wake cycles, is the main cause for concern. Physical, mental and emotional health may suffer significantly from the circadian system's misalignment with irregular work schedules such night shifts, rotating shifts and long work hours (Arendt, 2010; Puttonen et al., 2010). The main causes of these health concerns include irregular circadian rhythms, exhaustion and sleep disruptions (Akerstedt, 2003; Vetter et al., 2016).

Long-term sleep deprivation is a prevalent problem among shift workers and it has been connected to immune system weakness, elevated stress and decreased cognitive function (Bowers & Moyer, 2017). Many health conditions such as sleep disorders, metabolic disorders, cardiovascular diseases, mental health problems and even cancers, are more likely to affect shift workers. A series of detrimental impacts on different biological processes are caused by the body's incapacity to synchronize with the inconsistent work patterns. Long-term health hazards are also increased by shift employment, which frequently exacerbates lifestyle variables like stress, sedentary behaviour and less nutritive diet.

There is also a need for further exploration of how these health problems can be decreased both through organizational changes in work scheduling and individual strategies that promote health and well-being. The aim of this review is to explore the relationship between shift work and health issues, revising existing research and examining the physiological and psychological consequences faced by shift workers. In addition, potential solutions and preventive measures will be discussed to help these health risks and promote a healthier work-life balance for shift workers. This review paper will contribute to a broader understanding of how shift work affects health, providing insight into effective policies for addressing these issues and improving the quality of life for workers engaged in non-standard work schedules. This review will offer an understanding of the wider effects of shift work on worker health by observing number of researches and conclusions. This review's ultimate objective is to provide guidance for future studies and workplace regulations that can better promote shift workers' health and wellbeing and will suggest possible solutions to these issues.

METHODOLOGY

The methodology for writing a review study on shift work and health issues involves several key steps to ensure a detailed, systematic and organized approach to analyze existing research. This section details the processes and criteria used to identify, select and analyze relevant studies, as well as the style for analyzing the data and these findings. The objective of the study is to provide a comprehensive overview of the current literature on the impact of shift work on health. A systematic approach was used to identify peer-reviewed articles, books and other scholarly sources related to shift work and its associated health issues.

DISCUSSION

Shift work has become a worldwide feature of current labor markets chiefly in industries that demand uninterrupted operation such as healthcare, manufacturing, transportation and emergency services. Though shift work is vital to maintain round-the-clock services and meeting societal needs, but it impacts the disruption of the body's circadian rhythm - the natural internal clock that regulates sleep-wake cycles. The disconnect between the body's natural biological functions and irregular work hours can result in numerous physical, mental and emotional health challenges. Circadian rhythms are governed by the Suprachiasmatic Nucleus (SCN) located in the hypothalamus, which primarily reacts to light and dark cues. The SCN orchestrates the secretion of hormones such as melatonin and cortisol, which are essential for regulating sleep, wakefulness, and alertness (Saper et al, 2005). This system is disrupted when shift workers encounter artificial light during night time and attempt to rest during daylight hours, creating a misalignment between their internal biological clock and the external environment (Cajochen et al, 2003). Irregular sleep patterns, along with factors such as poor nutrition, insufficient physical activity and heightened stress levels, lead to a decline in quality of life and potential long-term health repercussions. Shift workers, especially those on night or rotating shifts, frequently suffer from sleep deprivation, fatigue and reduced cognitive function. Although there is an increasing amount of research linking shift work to negative health effects, the specific mechanisms by which shift work contributes to these health problems remain inadequately understood. Additionally, there is still a lack of clarity on effective strategies to alleviate the adverse impacts of shift work and enhance the health outcomes for these workers.

Given the extensive occurrence of shift work across various industries, it is crucial to conduct a thorough review of the existing literature to pinpoint the primary health risks linked to shift work and to explore methods for mitigating these risks. Research has consistently indicated that individuals engaged in shift work face a heightened risk of developing sleep disorders, cardiovascular issues, metabolic conditions, mental health disorders (including depression and anxiety) and certain types of cancer. Additionally, shift work has been correlated with long-term health challenges such as obesity, type-2 diabetes and cardiovascular diseases, likely stemming from alterations in lifestyle factors such as diet and physical activity (Kivimaki et al., 2011; Pan et al., 2011). The relationship between shift work and cancer risk, particularly concerning breast and prostate cancer, has received considerable attention, with studies proposing that the disruption of melatonin production—a hormone that regulates sleep and functions as an antioxidant—may contribute to an increased risk of cancer (Haus & Smolensky, 2006; Schernhammer et al., 2001). Despite the growing body of evidence, the precise mechanisms that connect shift work to these health outcomes remain inadequately understood, necessitating further research to investigate the underlying biological processes. This review seeks to provide a comprehensive synthesis of the current literature regarding the health risks associated with shift work. This paper will analyze the physiological, psychological and social impacts of shift work, emphasizing the significant health challenges faced by individuals in various shift-oriented occupations. Furthermore, the review will investigate possible strategies to alleviate these risks including modifications to work schedules, enhancements in sleep hygiene and the introduction of workplace health initiatives. The ultimate goal of this review is to aid in the formulation of policies and interventions aimed at enhancing the well-being of shift workers and reducing the long-term health hazards linked to irregular work hours.

1. Sleep disorders

The impact of shift work on sleep is one of its most immediate and significant consequences. Irregular work hours disrupt the body's circadian rhythm, leading to common sleep disturbances, especially among those on night or rotating shifts. Shift workers frequently suffer from insomnia, which manifests as difficulty in both falling asleep and maintaining sleep. Akerstedt (2003) found that these workers tend to have shorter sleep durations and lower sleep quality compared to those who work standard daytime hours. The disruption of circadian rhythms results in sleep deprivation, adversely affecting cognitive abilities, alertness, and overall productivity (Puttonen et al., 2010).

A major health concern linked to shift work is the prevalence of sleep disorders, particularly Shift Work Sleep Disorder (SWSD), characterized by chronic insomnia, excessive daytime sleepiness and diminished cognitive function. Disrupted sleep patterns lead to a decrease in restorative sleep, which can hinder both cognitive and physical performance (Vetter et al., 2016).

Additionally, prolonged sleep deprivation and circadian misalignment have been associated with a heightened risk of cardiovascular issues like hypertension, coronary heart disease and stroke. These issues are often compounded by metabolic disruptions as obesity, insulin resistance and type -2 diabetes.

2. Cardiovascular diseases

Shift work has been consistently linked to an increased risk of cardiovascular diseases (CVD), hypertension, heart disease and stroke. The association between shift work and CVD is thought to be influenced by several factors like stress, disrupted sleep, poor diet and lack of physical activity. Kivimaki et al. (2011) have established that shift workers are at greater risk of developing hypertension compared to those on regular daytime schedules. The long-term effects of shift work on cardiovascular health may be due to chronic inflammation, increased cortisol levels (a stress hormone) and dysregulation of the autonomic nervous system, all of which are related with cardiovascular disease. The nurses' health study conducted by Schernhammer et al. (2001) found that rotating shift work increased the risk of coronary heart disease among women.

3. Metabolic and Endocrine Disorders

The interruption of circadian rhythms associated with shift work also affects metabolic processes. Studies have shown that shift workers are at greater risk of developing metabolic disorders, particularly obesity and type-2 diabetes. Shift work, by contributing to these factors, plays important role in the development of metabolic syndrome (Kivimaki et al., 2011) that increase the risk of heart disease, stroke and diabetes. Irregular eating habits, high-calorie snacks or meals combined with reduced physical activity due to fatigue, increases the risk of obesity (Pan et al., 2011).

4. Mental health issues

Shift work is associated with a range of mental health challenge as depression, anxiety, stress and cognitive loss. The chronic disruption of sleep, irregular work schedules, social-isolation and lack of rest can significantly affect emotional well-being (Bowers & Moyer, 2017). The constant pressure of adjusting to irregular hours, coupled with the physical and mental strain of night shifts or extended work hours, increases the risk of chronic stress. Stress adversely affects both mental and physical well-being, leading to issues such as cardiovascular diseases, metabolic disorders and sleep disturbances (Arendt, 2010). Prolonged sleep deprivation and fatigue resulting from inconsistent shift patterns can hinder cognitive performance, especially in areas like attention, memory, and decision-making. These cognitive deficits are particularly significant among individuals engaged in long-term shift work and those facing ongoing sleep interruptions (Akerstedt, 2003).

5. Cancers' Risk

Numerous research studies indicate that shift work, particularly during night hours, may be linked to a heightened risk of specific cancers, such as breast and prostate cancer. This association is thought to stem from the disruption of melatonin, a hormone that is produced at night and possesses anticancer properties. The World Health Organization has classified night shift work as a probable carcinogen highlighting evidence of an increased incidence of breast cancer among female shift workers. Research has demonstrated that light exposure at night can suppress melatonin production, potentially hindering the body's cancer-fighting capabilities (Haus & Smolensky, 2006). Additionally, male shift workers have been observed to have a higher risk of developing prostate cancer. Factors such as circadian rhythm disruption, hormonal imbalances and decreased daylight exposure are believed to contribute to this elevated risk (Schernhammer et al., 2001).

To lessen these risks, both individuals and organizations can implement various strategies and preventive measures. Shift workers are encouraged to maintain a regular sleep schedule, even on days off, to help stabilize their circadian rhythms. Utilizing blackout curtains or eye masks can create a dark sleeping environment, signaling to the body that it is time to rest. Since light exposure can inhibit melatonin production, it is essential for shift workers to manage their light environment effectively. Consuming small, balanced meals throughout the day and night can help maintain stable energy levels. Engaging in physical activity may alleviate some adverse effects of shift work, including lowering the risk of cardiovascular diseases, diabetes, and obesity. Techniques such as meditation, deep breathing exercises and progressive muscle relaxation can assist shift workers in managing stress and anxiety, particularly before sleep or during breaks, thereby enhancing mental well-being.

CONCLUSION

The health challenges associated with shift work are extensive and complex, affecting nearly every aspect of both mental and physical health. Among the sleep disorders linked to this work pattern is shift work sleep disorder (SWSD) which manifests as excessive sleepiness, fatigue and insomnia, alongside delayed sleep phase syndrome, both of which are tied to chronic disruptions in circadian rhythms. Beyond physical health implications shift work significantly impacts mental well-being, with higher rates of depression, anxiety, and cognitive impairments observed in shift workers compared to those on regular daytime schedules. These mental health issues are often exacerbated by the social isolation and lifestyle changes that many shift workers experience. Moreover, while further research is necessary to confirm definitive connections. Emerging studies suggest that prolonged exposure to shift work may increase the risk of certain cancers, such as breast and prostate cancer, due to hormonal disruptions and circadian rhythm disturbances. The primary factor contributing to many of these health concerns is the disruption of the circadian rhythm. Due to the prevalence of shift work across various sectors, it is essential for employers and healthcare professionals to recognize these risks and develop strategies to decrease them. Shift workers can improve their health by prioritizing good sleep practices, maintaining a nutritious diet, engaging in regular physical activity, managing stress and utilizing technology to monitor their health. Additionally, organizations should proactively design shift schedules that minimize adverse health effects and provide support services. To fully understand the extent of these health risks and explore effective prevention methods, further research is also warranted.

Future prospects of research on health issues related to shift work

Shift work is associated with various health issues, yet there remains a need for further investigation in several key areas. Given its widespread occurrence in sectors such as healthcare, transportation and manufacturing, it is essential to comprehend the long-term implications of shift work and to devise effective interventions. This section explores future research directions concerning the health impacts of shift work, emphasizing areas that require innovative strategies and deeper exploration. One of the most critical yet insufficiently studied effects of shift work is its impact on mental health. Additional research is necessary to clarify how disruptions to circadian rhythms influence different biological systems and lead to health problems. There is an increasing demand for evidence-based regulations and policies aimed at decreasing the health risks associated with shift work, particularly in high-risk fields such as emergency services, healthcare and transportation.

REFERENCES

1. Akerstedt, T. Shift work and health: A critical review of the literature. *Scandinavian Journal of Work, Environment & Health*, 2003; 29(4): 233-238.
2. Arendt, J. Shift work: Health, sleep, and performance. *Occupational Medicine*, 2010; 60(1): 10-18.
3. Bowers, A., & Moyer, A. Effects of shift work on sleep quality and mood disorders in hospital nurses: A meta-analysis. *Journal of Occupational Health Psychology*, 2017; 22(1): 10-24.
4. Cajochen, C., Frey, S., Anders, D., & Brunner, D. P. The effects of light on alertness and performance in shift workers: A review. *Chronobiology International*, 2011; 28(8): 669-680.
5. Haus, E., & Smolensky, M. H. Biological clocks and shift work: Circadian dysregulation and the risk of cardiovascular disease. *Chronobiology International*, 2006; 23(6): 1155-1169.
6. Kivimaki, M., Virtanen, M., Vahtera, J., Elovainio, M., & Keltikangas-Järvinen, L. Long working hours and risk of coronary heart disease and well-being: A systematic review and meta-analysis. *Scandinavian Journal of Work, Environment & Health*, 2011; 37(4): 245-252.
7. Pan, A., Schernhammer, E. S., Sun, Q., & Hu, F. B. Rotating night shift work and risk of type 2 diabetes: Two prospective cohort studies in women. PLoS Medicine, 2011; 8(12): e1001141.
8. Puttonen, S., Harma, M., & Hublin, C. Shift work and cardiovascular disease—A systematic review. *Scandinavian Journal of Work, Environment & Health*, 2010; 36(3): 163-179.
9. Saper B.Clifford, Thomas E. Scammell & Jun Lu. Hypothalamic regulation of sleep and circadian rhythms. Nature, 2005; 437: 1257–1263.
10. Schernhammer, E. S., Laden, F., Speizer, F. E., Willett, W. C., & Colditz, G. A. Rotating night shifts and risk of breast cancer in women participating in the Nurses' Health Study. *Journal of the National Cancer Institute*, 2001; 93(20): 1563-1568.
11. Vetter, C., Devore, E. E., & Czeisler, C. A. Shift work and the risk of cardiovascular disease. Current Cardiology Reports, 2016; 18(12): 1-9.