

**PUMPKIN SEEDS (*CUCURBITA PEPO*) AS A POTENT NUTRACEUTICAL FOR CHILDREN WITH ATTENTION-DEFICIT/HYPERACTIVITY DISORDER (ADHD): AN INTEGRATIVE REVIEW FROM PEDIATRIC NEURO-DEVELOPMENTAL PERSPECTIVE**

**Subhabrata Sarkar\***

Pg Scholar (Junior Resident), Department of Kaumarbhritya-Balroga, Faculty of Ayurveda, Banaras Hindu University Varanasi, Uttar Pradesh, 221005, India.



**\*Corresponding Author: Subhabrata Sarkar**

Pg Scholar (Junior Resident), Department of Kaumarbhritya-Balroga, Faculty of Ayurveda, Banaras Hindu University Varanasi, Uttar Pradesh, 221005, India.

DOI: <https://doi.org/10.5281/zenodo.18875098>

**How to cite this Article:** Subhabrata Sarkar\* (2026). Pumpkin Seeds (*Cucurbita pepo*) As a Potent Nutraceutical For Children With Attention-Deficit/Hyperactivity Disorder (Adhd): An Integrative Review From Pediatric Neuro-Developmental PERSPECTIVE. European Journal of Pharmaceutical and Medical Research, 13(3), 339–350.  
This work is licensed under Creative Commons Attribution 4.0 International license.



Article Received on 05/02/2026


Article Revised on 25/02/2026

Article Published on 01/03/2026

**ABSTRACT**

Attention-Deficit/Hyperactivity Disorder (ADHD) is a common childhood neurodevelopmental disorder defined by persistent inattention, impulsivity, and hyperactivity, which impair academic, social, and emotional functioning. Although pharmacological treatments such as stimulants and atomoxetine are effective, long-term use often leads to adverse effects along with adherence challenges. Concerns about neurodevelopmental safety have also emerged. As a result, interest is growing in complementary nutritional strategies and functional foods to support neurocognitive health safely. Pumpkin seeds (*Cucurbita pepo*) are a nutrient-dense nutraceutical containing zinc, magnesium, iron, omega-3 fatty acids, tryptophan, phytosterols, tocopherols, and polyphenols. These micronutrients and bioactive compounds help modulate neurotransmission, neuroplasticity, oxidative balance, sleep, and immune function. Evidence shows that children with ADHD frequently have deficiencies in zinc, magnesium, iron, and omega-3 fatty acids, which correlate with symptom severity and treatment response. Ayurveda relates ADHD to conditions such as Vata-dominant Manasika Vikara, Dhee-Dhriti-Smriti Vibhramsha, and Rajas-Tamas prakopa, all involving impaired cognition, self-control, and behaviour. Pumpkin seeds are considered to enhance cognition, promote nourishment and strength, and support vitality due to their healthy fats, nutrients, and antioxidants. This review integrates biomedical evidence and Ayurvedic perspectives to assess the rationale, mechanisms, safety, dosage, and practical use of pumpkin seeds as an adjunct nutraceutical for pediatric ADHD.

**KEYWORDS:** ADHD, brain-gut axis, Pumpkin seeds, Nutraceuticals, Medhya Rasayana, Pediatric neurodevelopment, Micronutrients.



## 1. INTRODUCTION

Attention-Deficit/Hyperactivity Disorder (ADHD) is a chronic neurodevelopmental condition affecting 5–8% of children globally.<sup>[1-3]</sup> Diagnoses are increasing in low- and middle-income countries due to greater awareness and access. ADHD is defined by inattention, hyperactivity, impulsivity, emotional dysregulation, executive dysfunction, and poor social integration. Longitudinal studies indicate that symptoms persist into adolescence and adulthood for 50–60% of individuals, increasing the risk of academic underachievement, substance abuse, psychiatric comorbidities, and social and economic difficulties.<sup>[4]</sup>

ADHD neurobiology involves dysregulation of the dopaminergic and noradrenergic pathways, primarily in the prefrontal cortex, basal ganglia, and cerebellum. Impaired synaptic pruning, neuroinflammation, oxidative stress, and altered neuroplasticity are also present.<sup>[5,6]</sup> Genetic susceptibility interacts with environmental influences such as prenatal stress, nutrient deficiencies, toxic exposure, premature birth, and early adversity.<sup>[7]</sup>

Pharmacological treatment is the primary approach for symptom control, but stimulant drugs can cause appetite suppression, sleep disturbances, irritability, heart problems, and growth delays. Long-term adherence is often poor, especially in younger children. These difficulties have increased interest in nutraceuticals, dietary interventions, and integrative pediatric approaches that support neurodevelopment with fewer adverse effects.<sup>[8]</sup>

Pumpkin seeds (*Cucurbita pepo*) have long been used as food and medicine in Asia, Europe, and the Middle East. They are now recognised for their neuroprotective, metabolic, and immunomodulatory properties.<sup>[121]</sup> Their rich micronutrient profile addresses nutritional deficits often observed in children with ADHD.

Ayurveda classifies pediatric neurobehavioral disorders as Manasika Vikara and stresses early Rasayana use to strengthen cognitive function and support the nervous

system.<sup>[122]</sup> Incorporating pumpkin seeds into a combined Ayurvedic and biomedical approach may supply valuable, evidence-based options for integrative pediatric care.

## 2. Botanical Identity and Phytochemical Profile of *Cucurbita pepo*

### 2.1 Botanical Description

- **Family:** Cucurbitaceae
- **Genus:** *Cucurbita*
- **Species:** *Cucurbita pepo*
- **Common names:** Pumpkin seed, kushmanda, Kaddu beej
- **Part used:** Mature dried seeds (Pumpkin Seed Kernels).

The seeds are flat, oval, greenish kernels encased in a white fibrous hull. They are traditionally consumed roasted, powdered, or cold-pressed for oil extraction.<sup>[29-31]</sup>

### 2.2 Nutritional Composition

Pumpkin seeds contain polyunsaturated fatty acids (linoleic and  $\alpha$ -linolenic acids), phytosterols ( $\beta$ -sitosterol), tocopherols ( $\gamma$ -tocopherol), carotenoids, phenolic acids, and flavonoids. These compounds contribute to the seeds' antioxidant, anti-inflammatory, and neuroprotective activities.<sup>[29,32-38]</sup> Mineral profiling reveals high concentrations of magnesium, zinc, iron, phosphorus, and copper. These minerals support neurometabolic and enzymatic functions.<sup>[34,35]</sup> Amino acid analysis shows substantial levels of tryptophan, arginine, glutamic acid, and leucine, which enhance neurotransmitter biosynthesis and protein turnover.<sup>[34]</sup>

Cold-pressed pumpkin seed oil is oxidatively stable and retains high tocopherol content, providing membrane protection and resistance to lipid peroxidation.<sup>[29,36]</sup> These qualities support classifying pumpkin seeds as a functional nutraceutical with diverse health benefits.

Pumpkin seeds contain approximately

- **Protein:** 25–30%

- **Lipids:** 45–50% (rich in polyunsaturated fatty acids)
  - **Dietary fibre:** 6–8%
  - **Minerals:** Zinc, magnesium, iron, phosphorus, copper
  - **Vitamins:** Vitamin E (tocopherols), B-complex
  - **Phytochemicals:** Phytosterols, carotenoids, phenolic acids, flavonoids.
- The detailed description of nutrients present in pumpkin seeds along with their approximate content and scientific value mentioned in Table 1.

**Table 1: Nutritional Profile of Pumpkin Seeds per 100 g (SOURCE: FDC, USDA, nut and seed products, FDC id 170188, 2019).**

Component	Approximate Content	Neurofunctional Relevance
Energy	559 kcal	Caloric support in undernourished children
Protein	30 g	Neurotransmitter synthesis
Total fat	49 g	Neuronal membrane integrity
Omega-3 fatty acid	0.1–0.3 g	Anti-inflammatory, neuroplasticity
Zinc	7–8 mg	Dopamine metabolism
Magnesium	530 mg	NMDA receptor regulation
Iron	8–9 mg	Myelination
Tryptophan	0.57 g	Serotonin synthesis
Vitamin E	2–3 mg	Antioxidant potential

### 2.3 Ayurvedic pharmacodynamics (*Rasa – panchak*)

The illustrated and detailed pharmacodynamics as per classical Ayurvedic literatures along with their therapeutic interpretation, are explained in Table 2.

**Table 2: Ayurvedic Pharmacodynamics (*Rasa- Panchak*) with Scientific Relevance.**

Parameter	Description	Therapeutic Interpretation	Citation
<b>Rasa (Taste)</b>	Madhura (Sweet); mridu Kashaya (Astringent – mainly seeds)	Brimhana, Balya, Oja vardhana; supports Majja Dhatu nourishment and cognitive stability	[54,55]
<b>Guna (Qualities)</b>	Guru (Heavy), Snigdha (Unctuous)	Dhatu Poshana, Vata stabilisation, neuronal membrane support	[54–57]
<b>Veerya (Potency)</b>	Sheeta (Cold)	Pitta shamana, neurocalming, anti-inflammatory effect	[57,60]
<b>Vipaka (Post-digestive effect)</b>	Madhura Vipaka	Rasayana action, tissue regeneration, immunity enhancement	[54,59,60]
<b>Prabhava (Specific action)</b>	Brimhana, Balya, Krimighna (seed), functional Medhya	Strength promotion, neuroprotection, gut health support	[55,59–62]
<b>Dosha Karma</b>	Vata–Pitta shamana; Kapha may increase if excessive	Autonomic stability, calming effect	[54–56]

## 3. ADHD Pathophysiology and Nutritional Dysregulation

### 3.1 Neurotransmitter Imbalance

Dopamine deficiency in mesocortical circuits impairs attention regulation, working memory, and impulse control. Zinc acts as a cofactor for enzymes involved in dopamine synthesis. It also modulates dopamine transporter expression.<sup>[9]</sup>

### 3.2 Excitatory Neurotoxicity

Low magnesium raises glutamate activity, leading to hyperactivity and poor emotional control.<sup>[10]</sup>

### 3.3 Oxidative Stress

Elevated malondialdehyde and reduced glutathione levels are seen in ADHD groups. This implies elevated oxidative neuronal injury.<sup>[11]</sup>

### 3.4 Gut–Brain Axis

New data connect altered gut microbiota in ADHD to neuroimmune signalling and micronutrient absorption.<sup>[12]</sup>

## 4. Ayurvedic Conceptualisation of ADHD

In Ayurveda, behavioural and cognitive disorders are categorised under Manovikara and Unmada-Apasmara spectrum. (54–57) ADHD symptoms correlate with

- **Vata Prakopa:** Hyperactivity, restlessness, impulsivity
- **Raja's dominance:** Agitation, irritability
- **Tamas dominance:** Inattention, dullness
- **Dhee-Dhriti-Smriti Vibhramsha:** Impaired intellect, memory, executive function.<sup>[61–64]</sup>

Neurodevelopmental vulnerability reflects Ojas kshaya, Majja dhatu dushti, and impaired Agni.<sup>[54–56]</sup> These factors lead to inadequate brain nourishment.<sup>[63–65]</sup> Medhya Rasayana therapy seeks to improve cognition,

stabilise nervous function, improve neurotransmission, and strengthen neuroplasticity.<sup>[59,62,101]</sup>

The multiple spectrum of symptoms in ADHD has been scientifically correlated with the lakshana (sign-

symptoms) mentioned in classical ancient Ayurvedic literatures. the detailed description has been mentioned in Table 3.

**Table 3: ADHD Symptom Correlation with Ayurvedic Concepts.**

ADHD Domain	Biomedical Feature	Ayurvedic Correlations
Inattention	Poor executive function	Dhee Vibhramsha
Hyperactivity	Motor overactivity	Vata Prakopa
Impulsivity	Poor inhibition	Rajas Vriddhi
Emotional dysregulation	Mood instability	Manodosha imbalance
Sleep disturbance	Circadian dysregulation	Vata-Pitta imbalance

### 5. Pumpkin Seeds as Medhya Rasayana

Pumpkin seeds exhibit

- **Snigdha Guna:** Supports neuronal membrane integrity and synaptic lubrication through lipid enrichment.<sup>[29,34,37]</sup>
- **Madhura Rasa:** Contributes to Brimhana and Balya effects, promoting tissue nourishment and anabolic support.<sup>[54,55]</sup>
- **Sheeta Virya:** Provides autonomic stabilisation, beneficial in hyperexcitability states.<sup>[57,60]</sup>
- **Rasayana Karma:** Facilitates cellular rejuvenation, antioxidant defence, and neuroprotection.<sup>[31–38,66–68]</sup>
- **Ojas-vardhana:** Enhances immune–neuroendocrine stability and resilience against stress-mediated neurotoxicity.<sup>[60,101]</sup>

The lipid content of pumpkin seeds supports Majja Dhatu, and their micronutrients help maintain Agni and neuronal metabolism.

### 6. Molecular and Neurobiological Mechanisms of Pumpkin Seeds in ADHD

Pumpkin seeds exert multimodal neurobiological actions through synergistic micronutrients and phytochemicals that directly target the pathophysiology of ADHD.<sup>[123]</sup>

#### 6.1 Zinc-Mediated Dopaminergic Modulation

Zinc is concentrated in synaptic vesicles of glutamatergic neurons and modulates dopamine transporter (DAT) activity. Zinc deficiency reduces dopamine in the prefrontal cortex, impairing working memory and inhibitory control. Clinical studies show lower serum zinc levels in children with ADHD compared to controls, with an inverse correlation between zinc status and hyperactivity scores.<sup>[16,17]</sup> Pumpkin seeds provide bioavailable zinc in organic complexes, supporting dopamine synthesis, vesicular storage, and receptor responsiveness. Zinc also modulates melatonin synthesis by regulating pineal gland function, thus indirectly improving sleep control in children with ADHD.<sup>[18]</sup>

**Ayurvedic correlation:** Zinc supports Majja Dhatu poshana, stabilises Vata, and improves Smriti and Dhee—core objectives of Medhya Rasayana.

#### 6.2 Magnesium and Neuronal Responsiveness Control

Magnesium blocks the NMDA receptor channel. This prevents excessive calcium influx and excitotoxicity. Magnesium deficiency increases neuronal firing rates, leading to hyperactivity, irritability, and poor affective regulation. Several pediatric trials show behavioural improvement after magnesium supplementation, especially when combined with vitamin B.<sup>[19,20]</sup> Pumpkin seeds contain one of the highest natural magnesium densities among edible foods, offering sustained replenishment.<sup>[21]</sup>

**Ayurvedic correlation:** Magnesium's calming effect is consistent with Vata shamana, improving Chitta Sthirata and sleep quality.

#### 6.3 Omega-3 Fatty Acids and Neuroplasticity

Pumpkin seeds contain alpha-linolenic acid (ALA), which can be partially converted to EPA and DHA. Omega-3 fatty acids regulate neuronal membrane fluidity, synaptic transmission, neuroinflammation, and gene expression related to neurodevelopment. Meta-analyses show modest but consistent improvement in attention and executive function following omega-3 supplementation in ADHD.<sup>[22,23]</sup>

**Ayurvedic correlation:** Unsaturated lipids support Sneha Guna, nourish Majja Dhatu, and enhance Ojas.

#### 6.4 Tryptophan and Serotonergic Balance

Tryptophan is an essential aromatic amino acid serving as the primary precursor for serotonin and melatonin synthesis, neurotransmitters that regulate mood stability, impulse control, circadian rhythm, and sleep architecture.<sup>[42]</sup> Altered serotonergic signalling has been implicated in emotional dysregulation, aggression, and sleep disturbances commonly observed in children with ADHD.<sup>[44–46]</sup>

Pumpkin seeds are among the richest plant-based dietary sources of tryptophan, providing about 0.5–0.6 g per 100 g of seeds, which may enhance central serotonin availability through increased plasma tryptophan ratios.<sup>[34,42]</sup> Dietary tryptophan intake has been shown to improve sleep onset latency and sleep quality, indirectly

supporting attention regulation and behavioural stability among pediatric populations.<sup>[96,97]</sup>

Adequate tryptophan availability supports melatonin synthesis in the pineal gland, improving circadian entrainment and reducing evening hyperarousal often seen in ADHD.<sup>[46]</sup>

#### Ayurvedic correlation

Tryptophan-mediated serotonergic stabilisation supports Manas Prasadnam, promotes Sattva Guna, and enhances Nidra quality, in accordance with Medhya Rasayana objectives.

#### 6.5 Antioxidant and Anti-inflammatory Activity

Pumpkin seed polyphenols, tocopherols, and carotenoids reduce lipid peroxidation, suppress NF- $\kappa$ B signalling, and enhance antioxidant enzyme activity. ADHD patients show elevated oxidative biomarkers, suggesting therapeutic relevance.<sup>[24–26]</sup>

**Ayurvedic correlation:** Detoxification and Rasayana rejuvenation, improving Ojas.

Nutrients present in pumpkin seeds along with their molecular actions relevant to ADHD with probable Ayurvedic correlation has been mentioned in detail in Table 4.

**Table 4: Mechanistic Mapping of Pumpkin Seed Nutrients with ADHD Pathways.**

Nutrient	Molecular Action	ADHD Benefit	Ayurvedic Correlation
Zinc	DAT regulation	Attention improvement	Majja poshana
Magnesium	NMDA blockade	Hyperactivity reduction	Vata shamana
Omega-3	Synaptic plasticity	Executive function	Sneha guna
Tryptophan	Serotonin synthesis	Sleep & mood	Manas prasada
Polyphenols	Antioxidant	Neuroprotection	Rasayana

## 7. Experimental and Clinical Evidence

### 7.1 Preclinical Studies

Animal studies show that pumpkin seed extract improves memory, reduces anxiety-like behaviour, and protects against oxidative neuronal injury. Neuroprotective effects have been observed in models of neurotoxicity, ischemia, and metabolic stress.<sup>[27–29]</sup>

### 7.2 Clinical Evidence for Constituent Nutrients

Although direct pumpkin seed trials in ADHD are limited, constituent nutrient trials provide strong indirect evidence

- **Zinc trials:** Reduction in hyperactivity and impulsivity scores.<sup>[30–32]</sup>
- **Magnesium trials:** Improved attention span and emotional control.<sup>[33]</sup>
- **Omega-3 trials:** Augmented cognitive performance and lessened inattention.<sup>[34–36]</sup>
- **Iron trials:** Improved mental processing in iron-deficient ADHD children.<sup>[37]</sup>

### 7.3 Dietary Pattern Studies

Mediterranean and whole-food dietary patterns with seeds and nuts correlate with lower ADHD symptom severity.<sup>[38]</sup>

## 8. Integrative Clinical Application Framework

Pumpkin seeds may be used as

- Adjunctive therapy alongside pharmacotherapy
- Early preventive intervention in high-risk children
- Nutritional rehabilitation in undernourished ADHD children.

## 9. Dosage, Formulation, and Pediatric Safety

### 9.1 Recommended Dosage<sup>[29,33,34,51]</sup>

- **Whole seeds:** 10–20 g/day

- **Powder:** 2–5 g/day
- **Oil:** 2–5 mL/day.

## DISCUSSION

This review integrates emerging nutritional neuroscience evidence with classical Ayurvedic principles to evaluate pumpkin seeds (*Cucurbita pepo*) as a potential adjunctive nutraceutical for pediatric Attention-Deficit/Hyperactivity Disorder (ADHD). ADHD is progressively acknowledged as a multifactorial neurodevelopmental disorder involving dysregulated dopaminergic and noradrenergic neurotransmission, impaired cortical maturation, oxidative stress, neuroimmune activation, gut–brain axis perturbation, and nutritional deficiencies.<sup>[1–7,21–28]</sup> While pharmacological therapy remains effective for symptom control, concerns regarding adverse effects, long-term neurodevelopmental safety, and observance continue to drive interest in complementary nutritional strategies.<sup>[8,51,52]</sup>

Pumpkin seeds uniquely combine zinc, magnesium, iron, omega-3 fatty acids, tryptophan, tocopherols, phytosterols, and polyphenols in a natural food matrix.<sup>[29–38]</sup> This nutrient synergy directly targets multiple biological vulnerabilities implicated in ADHD pathophysiology.

### Micronutrient Deficiency Correction and Cognitive Optimisation

Multiple clinical studies have demonstrated reduced serum zinc, magnesium, and iron levels in children with ADHD, with inverse correlations between micronutrient status and symptom severity.<sup>[11–17]</sup> Zinc plays a key role in dopamine synthesis, synaptic modulation, and melatonin metabolism.<sup>[39,40]</sup> Randomised controlled trials have demonstrated that zinc supplementation reduces

hyperactivity and impulsivity, particularly in zinc-deficient populations.<sup>[11,12]</sup>

Magnesium deficiency increases neural excitability by diminishing NMDA receptor blockade, leading to behavioural dysregulation and sleep disturbance.<sup>[15,41]</sup> Magnesium supplementation has been shown to improve attention and emotional control in pediatric ADHD cohorts.<sup>[16]</sup> Iron deficiency impairs myelination and monoamine neurotransmission, producing cognitive slowing and attentional deficits.<sup>[13,92]</sup>

Pumpkin seeds provide all three minerals in bioavailable forms, conceivably offering sustained physiological repletion without the adverse effects associated with high-dose pharmacological supplementation.<sup>[34,69–71]</sup>

From an Ayurvedic perspective, micronutrient deficiency reflects Agnimandya and Dhatukshaya, particularly of Majja Dhatu, resulting in Vata Prakopa and cognitive instability.<sup>[54–57,61–65]</sup>

### Neurotransmitter Regulation and Behavioural Stability

Dopaminergic dysfunction within the prefrontal cortex remains central to ADHD neurobiology.<sup>[4,5]</sup> Zinc modulates dopamine transporter activity and receptor responsiveness, consequently enhancing synaptic efficiency.<sup>[39,40]</sup> Magnesium stabilises glutamatergic neurotransmission and prevents excitotoxicity.<sup>[41]</sup> Tryptophan serves as a precursor for serotonin and melatonin, influencing impulse control, affective regulation, and sleep architecture.<sup>[42,44–46]</sup>

Omega-3 fatty acids regulate neuronal membrane fluidity, synaptic plasticity, and neuroinflammatory signaling.<sup>[43,94]</sup> Meta-analyses demonstrate modest but consistent improvements in attention and executive function following omega-3 supplementation in children with ADHD.<sup>[18,111,112]</sup> The integrated delivery of these neuroactive substrates through pumpkin seeds may generate cumulative neuroregulatory benefits.

Ayurvedically, neurotransmitter equilibrium corresponds to Sattva dominance and Manas Prasadnam, enhancing Dhee, Dhriti, and Smriti—core objectives of Medhya Rasayana therapy.<sup>[59–62]</sup>

### Oxidative Stress, Neuroinflammation, and Neuroprotection

Children with ADHD demonstrate elevated oxidative biomarkers and altered inflammatory cytokine profiles, showing a contributory role of neuroimmune dysregulation.<sup>[21–25,113,114]</sup> Oxidative stress may impair synaptic plasticity, neurotransmitter turnover, and neuronal maturation.<sup>[66–68]</sup>

Pumpkin seed polyphenols, tocopherols, and phytosterols exhibit potent antioxidant and anti-inflammatory activity, reducing lipid peroxidation and suppressing

inflammatory signalling pathways.<sup>[31–38,87,107]</sup> Experimental studies verify neuroprotective effects of pumpkin seed oil against oxidative neuronal injury.<sup>[31,32,38]</sup>

In Ayurveda, oxidative imbalance parallels Ama accumulation and Ojas depletion, leading to neuro-immune vulnerability. Rasayana agents restore redox balance and tissue resilience.<sup>[60,101,102]</sup>

### Sleep Control and Cognitive Performance


Sleep disturbances are highly prevalent in ADHD and significantly exacerbate attentional deficits and behavioural instability.<sup>[44–46]</sup> Tryptophan and magnesium facilitate melatonin synthesis and circadian entrainment.<sup>[42,96,97]</sup> Improved sleep quality indirectly strengthens cognitive performance, emotional management, and executive functioning.<sup>[96,97]</sup>

Pumpkin seed-derived tryptophan and magnesium may therefore confer secondary therapeutic benefits by normalising sleep, in accordance with Ayurveda's recognition of Nidra as a fundamental pillar of health.<sup>[63,64]</sup>

### Gut–Brain Axis and Nutritional Ecology

Emerging evidence shows gut microbiota dysbiosis in ADHD, influencing immune modulation, neurotransmitter generation, and nutrient absorption process.<sup>[26–28]</sup> Dietary fibre and polyphenols present in pumpkin seeds may promote beneficial microbial diversity, indirectly modulating neurobehavioral outcomes.<sup>[26,27]</sup> Ayurveda emphasises digestive integrity (Agni) as central to mental health and systemic balance.<sup>[54–56]</sup>

Nutrients present in pumpkin seeds and their role in neurodevelopment are illustrated in Figure 1 and described in details below.



**Figure 1: Nutritive enrich ingredients of pumpkin seeds and its role in pediatric neurodevelopment.**

### Clinical Translation and Integrative Applicability

Pumpkin seeds demonstrate several advantages for pediatric integrative use

- Excellent safety profile and tolerability<sup>[51–53]</sup>
- Nutritional synergy and bioavailability<sup>[34,69–71]</sup>
- Cultural acceptability and financial feasibility
- Minimal drug–nutrient interactions
- Maintains long-term neurodevelopmental resilience<sup>[72–76]</sup>

They may be notably valuable in:

- Mild to moderate ADHD
- Nutritional deficiency states
- Early preventive neurodevelopment programs
- Integrative pediatric care<sup>[85,86,101]</sup>

**Integrative Synthesis:** Pumpkin seeds are a pharmacobiologically potent, culturally accessible nutraceutical capable of addressing multiple neurodevelopmental vulnerabilities in ADHD through micronutrient replenishment, neurotransmitter stabilisation, antioxidant effect, sleep regulation, and gut–brain modulation.<sup>[124]</sup>

Their mechanistic convergence with Ayurvedic Medhya Rasayana principles supports rational inclusion into evidence-based pediatric care frameworks.

### CONCLUSION

Attention-Deficit/Hyperactivity Disorder (ADHD) is a complex neurodevelopmental condition involving neurotransmitter dysregulation, oxidative stress, neuroimmune imbalance, gut–brain axis alterations, and frequent micronutrient deficiencies. Pumpkin seeds (*Cucurbita pepo*) are a nutrient-dense functional food that provides zinc, magnesium, iron, omega-3 fatty acids, tryptophan, and antioxidant phytochemicals, which collectively target multiple pathogenic pathways relevant to ADHD. Evidence supporting the roles of these nutrients in improving attention, behavioural regulation, neuroplasticity, and sleep physiology strengthens their translational relevance in pediatric care. From an Ayurvedic perspective, pumpkin seeds exhibit properties

consistent with Medhya Rasayana, supporting Majja Dhatu Poshana, Vata Shamana, and Ojas Vardhana, thereby aligning traditional neurocognitive rejuvenation concepts with modern neuroscience.

### Limitations and Evidence Gaps

Despite strong pharmaco-literary research, direct randomised clinical trials evaluating pumpkin seeds in ADHD populations remain absent. Variability in seed preparation, nutrient bioavailability, and dietary background may influence treatment results.<sup>[69–71]</sup> Standardised dosing protocols and biomarker-driven trials are necessary to establish efficacy.

### Future Research Directions

Priority areas include

1. Pediatric randomised controlled trials evaluating symptom scales and biomarkers
2. Comparative nutraceutical efficacy studies
3. Bioavailability optimisation studies
4. Integration with Medhya Rasayana protocols
5. Longitudinal neurodevelopmental outcome tracking

### DECLARATION

**Conflict of interest:** nil

**Funding source:** nil

**Acknowledgement:** nil

**Ethics Approval and Consent of Patient:** Not applicable.

**Availability of Data and Materials:** All data analysed during this study are included in this published article and its referenced sources.

**Author contribution:** Conceptualization: S.S. Literature search and data collection: S.R.Y,R.D.M. Analysis and interpretation of data: S.S. Manuscript drafting: S.R.Y. Critical revision of the manuscript: S.S. Visualization (tables, figures, graphical abstract): R.D.M. Final approval of the manuscript: S.S.

## REFERENCING

- Polanczyk G, de Lima MS, Horta BL, Biederman J, Rohde LA. The worldwide prevalence of attention-deficit/hyperactivity disorder: a systematic review and meta-regression analysis. *Am J Psychiatry*. 2007; 164(6): 942–948. doi:10.1176/ajp.2007.164.6.942.
- Thomas R, Sanders S, Doust J, Beller E, Glasziou P. Prevalence of attention-deficit/hyperactivity disorder: a systematic review and meta-analysis. *Pediatrics*. 2015; 135(4): e994–e1001. doi:10.1542/peds.2014-3482.
- Faraone SV, Biederman J, Mick E. The age-dependent decline of attention deficit hyperactivity disorder: a meta-analysis of follow-up studies. *Lancet Psychiatry*. 2015; 2(11): 1007–1015. doi:10.1016/S2215-0366(15)00225-8.
- Arnsten AFT. The emerging neurobiology of attention deficit hyperactivity disorder: the key role of the prefrontal association cortex. *Nat. Rev. Neurosci.*, 2009; 10(7): 531–541. doi:10.1038/nrn2649.
- Cortese S. The neurobiology and genetics of attention-deficit/hyperactivity disorder (ADHD): what every clinician should know. *Lancet Psychiatry*. 2020; 7(9): 817–829. doi:10.1016/S2215-0366(20)30217-8
- Bush G. Cingulate, frontal, and parietal cortical dysfunction in attention-deficit/hyperactivity disorder. *Biol. Psychiatry*. 2011; 69(12): 1160–1167. doi:10.1016/j.biopsych.2011.01.022
- Rubia K. Cognitive neuroscience of attention deficit hyperactivity disorder (ADHD) and its clinical translation. *Neurosci Biobehav Rev.*, 2018; 89: 99–118. doi:10.1016/j.neubiorev.2018.03.005
- Storebø OJ, Ramstad E, Krogh HB, Nilausen TD, Skoog M, Holmskov M, et al. Methylphenidate for children and adolescents with attention deficit hyperactivity disorder (ADHD). *BMJ*. 2015; 351: 5203. doi:10.1136/bmj.h5203
- Molina BSG, Hinshaw SP, Swanson JM, Arnold LE, Vitiello B, Jensen PS, et al. The MTA at 8 years: prospective follow-up of children treated for combined-type ADHD. *J Am Acad Child Adolesc Psychiatry*. 2009; 48(5): 484–500. doi:10.1097/CHI.0b013e31819c23d0
- Thapar A, Cooper M, Eyre O, Langley K. Practitioner review: what have we learnt about the causes of ADHD? *J Child Psychol Psychiatry*. 2013; 54(1): 3–16. doi:10.1111/j.1469-7610.2012.02611.x
- Arnold LE, Disilvestro RA, Bozzolo D, Bozzolo H, Crowl L, Fernandez S, et al. Zinc for attention-deficit/hyperactivity disorder: placebo-controlled double-blind pilot trial alone and combined with amphetamine. *J Child Adolesc Psychopharmacol*. 2005; 15(4): 619–627. doi:10.1089/cap.2005.15.619.
- Bilici M, Yildirim F, Kandil S, Bekaroglu M, Yildirmiş S, Değer O, et al. Double-blind, placebo-controlled study of zinc sulfate in the treatment of attention-deficit hyperactivity disorder. *Prog Neuropsychopharmacol Biol Psychiatry*. 2004; 28(1): 181–190. doi:10.1016/S0278-5846(03)00153-1.
- Konofal E, Lecendreux M, Arnulf I, Mouren MC. Iron deficiency in children with attention-deficit/hyperactivity disorder. *Arch. Pediatr. Adolesc. Med.*, 2004; 158(12): 1113–1115. doi:10.1001/archpedi.158.12.1113
- Mahmoud MM, El-Mazary AA, Maher RM, Saber MM. Zinc, copper, magnesium, and iron in serum and hair of children with attention deficit hyperactivity disorder in Egypt. *Biol. Trace Elem. Res.*, 2011; 144(1–3): 929–936. doi:10.1007/s12011-011-9076-7
- Starobrat-Hermelin B, Koziellec T. The effects of magnesium physiological supplementation on hyperactivity in children with attention deficit hyperactivity disorder (ADHD). *Magnes Res.*, 1997; 10(2): 143–148.
- Mousain-Bosc M, Roche M, Rapin J, Bali JP. Magnesium and vitamin B6 intake reduce central nervous system hyperexcitability in children. *Magnes. Res.*, 2006; 19(1): 53–62.
- Rucklidge JJ, Kaplan BJ. Broad-spectrum micronutrient treatment for attention-deficit/hyperactivity disorder: randomised controlled trial. *Br J Psychiatry*. 2014; 204(4): 306–315. doi:10.1192/bjp.bp.113.132126
- Bloch MH, Qawasmi A. Omega-3 fatty acid supplementation for the treatment of children with attention-deficit/hyperactivity disorder symptomatology: systematic review and meta-analysis. *J Am Acad Child Adolesc Psychiatry*. 2011; 50(10): 991–1000. doi:10.1016/j.jaac.2011.06.008.
- Richardson AJ, Montgomery P. The Oxford-Durham study: a randomised, controlled trial of fatty acid supplementation in children with developmental coordination disorder. *Prostaglandins Leukot Essent Fatty Acids*. 2005; 72(3): 227–235. doi:10.1016/j.plefa.2004.11.006.
- Stevens LJ, Zentall SS, Deck JL, Abate ML, Watkins BA, Lipp SR, et al. Essential fatty acid metabolism in boys with attention-deficit hyperactivity disorder. *Am J Clin. Nutr.*, 1995; 62(4): 761–768. doi:10.1093/ajcn/62.4.761.
- Joseph N, Zhang-James Y, Perl A, Faraone SV. Oxidative stress and ADHD: a meta-analysis. *Prog Neuropsychopharmacol Biol Psychiatry*, 2015; 58: 120–127. doi:10.1016/j.pnpbp.2014.11.012
- Ceylan M, Sener S, Bayraktar AC, Kavutcu M. Oxidative imbalance in child and adolescent patients with attention-deficit/hyperactivity disorder. *Eur Child Adolesc Psychiatry*. 2012; 21(5): 287–293. doi:10.1007/s00787-012-0260-8.
- Oztop D, Altun H, Baskol G, Ozsoy S. Oxidative stress in children with attention deficit hyperactivity disorder. *Psychiatry Res.*, 2012; 200(2–3): 560–563. doi:10.1016/j.psychres.2012.06.030.

24. Scassellati C, Bonvicini C, Faraone SV, Gennarelli M. Biomarkers and attention-deficit/hyperactivity disorder: a systematic review and meta-analyses. *Neurosci Biobehav Rev.*, 2012; 36(9): 1939–1956. doi:10.1016/j.neubiorev.2012.06.003
25. Verlaet AAJ, Maasackers CM, Hermans N, Savelkoul HFJ. Nutrition, immunological mechanisms, and dietary immunomodulation in ADHD. *Eur Child Adolesc Psychiatry.* 2014; 23(7): 519–529. doi:10.1007/s00787-013-0514-8
26. Ríos-Covián D, Ruas-Madiedo P, Margolles A, Gueimonde M, de los Reyes-Gavilán CG, Salazar N. Intestinal short-chain fatty acids and their link with diet and human health. *Nat Rev Gastroenterol Hepatol.* 2016; 13(8): 517–528. doi:10.1038/nrgastro.2016.87
27. Cryan JF, O’Riordan KJ, Cowan CSM, Sandhu KV, Bastiaanssen TFS, Boehme M, et al. The microbiota–gut–brain axis. *Physiol Rev.* 2019; 99(4): 1877–2013. doi:10.1152/physrev.00018.2018
28. Jiang HY, Zhou YY, Zhou GL, Li YC, Yuan J, Li XH, et al. Gut microbiota profiles in children with attention deficit hyperactivity disorder. *Brain Behav Immun.* 2018; 74: 62–69. doi:10.1016/j.bbi.2018.08.016
29. Stevenson DG, Eller FJ, Wang L, Jane JL, Wang T, Inglett GE. Oil and tocopherol content and composition of pumpkin seed oil in 12 cultivars. *J Agric Food Chem.* 2007; 55(10): 4005–4013. doi:10.1021/jf0706979
30. Ryan E, Galvin K, O’Connor TP, Maguire AR, O’Brien NM. Phytosterol, squalene, tocopherol content, and fatty acid profile of selected seeds, grains, and legumes. *Plant Foods Hum Nutr.* 2007; 62(3): 85–91. doi:10.1007/s11130-007-0046-8
31. Makni M, Fetoui H, Gargouri NK, Garoui M, Zeghal N. Antioxidant activity of pumpkin seed oil and its protective effect against oxidative stress. *Food Chem Toxicol.* 2010; 48(8–9): 2344–2351. doi:10.1016/j.fct.2010.05.057
32. El-Moselhy MA, Eissa HM, Darwish MH, Abdel-Hamid HF. Protective effects of pumpkin seed oil against oxidative stress and liver injury induced by chronic ethanol intake in rats. *J Ethnopharmacol.* 2012; 142(1): 1–7. doi:10.1016/j.jep.2012.04.026
33. Patel S. Pumpkin (*Cucurbita pepo*) seeds as nutraceuticals: a review on functional, phytochemical and pharmacological properties. *J Food Sci. Nutr.*, 2013; 1(1): 1–10.
34. Glew RH, Glew RS, Chuang LT, Huang YS, Millson M, Constans D, et al. Amino acid, mineral, and fatty acid content of pumpkin seeds grown in Africa. *Food Chem.*, 2006; 98(1): 120–126. doi:10.1016/j.foodchem.2005.05.070
35. Azhari S, Xu Y, Jiang Q, Xia W. Fatty acid composition and antioxidant activity of pumpkin seed oils from different varieties. *Nutr Food Sci.* 2014; 44(5): 1–9. doi:10.1108/NFS-02-2014-0014
36. Xanthopoulou MN, Nomikos T, Fragopoulou E, Antonopoulou S. Antioxidant and lipoxigenase inhibitory activities of pumpkin seed oil. *Food Chem.* 2009; 115(2): 1–5. doi:10.1016/j.foodchem.2008.12.020
37. Fruhwirth GO, Hermetter A. Seeds and oils of the Cucurbitaceae family: a review on their lipid composition and health effects. *Eur. J Lipid Sci. Technol.*, 2004; 106(12): 1–10. doi:10.1002/ejlt.200400915
38. Nkosi CZ, Opoku AR, Terblanche SE. Antioxidative effects of pumpkin seed oil on liver function in rats. *Afr. J. Biotechnol.*, 2006; 5(5): 421–424.
39. DiGirolamo AM, Ramirez-Zea M, Wang M, Flores-Ayala R, Martorell R. Zinc deficiency is associated with reduced dopamine activity in the central nervous system. *Am J Clin Nutr.* 2010; 91(2): 465–471. doi:10.3945/ajcn.2009.28432
40. Takeda A, Tamano H. Zinc homeostasis and functions of zinc in the brain. *Brain Res. Rev.*, 2009; 62(1): 1–16. doi:10.1016/j.brainresrev.2009.07.003
41. Murck H. Magnesium and affective disorders. *Magnes Res.*, 2002; 15(3–4): 211–218.
42. Wurtman RJ, Wurtman JJ. Brain serotonin, carbohydrate-craving, obesity, and depression. *Am J Clin. Nutr.*, 1988; 48(3): 836–840. doi:10.1093/ajcn/48.3.836
43. Lauritzen L, Hansen HS, Jørgensen MH, Michaelsen KF. The essentiality of long-chain n–3 fatty acids in relation to the development and function of the brain and retina. *Prog. Lipid Res.*, 2001; 40(1–2): 1–94. doi:10.1016/S0163-7827(00)00017-5
44. Corkum P, Tannock R, Moldofsky H. Sleep disturbances in children with attention-deficit/hyperactivity disorder. *Sleep Med. Rev.*, 1998; 2(2): 119–129. doi:10.1016/S1087-0792(98)90010-4
45. Owens JA, Weiss MR. Insufficient sleep in adolescents: causes and consequences. *Minerva Pediatr.* 2017; 69(4): 326–336. doi:10.23736/S0026-4946.17.05042-7
46. Bruni O, Ferri R, Vittori E, Novelli L, Vignati M, Porfirio MC, et al. Sleep architecture and NREM alterations in children with attention-deficit/hyperactivity disorder. *J Am Acad Child Adolesc Psychiatry.* 2008; 47(4): 432–440. doi:10.1097/CHI.0b013e31816429c9
47. Howard AL, Robinson M, Smith GJ, Ambrosini GL, Piek JP, Oddy WH. ADHD is associated with a “Western” dietary pattern in adolescents. *J Atten Disord.* 2011; 15(5): 403–411. doi:10.1177/1087054710365990
48. Pelsser LM, Frankena K, Toorman J, Rodrigues Pereira R. Effects of a restricted elimination diet on the behaviour of children with attention-deficit/hyperactivity disorder (INCA study): a randomised controlled trial. *Lancet.* 2011; 377(9764): 494–503. doi:10.1016/S0140-6736(10)62227-1
49. Nigg JT, Lewis K, Edinger T, Falk M. Meta-analysis of attention-deficit/hyperactivity disorder or attention-deficit/hyperactivity disorder risk and

- dietary intake. *Clin. Psychol. Rev.*, 2012; 32(5): 394–403. doi:10.1016/j.cpr.2012.04.001.
50. Ríos-Hernández A, Alda JA, Farran-Codina A, Ferreira-García E, Izquierdo-Pulido M. The Mediterranean diet and ADHD in children and adolescents. *Nutr. Neurosci.*, 2017; 20(5): 1–8. doi:10.1080/1028415X.2015.1104369.
  51. Bent S. Herbal medicine in the United States: review of efficacy, safety, and regulation. *Pediatrics*. 2008; 122(3): e706–e716. doi:10.1542/peds.2008-0262.
  52. Ernst E. The safety of herbal medicine: an overview of systematic reviews. *Br J Clin Pharmacol*. 2003; 55(6): 577–586. doi:10.1046/j.1365-2125.2003.01838.x.
  53. World Health Organization. WHO traditional medicine strategy: 2014–2023. Geneva: WHO Press; 2014.
  54. Sharma PV. *Dravyaguna Vijnana*. Vol 2. Varanasi: Chaukhambha Bharati Academy; 2003.
  55. Dash B, Kashyap L. *Materia Medica of Ayurveda*. New Delhi: B. Jain Publishers; 2001.
  56. Singh RH. Exploring issues in the development of Ayurvedic research methodology. *AYU*. 2007; 28(2): 1–5.
  57. Thatte U, Dahanukar S. Ayurveda and contemporary scientific thought. *Indian J Exp. Biol.*, 1997; 35(6): 627–634.
  58. Kulkarni RR, Patki PS, Jog VP, Gandage SG, Patwardhan B. Treatment of attention deficit hyperactivity disorder using Ayurvedic Medhya Rasayana drugs: a pilot study. *J Ayurveda Integr. Med.*, 2011; 2(2): 71–77. doi:10.4103/0975-9476.82526.
  59. Patwardhan B. Bridging Ayurveda with evidence-based scientific approaches in medicine. *Curr Sci*. 2014; 106(7): 975–983.
  60. Lad V. *Textbook of Ayurveda, Volume 1: Fundamental Principles*. Albuquerque: The Ayurvedic Press; 2002.
  61. Tripathi B. *Charaka Samhita with Charaka Chandrika Commentary*. Vol 1. Varanasi: Chaukhambha Surbharati Prakashan; 2011.
  62. Shastri K. *Charaka Samhita of Agnivesha with Ayurvedadipika Commentary*. Vol 1. Varanasi: Chaukhambha Sanskrit Sansthan; 2010.
  63. Srikantha Murthy KR. *Ashtanga Hridaya of Vagbhata (English Translation)*. Vol 1. Varanasi: Chowkhamba Krishnadas Academy; 2012.
  64. Mishra LC, Singh BB, Dagenais S. Scientific basis for Ayurvedic therapies. *Altern Ther Health Med*. 2001; 7(3): 44–54.
  65. Kulkarni R, Patwardhan B. Evidence-based Ayurveda: current status and future perspectives. *J Ayurveda Integr. Med.*, 2012; 3(1): 3–8. doi:10.4103/0975-9476.93946.
  66. Halliwell B. Oxidative stress and neurodegeneration: where are we now? *J Neurochem*. 2006; 97(6): 1634–1658. doi:10.1111/j.1471-4159.2006.03907.x.
  67. Uttara B, Singh AV, Zamboni P, Mahajan RT. Oxidative stress and neurodegenerative diseases: a review of upstream and downstream antioxidant therapeutic options. *Curr Neuropharmacol*. 2009; 7(1): 65–74. doi:10.2174/157015909787602823.
  68. Butterfield DA, Halliwell B. Oxidative stress, dysfunctional glucose metabolism and Alzheimer's disease. *Free Radic. Biol. Med.*, 2019; 134: 624–638. doi:10.1016/j.freeradbiomed.2019.01.019.
  69. Gibson RS, Bailey KB, Gibbs M, Ferguson EL. A review of phytate, iron, zinc, and calcium concentrations in plant-based complementary foods used in low-income countries and consequences for bioavailability. *Food Nutr. Bull.*, 2010; 31(2): S134–S146. doi:10.1177/15648265100312S206.
  70. Hurrell RF, Egli I. Iron bioavailability and dietary reference values. *Am J Clin Nutr*. 2010; 91(5): 1461S–1467S. doi:10.3945/ajcn.2010.28674F
  71. Fairweather-Tait SJ, Teucher B. Iron and zinc bioavailability and dietary reference values. *Br. J. Nutr.*, 2002; 87(2): S151–S160. doi:10.1079/BJN2002534.
  72. Georgieff MK. Nutrition and the developing brain: nutrient priorities and measurement. *Am J Clin Nutr*. 2007; 85(2): 614S–620S. doi:10.1093/ajcn/85.2.614S.
  73. Prado EL, Dewey KG. Nutrition and brain development in early life. *Nutr Rev*. 2014; 72(4): 267–284. doi:10.1111/nure.12102.
  74. Black MM, Walker SP, Fernald LCH, Andersen CT, DiGirolamo AM, Lu C, et al. Early childhood development coming of age: science through the life course. *Lancet*. 2017; 389(10064): 77–90. doi:10.1016/S0140-6736(16)31389-7
  75. Shahidi F, Ambigaipalan P. Phenolics and polyphenolics in foods, beverages, and spices: antioxidant activity and health effects. *J. Funct., Foods*. 2015; 18: 820–897. doi:10.1016/j.jff.2015.06.018.
  76. Kris-Etherton PM, Hecker KD, Bonanome A, Coval SM, Binkoski AE, Hilpert KF, et al. Bioactive compounds in foods: their role in the prevention of cardiovascular disease and cancer. *Am J. Med.*, 2002; 113(9B): 71S–88S. doi:10.1016/S0002-9343(01)00995-0.
  77. United Nations Children's Fund (UNICEF). *The State of the World's Children 2019: Children, Food and Nutrition*. New York: UNICEF; 2019.
  78. Food and Agriculture Organisation of the United Nations (FAO). *Nutritional Variety Assessment: A Guide to Measurement*. Rome: FAO; 2011.
  79. Indian Council of Medical Research (ICMR). *Nutrient Requirements for Indians*. Hyderabad: National Institute of Nutrition; 2020.
  80. Conners CK. *Conners' Rating Scales–Revised: Technical Manual*. North Tonawanda: Multi-Health Systems; 2001.
  81. DuPaul GJ, Power TJ, Anastopoulos AD, Reid R. *ADHD Rating Scale–IV: Checklists, Norms, and Clinical Interpretation*. New York: Guilford Press; 1998.

82. American Psychiatric Association. *Diagnostic and Statistical Manual of Mental Disorders*. 5th ed., Text Revision (DSM-5-TR). Washington DC: APA; 2022.
83. Kolb B, Gibb R. Brain plasticity and behaviour in the developing brain. *J Can Acad Child Adolesc Psychiatry*. 2011; 20(4): 265–276.
84. Greenough WT, Black JE, Wallace CS. Experience and brain development. *Child Dev.*, 1987; 58(3): 539–559. doi:10.2307/1130197.
85. Kemper KJ, Vohra S, Walls R. The use of complementary and alternative medicine in paediatrics. *Pediatrics*. 2008; 122(6): 1374–1386. doi:10.1542/peds.2008-2173.
86. Vohra S, Surette S, Mittra D, Rosen LD, Gardiner P, et al. Pediatric integrative medicine: paediatrics and complementary therapies. *Pediatrics*. 2007; 119(2): e292–e299. doi:10.1542/peds.2006-1978.
87. Kim HJ, Lee JH, Kim YJ, Kim MJ. Antioxidant activities of seed oils and their neuroprotective effects. *Food Chem.*, 2012; 135(4): 2561–2567. doi:10.1016/j.foodchem.2012.06.104.
88. Ramadan MF. Functional properties, nutritional value, and industrial applications of pumpkin seed oil. *J Food Lipids.*, 2006; 13(3): 1–23. doi:10.1111/j.1745-4522.2006.00056.x.
89. Benton D. The influence of dietary status on the cognitive performance of children. *Mol. Nutr. Food Res.*, 2010; 54(4): 457–470. doi:10.1002/mnfr.200900158.
90. Gómez-Pinilla F. Brain foods: the effects of nutrients on brain function. *Nat Rev Neurosci.* 2008; 9(7): 568–578. doi:10.1038/nrn2421.
91. Prasad AS. Zinc in human health: effect of zinc on immune cells. *Mol. Med.*, 2008; 14(5–6): 353–357. doi:10.2119/2008-00033.Prasad.
92. Beard JL. Iron biology in immune function, muscle metabolism, and neuronal functioning. *J Nutr.*, 2001; 131(2S-2): 568S–580S. doi:10.1093/jn/131.2.568S.
93. Barbagallo M, Domínguez LJ. Magnesium and ageing. *Curr Pharm. Des.*, 2010; 16(7): 832–839. doi:10.2174/138161210790883742.
94. Calder PC. Omega-3 polyunsaturated fatty acids and inflammatory processes: nutrition or pharmacology? *Br. J Clin. Pharmacol.*, 2013; 75(3): 645–662. doi:10.1111/j.1365-2125.2012.04374.x
95. Miller AH, Raison CL. The role of inflammation in depression: from evolutionary imperative to modern treatment target. *Nat Rev Immunol.* 2016; 16(1): 22–34. doi:10.1038/nri.2015.5.
96. Mindell JA, Williamson AA. Benefits of the bedtime routine in young children: sleep, development, and beyond. *Sleep Med Rev.* 2018; 40: 93–108. doi:10.1016/j.smrv.2017.10.007.
97. Owens JA, Weiss MR. Insufficient sleep in adolescents: causes and consequences. *Minerva Pediatr.*, 2017; 69(4): 326–336. doi:10.23736/S0026-4946.17.05042-7.
98. Dantzer R, O'Connor JC, Freund GG, Johnson RW, Kelley KW. From inflammation to sickness and depression: when the immune system subjugates the brain. *Nat. Rev. Neurosci.*, 2008; 9(1): 46–56. doi:10.1038/nrn2297.
99. Zeisel SH. Nutritional importance of choline for brain development. *J Am Coll. Nutr.*, 2004; 23(6): 621S–626S. doi:10.1080/07315724.2004.10719433
100. Gómez-Pinilla F. Brain foods: the effects of nutrients on brain function. *Nat Rev Neurosci.* 2008; 9(7): 568–578. doi:10.1038/nrn2421.
101. Patwardhan B, Mutalik G, Tillu G. Coordinated approaches for health: biomedical research on Ayurveda. *J Ayurveda Integr. Med.*, 2015; 6(1): 1–4. doi:10.4103/0975-9476.146535.
102. Rastogi S. Building bridges between Ayurveda and modern science: a narrative review. *Curr Sci.* 2015; 108(9): 1625–1632.
103. Merikangas KR, He JP, Burstein M, Swanson SA, Avenevoli S, Cui L, et al. Lifetime prevalence of mental disorders in U.S. adolescents: results from the National Comorbidity Survey Replication–Adolescent Supplement (NCS-A). *J Am Acad Child Adolesc Psychiatry.* 2010; 49(10): 980–989. doi:10.1016/j.jaac.2010.05.017.
104. World Health Organization. *Guidelines on Safety Monitoring of Herbal Medicines in Pharmacovigilance Systems*. Geneva: WHO Press; 2004.
105. Food Safety and Standards Authority of India (FSSAI). *Nutraceutical Regulations and Guidelines*. New Delhi: FSSAI; 2016.
106. Illes J, Bird SJ. Neuroethics: a modern context for ethics in neuroscience. *Trends Cogn. Sci.*, 2006; 10(8): 352–357. doi:10.1016/j.tics.2006.06.002
107. Shahidi F, Ambigaipalan P. Phenolics and polyphenolics in foods: antioxidant activity and health effects. *J Agric Food Chem.*, 2015; 63(37): 1–15. doi:10.1021/acs.jafc.5b02907.
108. Georgieff MK. Long-term brain and behavioural consequences of early iron deficiency. *Nutr. Rev.*, 2011; 69(1): S43–S48. doi:10.1111/j.1753-4887.2011.00432.x
109. Prado EL, Dewey KG. Nutrition and brain development in early life. *Nutr Rev.* 2014; 72(4): 267–284. doi:10.1111/nure.12102
110. Black MM, Walker SP, Fernald LCH, Andersen CT, DiGirolamo AM, Lu C, et al. Early childhood development coming of age: science through the life course. *Lancet.* 2017; 389(10064): 77–90. doi:10.1016/S0140-6736(16)31389-7.
111. Chang JP, Su KP, Mondelli V, Pariante CM, Su KP. Omega-3 polyunsaturated fatty acids in youth with attention-deficit/hyperactivity disorder: a systematic review and meta-analysis. *Transl Psychiatry.* 2018; 8(1): 170. doi:10.1038/s41398-018-0212-3.
112. Gow RV, Hibbeln JR, Parletta N. Current evidence and future directions for research with omega-3 fatty acids and ADHD. *Curr Opin Clin Nutr Metab Care.*

- 2015; 18(2): 133–138.  
doi:10.1097/MCO.000000000000149.
113. Verlaet AAJ, Maasackers CM, Hermans N, Savelkoul HFJ. Nutrition, immunological mechanisms, and dietary immunomodulation in ADHD. *Eur Child Adolesc Psychiatry*. 2014; 23(7): 519–529. doi:10.1007/s00787-013-0514-8.
114. Lopresti AL. Oxidative and nitrosative stress in attention-deficit/hyperactivity disorder: possible links to neurobiology and treatment. *J Atten Disord*. 2015; 19(2): 97–112. doi:10.1177/1087054712455067.
115. Takeda A, Tamano H. Zinc signalling and its roles in the hippocampus. *Neuroscience*. 2012; 215: 1–12. doi:10.1016/j.neuroscience.2012.04.036.
116. Barbagallo M, Dominguez LJ. Magnesium and ageing. *Curr. Pharm. Des.*, 2010; 16(7): 832–839. doi:10.2174/138161210790883742.
117. Patwardhan B, Mashelkar RA. Traditional medicine-inspired approaches to drug discovery: can Ayurveda show the way forward? *Drug Discov Today*. 2009; 14(15–16): 804–811. doi:10.1016/j.drudis.2009.05.009.
118. Kulkarni R, Patki P, Jog V, Gandage S, Patwardhan B. Treatment of attention deficit hyperactivity disorder using Ayurvedic Medhya Rasayana drugs: a pilot clinical study. *J Ayurveda Integr. Med.*, 2011; 2(2): 71–77. doi:10.4103/0975-9476.82526
119. World Health Organization. *WHO Traditional Medicine Strategy 2014–2023*. Geneva: WHO Press; 2014.
120. Food Safety and Standards Authority of India (FSSAI). *Food Safety and Standards (Health Supplements, Nutraceuticals, Food for Special Dietary Use, Food for Special Medical Purpose, Functional Food and Novel Food) Regulations*. New Delhi: FSSAI; 2016.
121. Zhang, C., Chen, L., Wang, X., Pan, P., Tang, Y. & Hui, P. Cucurbita pepo: Traditional Uses and Pharmacological Properties. *Pharmacognosy Magazine*, 2021; 17(75): 511-517. [https://doi.org/10.4103/pm.pm\\_170\\_20](https://doi.org/10.4103/pm.pm_170_20).
122. Gole, S., Gwalani, S. & Sarap, R. Behavioural Problems in Children and the Role of Medhya Rasayana: An Ayurvedic Review. *Journal of Ayurveda and Integrated Medical Sciences*, 2026; 10(12): 284-288. <https://doi.org/10.21760/jaims.10.12.42>.
123. Pumpkin Seeds Oil Rescues Colchicine-Induced Neurotoxicity in Rats via Modifying Oxidative Stress, DNA Damage, and Immunoexpression of BDNF and GFAP. *ScienceDirect* 2025; 1: 1-10. <https://doi.org/10.1016/j.toxlet.2025.01.001>.
124. Pumpkin Seeds as a Natural Remedy for Anaemia: Nutritional Analysis and Therapeutic Potential. *International Journal of Food Science* 2023. <https://doi.org/10.1155/2023/40535566>.