



SUDDEN CARDIAC DEATH IN 15 TO 39-YEAR-OLD TAIWANESE POPULATION

Shu-Ching Chiu, MSN^{a,d}, Mei-Ling Lin, Ph.D^a, Shu-Chen Yang, MSN^a, Lee-Wen Pai, MSN^a, Chuan-Mei Chen, MSN^a, Horng-Mo Lee, Ph. D^{b,c}, Yueh-Chin Chung, Ph. D^{a,*}

^aDepartment of Nursing, Central Taiwan University of Science and Technology, Taichung, Taiwan.

^bInstitute of Pharmaceutical Science and Technology, Central Taiwan University of Science and Technology, Taichung, Taiwan.

^cSchool of Medical Laboratory Sciences and Biotechnology, Taipei Medical University, Taipei, Taiwan.

^dPh. D student, School of Nursing, National Taipei University of Nursing & Health Science, Taiwan.

Article Received on 18/05/2015

Article Revised on 10/06/2015

Article Accepted on 01/07/2015

***Correspondence for
Author**

Yueh- Chin Chung

Associate Professor,
Department of Nursing,
Central Taiwan University
of Science and Technology,
Taichung, Taiwan.

ABSTRACT

Background: Sudden cardiac death (SCD) among the young (15 to 39-year-old) population has gained increasing attention in recent years, above all when occurring in young people, remains a major clinical problem. **Objective:** The aim of this study is to examine the effects of age, sex, and season and causes on outpatient SCDs from 2009 to 2010 in Taiwan. **Methods:** Cross-sectional study. Dates were identified using the International Classification of Disease (ICD) codes of inpatient

SCDs recorded by National Health Insurance Research Database (HNIRDs) from 2009 to 2010. **Results:** A total 684 SCD patients were men (70.75%), with a mean age of 30.9 years. The annual incidence of SCD increased by 17.0% from 2009 (3.53 per 100,000 person-years) to 2010 (4.13 per 100,000 person-years). The incidence was higher for men (4.85 to 5.85 per 100,000 person-years) than women (2.18 to 2.37 per 100,000 person-years). After adjusting for sex and age, increased risk of SCD in winter was observed (RR=1.30, 95% CI 0.23-1.43; p=0.01). **Conclusions:** The study posits that the major determinants of SCDs in Taiwan were male, 30 to 39 years old, and winter. Clinical diagnoses of SCDs include cardiovascular diseases of cardiac arrest (n=555, 88.66%), mitral valve prolapse, aortic valve stenosis, acute

myocarditis, and coronary artery aneurysm disease. In relation to these, ECG to a history and physical examination prevention and cardiopulmonary resuscitation (CPR) programs may help reduce the risk of SCD.

KEY WORDS: sudden Cardiac Death (SCD), incidence, young, cardiac arrest, sex distribution.

INTRODUCTION

Sudden cardiac death (SCD) is a major public health problem.^[1,2] SCD is a leading cause of death in the US, but the relative public health burden remains unknown.^[3]

SCD is one of the major problems in the western world with approximately 70,000 to 100,000 SCD patients in Germany and 450,000 SCD victims in the US. SCD is not caused by a single factor, but is a multi-factorial problem. In 50% of SCD victims, SCD is the first manifestation of heart disease.^[4] Over the past few decades, improvements in primary and secondary prevention measures have led to significant declines in cardiac origin.^[5-7] In recent prospective studies that used multiple sources from the US, SCD rates have been found to range from 0.50 to 13 per 100,000 people in the general population.^[8] In the US, SCD is estimated to occur once every minute; it is the most common and often the first manifestation of coronary heart disease. in the United States.^[9] Applying the incidence rates to the European and Asian populations would result in incidence rates of 0.93 to 21.7 per 100,000 people,^[10-13] and 2.3 to 10.1 per 100,000 people, respectively.^[14-21] In general, the sudden death (SD) risk is higher in males than in females (2-4.3:1.14),^[19,22-24] and increases with age.^[25] In young adults, the mortality rate for SCD per 100,000 person-years during the study period was 6.7 for males and 1.4 for females, and the incidence of SCD was higher in men than in women.^[26]

Aside from being the most common and the typical first manifestation of coronary heart disease, SCD is responsible for 50% to 88% of the mortality from cardiovascular diseases in the US and in other developed countries.^[9] Several population-based studies have documented a 15% to 19% decline in the incidence of SCD caused by coronary heart disease since the early 1980s. However, the increasing incidence of coronary heart disease may halt this decline in the future.^[27] Continuous surveillance is necessary in clarifying future trends in SCD because of an increasing incidence of diabetes mellitus (2.2 to 13.7 per 100,000 people)

and hypertension (23.0%). The SCD risk factors have also been positively associated with increased SCD risk.^[26, 28]

To the research, no study has yet to identify the risk of SCD in the 15 to 39-year-old population using the outpatient SCD data recorded by HNIRDs in Taiwan. Although SCD is rare, it is usually fatal and has seen an increase in recent years. Therefore, a retrospective study was conducted to understand the incidence and contributing factors of SCD in Taiwan. We identify the young population, we examined the 15 to 39-year-old population as well as explored the influence of sex, age, seasonal patterns, and associated diseases on the incidence of SCD. Results of the study will be helpful in the development of preventive strategies.

MATERIALS AND METHODS

Definitions

A total 684 people were diagnosed with SCD during 2009-2010. SCD in the young is defined as an unexpected natural death within 24 h after the onset of symptoms in persons aged 15 to 39. All cases wherein death occurred within 24 h from the onset of the underlying cause were considered SCD as long as the cause of death was either of cardiac origin or unknown. SD included sudden cardiac death (SCD), sudden non-cardiac death (SNCD) and unknown SD, which were identified from the NHIRDs from 2009 to 2010 using the ninth revision of the International Classification of Diseases, Clinical Modification (ICD-9-CM) for research population. SCDs included atherosclerotic disease (familial hypercholesterolemia, myocardial infarction, and other acute ischemic heart diseases) (ICD-9-CM 272, 410, 411), cardiomyopathy (dilated cardiomyopathy, obstructive hypertrophic cardiomyopathy, other hypertrophic cardiomyopathy, other cardiomyopathies, and total cardiomyopathy) (ICD-9-CM 425), conduction disorders (Cardiac arrhythmia), unspecified.

Ventricular tachycardia. Ventricular fibrillation and flutter. Total number of conduction disorders. Preexcitation syndrome, Other specified conduction disorders) (ICD-9-CM 427, 426), and other cardiac causes (cardiac arrest, mitral valve prolapse, aortic valve stenosis, acute myocarditis, myocarditis in diseases classified elsewhere, and coronary artery aneurysm) (ICD-9-CM 424, 422, 427.5, 414). SD with unknown cause (unknown SD) (ICD-9-CM 798.1, 798.2, 798.9) and sudden non-cardiac death (SNCD) included epilepsy, pulmonary embolism, non-traumatic cerebral hemorrhage, and asthma (ICD-9-CM 345, 415, 430, 431, 432, 493). Comorbidities included hypertension, hyperlipidemia, diabetes mellitus, acute gastritis, and colitis (ICD-9-CM 401, 250, 535, 558).^[29] We have excluded congenital cardiac

diseases. This study protocol was approved by the institutional review board (IRB) of China Medical University Hospital (CMUH103-REC1-088).

Statistical analysis

The incidence of SCD, SD, unknown SD and SD of comorbidities were calculated by dividing the number of new cases during 2009-2010 year by the number of people at risk in the population at the beginning of the study. Descriptive data were presented as percentages. Poisson regression models were used to assess the relative risks (RRs) and 95% confidence intervals (CIs) of SCD incidence associated with sex, age, and seasonal variables. The Mantel-Haenszel (M-H) method was used to calculate odds ratios adjusted for confounders (sex and age). *p*-values <0.05 were considered statistically significant.

RESULTS

Demographic characteristics of SCD, unknown SD, SNCD, SCD and SCD of comorbidities

During the two-year period (2009 to 2010), 684 people from the general population were diagnosed with SCD (incidence rate: 3.83 per 100,000). Men comprised 70.75% (n: 481) of the cases, while women comprised 29.25% (n: 203). 2.37 times more men than women died from SCD, and the average age was 30.9 years. The 30 to 39-year-old group accounted for 60.00% (n: 406) of the cases, with the 20 to 29-year-old and 15 to 19-year-old groups accounting for 28.96% (n: 199) and 11.04% (n: 79), respectively. The major cause of SCD was other cardiac causes (cardiac arrest, mitral and aortic valve prolapse, acute myocarditis, and coronary artery aneurysm) 626 (incidence rate: 3.43 per 100,000), followed by conduction disorders 39 (incidence rate: 0.22 per 100,000), and atherosclerotic diseases 23 (incidence rate: 0.12 per 100,000) (Table 1). SD with unknown cause (unknown SD) had an incidence rate of 12.49 per 100,000 (n: 2229). SNCD had an incidence rate of 2.92 per 100,000 (n: 522).

From the general population, 3,435 were diagnosed with SD (incidence rate: 18.99 per 100,000). Men and women were involved in 71.69% (n: 2462) and 28.33% (n: 973) of the cases. The ratio of male to female incidence was 2.58:1. The average age was 31.2 years. The 30 to 39-year-old group accounted for 56.42% (n: 1938) of the cases, while the 20 to 29-year-old and 15 to 19-year-old groups accounted for 32.78% (n: 1,126) vs 10.80% (n: 371), respectively.

The SCD of comorbidities included hypertension, hyperlipidemia, diabetes mellitus, acute gastritis, and colitis. Men accounted for 70.75% (n: 208) of the cases, while women accounted for 29.25% (n: 86) (incidence rate: 1.65 per 100,000) (Table 1).

The incidences Association of SCD with age group and sex

The annual incidence of SCD increased by 17.0% from 2009 (3.53 per 100,000 person-years) to 2010 (4.13 per 100,000 person-years). The incidence was higher for men (4.85 to 5.85 per 100,000 person-years) than for women (2.18 to 2.37 per 100,000 person-years). In the incidence, the 30 to 39-year-old group (4.96 to 5.71 per 100,000 person-years) had the highest incidence, while the 15 to 19-year-old group (1.98 to 2.74 per 100,000 person-years) had the lowest incidence (Table 2).

The incidence of male in 2010 was higher than that of male in 2009 (OR = 1.34, 95% CI 1.12-1.60, $p < 0.001$). The incidence was higher in 2009 15 to 19-year-old group in 2010 than in the 15 to 19-year-old group in 2009 (OR = 1.53, 95% CI 1.24-1.88, $p < 0.000$). The incidence was significantly higher in the 30 to 39-year-old group in 2010 than in the 30 to 39-year-old group in 2009 (OR = 1.35, 95% CI 1.13-1.61, $p < 0.001$). The incidence was significantly higher in the all-year-old group in 2010 than in the all-year-old group in 2009 (OR = 1.29, 95% CI 1.08- 1.54, $p < 0.006$) (Table 2).

Association of SCD with seasonal variation

The incidence of SCD was 1.10, 1.02, 0.89 and 0.82 per 100,000 person-years in winter (December to February), summer (June to August), spring (March to May), and autumn (September to November). The percentage of SCD was highest in the winter and lowest in the summer. Correspondingly, the age-adjusted RR were 1.30 (95% CI -0.23-1.43, $p < 0.01$) in winter, 1.13 (95% CI -0.04-0.15) in summer, and -0.40 (95% CI -0.11- 0.08) in spring, (Table 3).

Table 1. Demographic characteristics of sudden cardiac death, sudden death, and comorbidities, 2009–2010.

Causes of death	ICD-9-CM	15-19 years N (%)	20-29 years N (%)	30-39 years N (%)	Male N (%)	Female N (%)	Total N (%)	incidence*
Sudden cardiac death(SCD)		79(11.04)	199(28.96)	406(60.00)	481(70.75)	203(29.25)	684(100.00)	3.83
Atherosclerotic disease								
(Familiar hypercholesterolaemi. Myocardial infarction. Other acute ischaemic heart diseases.)	272,410, 411	0(0.00)	5(21.74)	18(78.26)	21(91.30)	2(8.70)	23(100.00)	0.12
Cardiomyopathy								
Dilated cardiomyopathy, Obstructive hypertrophic cardiomyopathy, Other hypertrophic ,cardiomyopathy,Other cardiomyopathies ,Total cardiomyopathy)	425	0(0.00)	1(20.00)	2(40.00)	2(40.00)	0(0.00)	5(100.00)	0.01
Conduction disorders								
(Cardiac arrhythmia, unspecified. Ventricular tachycardia. Ventricular fibrillation and flutter. Total number of conduction disorders. Preexcitation syndrome , Other specified conduction disorders)	427,426	3(7.69)	11(28.21)	25(64.10)	20(51.28)	19(48.72)	39(100.00)	0.22
Other cardiac causes								
(Cardiac arrest Mitral (valve) prolapse Aortic (valve) stenosis Acute myocarditis Myocarditis in diseases classified	427.5,424, 422,414	77(11.76)	181(29.58)	363(58.66)	443(71.24)	183(28.76)	626(100.00)	3.43

elsewhere Coronary artery aneurysm)								
Sudden death with unknown cause (unknown SD)	798	243(10.90)	768(34.46)	1218(54.64)	1608(72.14)	621(27.86)	2229(100.00)	12.49
Sudden noncardiac death (SNCD)								
(Epilepsy.Pulmonary embolism.Nontraumatic cerebral hemorrhage. Asthma)	345,415, 430,431, 432,493	52(9.96)	156(29.89)	314(60.15)	370(70.88)	152(29.12)	522(100.00)	2.92
Sudden death (SD)†	798.1,798. 2,798.9	371(10.80)	1126(32.78)	1938(56.42)	2462(71.69)	973(28.33)	3435(100.00)	18.99
Comorbidities‡								
(hypertension hyperlipidemia, Diabetes mellitus, Acute gastritis and colitis.)	401,250, 535,558	10(3.40)	76(25.85)	208(70.75)	208(70.75)	86(29.25)	294(100.00)	1.65

*Incidence rates are per 100,000 person-years during 2009-2010.

†Sudden death (SD) include Sudden cardiac death (SCD), unknown SD, and Sudden noncardiac death (SNCD).

‡Comorbidities include hypertension, **hyperlipidemia, diabetes mellitus, acute gastritis and colitis.**

Table 2. Incidence rate* of sudden cardiac death, sudden death 2009–2010, by year, age and sex group.

Variable	2010 n(Incidence rate)	2009 n(Incidence rate)	Odds ratios (ORs)†	95% confidence intervals (CIs)	<i>p</i> value
Sex					
Male	264(5.58)	220(4.85)	1.34	1.12-1.60	0.001
Female	104(2.37)	96(2.18)	1.11	0.90-1.37	0.311
Age					
15-19	44(2.74)	32(1.98)	1.53	1.24-1.88	0.000
20-29	106(3.05)	96(2.71)	1.18	0.97-1.43	0.093
30-39	218(5.71)	188(4.96)	1.35	1.13-1.61	0.001
Total	368(4.13)	316(3.53)	1.29	1.08-1.54	0.006

* Incidence rate (per 100,000 person-years) is the number of new cases of SCD divided by the size of the population at risk in each year.

* Data are represented as the number of the incidence and the rate in the parenthesis.

†The Mantel-Haenszel (M-H) method was used to calculate odds ratios adjusted for confounders (sex and age).

Table 3. Incidence rate* of sudden cardiac death, 2009-2010, by months and season

Months	n	Season	Total/Months	Incidence Rate	RR (95%CI)†	<i>p</i> -value
3	54	Spring	0.30	0.89	-0.40(-0.11~-0.08)	0.69
4	55		0.31			
5	50		0.28			
6	53	Summer	0.30	1.02	1.13(-0.04~-0.15)	0.26
7	60		0.34			
8	69		0.39			
9	46	Autumn	0.26	0.82	1(reference)	—
10	53		0.30			
11	47		0.26			
12	59	Winter	0.33	1.10	1.30(-0.23~1.43)	0.01
1	70		0.43			
2	62		0.35			

*Incidence rates are per 100,000 person-years during 2009-2010.

†The relative risks (95% Confidence Intervals, CI) are derived from Poission regression model adjusted for sex and age.

DISCUSSION

In this study, the annual year incidence of SCD was 3.53 to 4.13 per 100,000 person-years in Taiwan. The present analyses demonstrate marked variations in the occurrence of SCD, with peaks in the 30 to 39-year-old groups, in males, and during the winter months. Clinical

diagnoses of SCDs include cardiovascular diseases of cardiac arrest, mitral valve prolapse, aortic valve stenosis, acute myocarditis, and coronary artery aneurysm disease. Our findings provide strong evidence of an ongoing epidemiological transition in Taiwan, where treatable cardiac causes the major of cardiac arrest were responsible for the SCD. Such evidence would be helpful in the development of preventive strategies.

In this study, 481 (70.75%) of the cases involved males and 203 (29.25%) involved females of SCD. The male-to-female ratio was 2.37:1.^[14,19,30,31] Thus, given that the latter predominantly affects males, SCD is perceived as a male problem.^[21,22] Similar to overall SCD, SCD in this study is more common in males than in females (mean age range: 24 to 32 years). Studies on the incidence and causes of SCD in the young (1–40 years) are often restricted to male athletes.^[32-34] The age group with the highest incidence was the 30 to 39-year-old group, while the 15 to 19-year-old group had the lowest incidence. Previous studies have shown that SCD incidence increases with age.^[34,35]

Our study confirmed that the incidence was at its lowest and highest in September and in January, respectively. The literature describes the relationship between cold weather and SCD^[36,37] which likely contributes to the number of deaths occurring in winter. A seasonal trend in cardiovascular diseases was observed in the current study, with the highest incidence occurring during the colder winter months. The same finding has been observed in many countries.^[38-40] The implications of this finding are important for testing the relative importance of the proposed mechanisms. Furthermore, understanding the influence of seasonal patterns and other factors is essential in implementing effective public health measures. These could result in increased blood pressure through increased heart rate and peripheral vascular resistance. The seasonal variability of CVDs was clearly demonstrated by the epidemiological data, which showed mostly a peak in winter months. Many theories explain the possible reasons underlying the development of a disease in winter more than in other times of the year. Many biological aspects are also known to contribute to the development of disorders. This finding may have important implications, and the risk of cardiovascular diseases appears to be the greatest during the winter months. The knowledge of the role of environmental and biological factors could be used to improve prevention measures and educational strategies, especially in people with a risk of diseases. People should be informed of the increased risk of cardiovascular diseases during the cold seasons; they should also be educated about the importance of having regular physical activities and

dressing warmly in the winter.^[37-40] Low temperature has a hazardous effect on blood pressure and may alter the ratio of myocardial oxygen supply to demand; increase ventricular wall stress, cardiac work and oxygen requirements; as well as reduce mechanical efficiency and coronary blood flow.^[37,39,40] For the effective prevention of SCD across Taiwan, such variations must be considered when developing programs to reduce risk. Multiple studies on screening have shown that adding an ECG to a history and physical examination is more sensitive than history and physical examination alone in identifying those who are potentially at risk for SCD.^[41]

In this study, the annual year incidence of SCD was 3.53 to 4.13 per 100,000 person-years, which is most likely because of the increasing number of cardiac arrest (n=555,88.66%) in the age groups.^[42] The activation of the sympathetic nervous system and secretion of catecholamine are increased in response to cold temperatures. In this study, we observed that clinical diagnoses of SCD major causes of cardiac arrest. In cardiac rehabilitation programs, cardiac arrests occur at a rate of 1 in 12 000 to 15 000, and during stress testing, cardiac arrest occurs at a rate of 1 per 2000, at least 6 times greater than the general incidence of sudden cardiac death for patients known to have heart disease.^[42] When the blood flow to the heart is stopped due to narrow or obstructed coronary arteries, a heart attack occurs. This may lead to a cardiogenic shock and cardiac arrest. The survival rate after bystander cardiopulmonary resuscitation (CPR) was 2.6-fold higher than the rate for those where no treatment was given until the ambulance arrived.^[43]

The SCD of comorbidities included hypertension, hyperlipidemia, diabetes mellitus, acute gastritis, and colitis. (incidence rate: 1.65 per 100,000). The number of SCD risk factors (hypertension and diabetes) was positively associated with increased SCD risk.^[44,45] Patients who have been identified to have the highest risk require the most immediate anti-chronic disease therapeutic intervention.

There are some limitations in using the Health Insurance Database in this study. First, claims data were used, which indicated that only those who visited a medical institution were enrolled. NHI database-registered cases of SCD and SD are not linked to death certificate data; therefore, we could not confirm the incidence of SCD and SD using only the relevant data from medical records. We could only confirm the cases of SCD and SD from the NHI database. Second, this database does not document job (athletes), family status and lifestyle.

The lack of such valuable information makes it difficult to evaluate the respective influences of these factors.

ACKNOWLEDGMENTS

This study was supported by a grant from the Central Taiwan University of Science and Technology. It was conducted partly using data from the National Health Insurance Research Database provided by the Bureau of National Health Insurance, Department of Health and managed by the National Health Research Institute. The interpretation and conclusions contained herein do not represent the opinions of the Bureau of National Health Insurance, the Department of Health or the National Health Research Institute.

REFERENCES

1. Go AS, Mozaffarian D, Roger VL, Benjamin EJ, Berry JD, Blaha MJ, et al. Executive summary: heart disease and stroke statistics-2014 update: a report from the American Heart Association. *Circulation*, 2014; 129: 399-410.
2. Deo R, Albert CM. Epidemiology and genetics of sudden cardiac death. *Circulation*, 2012; 125: 620–37.
3. Stecker EC, Reinier K, Marijon E, Narayanan K, Teodorescu C, Uy-Evanado A, et al. Public health burden of sudden cardiac death in the United States. *Circ Arrhythm Electrophysiol*, 2014; 7: 212-7.
4. Trappe HJ. Sudden cardiac death and automated external defibrillators. Where we are in 2012. *Herz*, 2012; 37: 416-23.
5. Ford ES, Ajani UA, Croft JB, Critchley JA, Labarthe DR, Kottke TE, et al. Explaining the decrease in U.S. deaths from coronary disease, 1980–2000. *N Engl J Med*, 2007; 356: 2388–98.
6. Rosamond WD, Chambless LE, Folsom AR, Cooper LS, Conwill DE, Clegg L, et al. Trends in the incidence of myocardial infarction and in mortality due to coronary heart disease, 1987 to 1994. *N Engl J Med*, 1998; 339: 861–7.
7. Eckart RE, Shry EA, Burke AP, McNear JA, Appel DA, Castillo-Rojas LM, et al. Sudden death in young adults: an autopsy-based series of a population undergoing active surveillance. *J Am Coll Cardiol*, 2011; 58: 1254-61.
8. Silva AC, Santos L, Dinis-Oliveira RJ, Magalhães T, Santos A. Sudden Cardiac Death in Young Adult. *Cardiovasc Toxicol*, 2014; 9. [Epub ahead of print].

9. Harmon KG, Asif IM, Klossner D, Drezner JA. Incidence of sudden cardiac death in national collegiate athletic association athletes. *Circulation*, 2011; 123: 1594–600.
10. Michael Papadakis, Sanjay Sharma, Steve Cox, Mary N Sheppard, Vasileios F Panoulas, Elijah R Behr. The magnitude of sudden cardiac death in the young: a death certificate-based review in England and Wales. *Europace*, 2009; 11: 1353–8.
11. Risgaard B, Winkel BG, Jabbari R, Behr ER, Ingemann-Hansen O, Thomsen JL, et al. Burden of sudden cardiac death in persons aged 1 to 49 years: nationwide study in Denmark. *Circ Arrhythm Electrophysiol*. 2014; 7: 205-11.
12. Margey R, Roy A, Tobin S, O'Keane CJ, McGorrian C, Morris V, et al. Sudden cardiac death in 14- to 35-year olds in Ireland from 2005 to 2007: a retrospective registry. *Europace*, 2011; 13: 1411-18.
13. Risgaard B, Nielsen JB, Jabbari R, Haunsø S, Holst AG, Winkel BG, et al. Prior myocardial infarction in the young: predisposes to a high relative risk but low absolute risk of a sudden cardiac death. *Europace*, 2013; 15: 48-54.
14. Wang H, Yao Q, Zhu S, Zhang G, Wang Z, Li Z, et al. The autopsy study of 553 cases of sudden cardiac death in Chinese adults. *Heart*, 2013 ; 9. [Epub ahead of print].
15. Siu CW, Pong V, Ho HH, Liu S, Lau CP, Li SW, et al. Are MADIT II criteria for implantable cardioverter defibrillator implantation appropriate for Chinese patients? *J Cardiovasc Electrophysiol*, 2010; 21: 231–5.
16. Hua W, Zhang LF, Wu YF, Liu XQ, Guo DS, Zhou HL, et al. Incidence of sudden cardiac death in China: analysis of 4 regional populations. *J Am Coll Cardiol*, 2009; 54: 1110–18.
17. Tanno K, Miyoshi F, Watanabe N, Minoura Y, Kawamura M, Ryu S, et al. Are the MADIT II criteria for ICD implantation appropriate for Japanese patients? *Circ J*, 2005; 69: 19–22.
18. Maruyama M, Ohira T, Imano H, Kitamura A, Kiyama M, Okada T, et al. Trends in sudden cardiac death and its risk factors in Japan from 1981 to 2005: the Circulatory Risk in Communities Study (CIRCS). *BMJ Open*, 2012; 22: 2:e000573.
19. Cho JG, Park HW, Rhew JY, Lee SR, Chung WK, Park OY, et al. Clinical characteristics of unexplained sudden cardiac death in Korea. *Jpn Circ J*, 2001; 65: 18-22.
20. Cho JG, Park HW, Rhew JY, Lee SR, Chung WK, Park OY, et al. Arrhythmogenic right ventricular cardiomyopathy and sudden cardiac death in young Koreans. *Circ J*, 2003; 67: 925-8.

21. Neuspiel DR, Kuller LH. Sudden and unexpected natural death in childhood and Adolescence. *JAMA*, 1985; 254: 1321–5.
22. Behr E, Wood DA, Wright M, Syrris P, Sheppard MN, Casey A, et al. Sudden Arrhythmic Death Syndrome Steering Group. Cardiological assessment of first-degree relatives in sudden arrhythmic death syndrome. *Lancet*, 2003; 362: 1457-9.
23. Wisten A, Forsberg H, Krantz P, Messner T. Sudden cardiac death in 15-35-years old in Sweden during 1992-99. *J Intern Med*, 2002; 252: 529-36.
24. Melissa HK, Gregg CF, Eric DP, Anne BC, Adrian F H, Gillian DS, et al. Systematic Review of the Incidence of Sudden Cardiac Death in the United States. *J Am Coll Cardiol*, 2011; 57: 794-801.
25. Silva AC, Santos L, Dinis-Oliveira RJ, Magalhães T, Santos A. Sudden Cardiac Death in Young Adult. *Cardiovasc Toxicol*, 2014; 9. [Epub ahead of print].
26. Reddy KS, Yusuf S. Emerging epidemic of cardiovascular disease in developing countries. *Circulation*, 1998; 97: 596–601.
27. de Vreede-Swagemakers JJ, Gorgels AP, Dubois-Arbouw WI, van Ree JW, Daemen MJ, Houben LG, et al. Out-of-hospital cardiac arrest in the 1990s: a population-based study in the Maastricht area on incidence, characteristics and survival. *J Am Coll Cardiol*, 1997; 30: 1500.
28. Vetter VL, Dugan NP. A discussion of electrocardiographic screening and sudden cardiac death prevention: evidence and consensus. *Curr Opin Cardiol*, 2013; 28: 139-51.
29. Vaartjes I, Hendrix A, Hertogh EM, Grobbee DE, Doevendans PA, Mosterd A, et al. Sudden death in persons younger than 40 years of age: incidence and causes. *Eur J Cardiovasc Prev Rehabil*, 2009; 16: 592-6.
30. Wisten A, Forsberg H, Krantz P, Messner T. Sudden cardiac death in 15–35-year olds in Sweden during 1992–1999. *J Intern Med*, 2002; 252: 529–36.
31. Chugh SS, Reinier K, Teodorescu C, Evanado A, Kehr E, Al SM, et al. Epidemiology of sudden cardiac death: clinical and research implications. *Prog Cardiovasc Dis*, 2008; 51: 213–28.
32. Neuspiel DR, Kuller LH. Sudden and unexpected natural death in childhood and Adolescence. *JAMA*, 1985; 254: 1321–5.
33. Behr E, Wood DA, Wright M, Syrris P, Sheppard MN, Casey A, et al. Sudden Arrhythmic Death Syndrome Steering Group. Cardiological assessment of first-degree relatives in sudden arrhythmic death syndrome. *Lancet*, 2003; 362: 1457-9.

34. Corrado D, Pelliccia A, Bjornstad HH, Vanhees L, Biffi A, Borjesson M, et al. Cardiovascular pre-participation screening of young competitive athletes for prevention of sudden death: proposal for a common European protocol. Consensus Statement of the Study Group of Sport Cardiology of the Working Group of Cardiac Rehabilitation and Exercise Physiology and the Working Group of Myocardial and Pericardial Diseases of the European Society of Cardiology. *Eur Heart J*, 2005; 26: 516–24.
35. Corrado D, Basso C, Rizzoli G, Schiavon M, Thiene G. Does sports activity enhance the risk of sudden death in adolescents and young adults? *J Am Coll Cardiol*, 2003; 42: 1959–63.
36. Akinwusi PO, Komolafe AO, Olayemi OO, Adeomi AA. Communicable disease-related sudden death in the 21st century in Nigeria. *Infect Drug Resist*, 2013; 6: 125-32.
37. Fares A. Winter cardiovascular diseases phenomenon. *N Am J Med Sci*, 2013; 5: 266-79.
38. Ben Ahmed H, Allouche M, Zoghlami B, Shimi M, Razghallah R, Gloulou F, et al. Diurnal, weekly and seasonal variation of sudden cardiac death in northern Tunisia. *Presse Med*, 2014; 43: e39-45.
39. Silverman RA, Ito K, Freese J, Kaufman BJ, De Claro D, Braun J, Prezant DJ. Association of ambient fine particles with out of hospital cardiac arrests in New York City. *Am J Epidemiol*, 2010; 172: 917- 23.
40. Sun RC, Yang SD, Zhou ZY, Liang JB. Clinicopathological analysis of sudden cardiac death cases by autopsy. *Zhonghua Yi Xue Za Zhi*, 2012; 92: 2707-9.
41. Yang J, Teehan D, Farioli A, Baur DM, Smith D, Kales SN. Sudden cardiac death among firefighters \leq 45 years of age in the United States. *Am J Cardiol*, 2013; 112: 1962-7.
42. Douglas P, Zipes, MD; Hein J.J. Wellens, MD. Sudden Cardiac Death. *Circulation*, 1998; 98: 2334-51.
43. Holmberg M, Holmberg S, Herlitz J. The problem of out-of-hospital cardiac-arrest prevalence of sudden death in Europe today. *Am J Cardiol*, 1999; 11; 83(5B): 88D-90D.
44. Kataoka M, Ito C, Sasaki H, Yamane K, Kohno N. Low heart rate variability is a risk factor for sudden cardiac death in type 2 diabetes. *Diabetes Res Clin Pract*, 2004; 64: 51-8.
45. Ohira T, Maruyama M, Imano H, Kitamura A, Kiyama M, Okada T, et al. Risk factors for sudden cardiac death among Japanese: the Circulatory Risk in Communities Study. *J Hypertens*, 2012; 30: 1137-43.