

**COMPARATIVE RELIABILITY OF THE TAU ANGLE VERSUS YEN, BETA, W, AND ANB ANGLES IN SAGITTAL SKELETAL ASSESSMENT USING FACAD SOFTWARE**

Dr. Suhas\*

India.

**\*Corresponding Author: Dr. Suhas**

India.

DOI: <https://doi.org/10.5281/zenodo.20022403>

**How to cite this Article:** Dr. Suhas\*. (2026). Comparative Reliability of The Tau Angle Versus Yen, Beta, W, and Anb Angles In Sagittal Skeletal Assessment Using Facad Software. European Journal of Pharmaceutical and Medical Research, 13(5), 356–362.

This work is licensed under Creative Commons Attribution 4.0 International license.



Article Received on 05/04/2026

Article Revised on 25/04/2026

Article Published on 04/05/2026

**INTRODUCTION**

Accurate assessment of sagittal skeletal relationships plays a pivotal role in orthodontic diagnosis and treatment planning. The ANB angle, introduced by Cecil C. Steiner, remains one of the most widely used parameters; however, it is influenced by cranial base length and jaw rotations. Similarly, the WITS appraisal proposed by Alexander Jacobson depends on the occlusal plane, limiting its reliability.

To overcome these limitations, alternative angular parameters such as the Beta angle, Yen angle, and W angle have been introduced. While these parameters attempt to reduce dependency on cranial landmarks, variability in landmark identification and geometric construction persists.

The TAU angle has recently been proposed as a novel parameter aimed at improving diagnostic reliability. However, its comparative performance with established sagittal indicators remains inadequately explored. Therefore, the present study aimed to evaluate the reliability, agreement, and diagnostic validity of the TAU angle using FACAD software.

**MATERIALS AND METHODS****Study Design**

A cross-sectional analytical study.

**Sample**

- Sample size: **40 subjects**
- Age range: **15–41 years**
- Inclusion criteria:
  - Good quality lateral cephalograms
  - No prior orthodontic treatment
- **Exclusion criteria**
  - Craniofacial anomalies
  - Distorted radiographs

**Data Collection**


Cephalometric analysis was performed using **FACAD software**. The following parameters were recorded:

- ANB angle
- WITS appraisal
- Beta angle
- Yen angle
- W angle
- TAU angle

**ANB ANGLE**

- Introduced by: Richard A. Riedel (1952), later Cecil C Steiner in 1953 used this analysis. It is the difference between the SNA angle and the SNB angle. It evaluates the sagittal relationship between the maxilla and mandible. This has been the most followed method of measuring sagittal discrepancy, but it has a major disadvantage in that differences were found between the interpretation of this angle and the actual discrepancy between the maxillary and mandibular bases. It is also affected by the slightest of changes in the point Nasion, length of the cranial base, and anterior face height. ANB increases clockwise and decreases with counterclockwise rotation of the jaws.<sup>[2]</sup>
- In Binder's study every 5 mm of anterior displacement of Nasion horizontally, the ANB angle reduces by 2.5°. A 5 mm upward displacement of Nasion decreases the ANB angle by 0.5°, and a 5


mm downward displacement increases the ANB angle by 1°



- Perpendiculars from Point A and B on the maxilla and mandible, respectively, are drawn onto the occlusal plane. The points of contact are labeled AO and BO, respectively. The distance between BO and AO is measured. According to Jacobson, in a skeletal Class I relationship, in females, AO and BO should coincide; in males, BO is ahead of AO by 1 mm in males. It changes with orthodontic treatment but does not change significantly with age (after permanent teeth have erupted completely) if untreated.<sup>[3]</sup>

- **Wit's Appraisal of Jaw Disharmony**
- Introduced by: Alex Jacobson (1975). Overcomes the shortcomings of the ANB angle.

| The mean value for Wits   |   |
|---------------------------|---|
| Skeletal Class I pattern: | BO coincides with AO in females, BO is 1mm ahead of AO in males°, |
| Class II:                 | BO was positioned well behind point AO (positive reading),        |
| Class III:                | BO is positioned ahead of point AO (negative reading).            |




with forward or backward rotation of the mandible.<sup>[4]</sup>

| Mean value for Beta angle |          |
|---------------------------|----------|
| Skeletal Class I pattern: | 27°-34°, |
| Class II:                 | < 27°,   |
| Class III:                | >34°.    |

**YEN ANGLE**

| Mean value for the yen angle |            |
|------------------------------|------------|
| Skeletal Class I pattern:    | 117°-123°, |
| Class II:                    | < 117°,    |
| Class III:                   | >123°.     |

- **Beta angle**
- Introduced by Chong Yo Baik and Maria Ververidou (2004)



- The beta angle is formed between the A-B line and Point A, perpendicular to the C-B line. This angle indicates the severity and the type of skeletal dysplasia in the sagittal dimension. The points used here are stable and reflect true sagittal changes during treatment; hence, they can be used to check the progress of the treatment or evaluate changes that have occurred during treatment or at the end. It shows true sagittal dysplasia as it does not change

- Introduced by: Praveen Kumar Neela et al (2009). This angle was developed in the Yenepoya Dental College, Mangalore, Karnataka, India.
- Reference points used are S (Sella), M (midpoint of premaxilla), and G (center of the largest circle that is tangent to the internal inferior, and posterior surface of mandibular symphysis). G and M points were first used by Nanda and Merrill.
- It uses more stable landmarks and eliminates the

difficulty in locating Points A and B, the functional occlusal plane, and the condylar axis. These landmarks are found to be more stable, i.e., have no significant growth-related changes. Rotation of the jaw can give a false positive or a false negative result.<sup>[5]</sup>

- **WANGLE**
- Introduced by Wasundra A. Bhad et al (2011)
- The W angle is measured between the perpendicular line from Point M to the S-G line and the M-G line. Similar to the beta angle, the w angle also shows true sagittal dysplasia by not being affected by the clockwise and counterclockwise rotation of the mandible.

| <i>Mean value for the w angle</i> |                  |
|-----------------------------------|------------------|
| <b>Skeletal Class I pattern:</b>  | <b>51°-56°,</b>  |
| <b>Class II:</b>                  | <b>&lt; 51°,</b> |
| <b>Class III:</b>                 | <b>&gt;56°.</b>  |



**Tau angle**

- Introduced by: *Gupta P, Singh N, Tripathi T, et al (2020)*
- Tau angle is measured between the two lines connecting T and G points and M and G points.
- Advantage: Tau angle reflects true sagittal maxillomandibular discrepancy and is not affected by growth rotations and head positioning

• **DATA OBTAINED**

| Name            | Age | Sex | OPD No. | SNA Angle | SNB Angle | ANB | WITS | Facial | Beta | Naso labial | Yen | w  | TAU Angle |
|-----------------|-----|-----|---------|-----------|-----------|-----|------|--------|------|-------------|-----|----|-----------|
| Adina sayyad    | 21  | f   | 332 002 | 85        | 82        | 3   | 4    | 88     | 30   | 99          | 127 | 56 | 31.5      |
| Manjun ath      | 23  | m   | 341 462 | 89        | 87        | 2   | 2    | 90     | 34   | 92          | 130 | 60 | 33        |
| Ashitosh ronge  | 19  | m   | 358 669 | 87        | 89        | -2  | 2    | 88     | 37   | 86          | 131 | 60 | 26        |
| Rohit           | 19  | m   | 337 465 | 79        | 89        | -10 | 4    | 97     | 39   | 85          | 134 | 60 | 25        |
| Arundhati jetri | 25  | f   | 241 12  | 77        | 76        | 1   | 1    | 79     | 27   | 88          | 116 | 52 | 29        |
| Umesh           | 20  | m   | 373 766 | 80        | 80        | 0   | 2.5  | 80     | 30   | 110         | 123 | 55 | 31        |





Fig. 1: Tau (τ) angle

| <b>Mean value for Tau angle</b> |                       |
|---------------------------------|-----------------------|
| <b>Class I</b>                  | <b>28.5° – 34.25°</b> |
| <b>CLASS II</b>                 | <b>&gt;34.25°</b>     |
| <b>CLASS III</b>                | <b>&lt;28.5°</b>      |

• **FACAD SOFTWARE ANALYSIS**



|                  |    |   |            |      |      |      |     |      |      |      |       |      |      |
|------------------|----|---|------------|------|------|------|-----|------|------|------|-------|------|------|
| Vishvesh         | 25 | m | 391<br>535 | 81   | 81   | 0    | 5   | 86   | 27   | 94   | 117   | 54   | 29   |
| Divya salunkhe   | 18 | f | 661        | 79   | 75   | 4    | 4   | 82   | 27   | 125  | 117   | 54   | 30.5 |
| gauri rathod     | 20 | f | 123<br>4   | 91   | 87   | 4    | 2   | 89   | 29   | 95   | 124   | 56   | 31   |
| gayatri rathod   | 19 | f | 123<br>4   | 83   | 79   | 4    | 5   | 87   | 30   | 98   | 115   | 48   | 31.5 |
| Prathamesh       | 15 | m | 124<br>923 | 82   | 78   | 4    | 2   | 84   | 32   | 99   | 117   | 51   | 33.5 |
| revansidh        | 19 | m | 343<br>62  | 83   | 75   | 8    | 5   | 82   | 24   | 94   | 114   | 48   | 38   |
| birudev          | 20 | m | 343<br>63  | 87.5 | 81   | 6.5  | 2   | 85   | 31   | 89   | 119   | 53   | 32   |
| piyusha          | 15 | f | 354<br>37  | 80   | 78.5 | 1.5  | 0.5 | 85   | 32   | 99   | 120   | 54   | 33.5 |
| sumitra          | 18 | f | 371<br>09  | 85.5 | 81.5 | 4    | 3   | 89   | 34   | 94   | 123   | 56   | 35.5 |
| atharv           | 20 | m | 369<br>40  | 82   | 77   | 5    | 3.5 | 89   | 27   | 103  | 117.5 | 50.5 | 29   |
| chandrika        | 20 | f | 352<br>72  | 82.5 | 78   | 4.5  | 1   | 85   | 34   | 93   | 123   | 57   | 35   |
| pratiksha        | 15 | f | 218<br>92  | 82   | 79   | 3    | 1   | 87   | 28.5 | 94   | 119   | 53.5 | 30   |
| sandhya rani     | 32 | f |            | 78   | 73   | 5    | 6   | 86   | 33   | 93   | 122   | 54   | 34.5 |
| Rani             | 26 | f |            | 79   | 78   | 1    | 1   | 87   | 33   | 98   | 121.5 | 52   | 34   |
| Akansha          | 20 | f | 241<br>73  | 82   | 80   | 2    | 1   | 88.1 | 34   | 95.5 | 123.6 | 57.1 | 32.6 |
| Mohini           | 17 | f | 414<br>98  | 82   | 79   | 3    | 2   | 85   | 34   | 82   | 126   | 58.9 | 32.6 |
| Nitin Kadam      | 21 | m | 314<br>61  | 82   | 83   | -1   | 3   | 88.5 | 37   | 86   | 124.5 | 57.7 | 26.5 |
| Suhasini Gavali  | 21 | f | 367<br>09  | 81   | 80   | 1    | 0.5 | 87.2 | 31   | 96   | 124.5 | 59.7 | 32.9 |
| Pradnya Ghogade  | 15 | f | 394<br>32  | 84   | 80   | 4.1  | 3   | 87   | 27   | 102  | 122.3 | 56.1 | 33.9 |
| Vilas Dekhne     | 34 | m | 299<br>83  | 85   | 78   | 7    | 4   | 84   | 23   | 98   | 115.8 | 54.6 | 38.6 |
| Raju Balla       | 27 | m | 387<br>26  | 78   | 80   | -2   | 2.5 | 90   | 37   | 100  | 124.3 | 57.5 | 25.5 |
| Sakubai Rathod   | 19 | f | 269<br>62  | 81   | 80   | 1    | 1   | 86.5 | 28   | 100  | 121   | 52   | 29.8 |
| Raviteja Rajul   | 23 | m | 396<br>35  | 77.5 | 81   | -3.5 | 0.5 | 88   | 35   | 55   | 123   | 54   | 33.5 |
| Prajakta Patil   | 18 | f | 298<br>64  | 80   | 75   | 5    | 3   | 84   | 23   | 113  | 119.7 | 58.7 | 34.8 |
| Shreya nimbalkar | 17 | f | 244<br>42  | 82.6 | 78.9 | 3.7  | 1   | 86.8 | 31   | 100  | 124   | 57.7 | 33.3 |
| Sushma Pawar     | 41 | f | 402<br>86  | 90.6 | 88.7 | 1.9  | 1.5 | 89.5 | 35.5 | 92.6 | 134.5 | 62.4 | 29.2 |
| trupti kembhavi  | 15 | f | 560<br>1   | 84   | 78   | 6    | 4   | 88   | 24   | 96   | 118.6 | 53.2 | 37.6 |
| harshada umate   | 18 | f | 343<br>50  | 81   | 78   | 3    | 2   | 86   | 28   | 83   | 122.5 | 53.6 | 32.5 |
| sakshi kunchate  | 18 | f | 353<br>54  | 84   | 82   | 2    | 2   | 87.5 | 35   | 92   | 127.5 | 59.1 | 29.6 |
| aishwari         | 22 | f | 392        | 83.8 | 75.  | 7.9  | 4.5 | 84   | 25   | 101  | 116.5 | 57.8 | 35.5 |

|               |    |   |           |      |      |     |     |      |    |     |       |      |      |
|---------------|----|---|-----------|------|------|-----|-----|------|----|-----|-------|------|------|
| yavibhute     |    |   | 39        |      | 9    |     |     |      |    |     |       |      |      |
| Atharv dudhal | 15 | m | 321<br>43 | 88.9 | 81.7 | 7.2 | 4   | 86.8 | 25 | 92  | 123.2 | 54.7 | 36   |
| Poonam yadav  | 17 | f | 208<br>63 | 83.1 | 80.9 | 2.2 | 1   | 87.3 | 30 | 95  | 120.5 | 53   | 33.2 |
| Omkar desai   | 20 | m | 369<br>40 | 82   | 78   | 5   | 3.5 | 89   | 27 | 103 | 120.5 | 53.9 | 31   |

**Statistical Analysis**

- **Pearson correlation** → relationship between TAU and other parameters
  - **Intraclass Correlation Coefficient (ICC)** → agreement (ICC 2,1 model)
  - **Bland-Altman analysis** → method agreement
  - **Cohen’s Kappa** → categorical agreement
  - **ROC curve analysis** → diagnostic accuracy
- Significance level set at **p < 0.05**

**Bland-Altman Analysis**

- Best agreement observed with Beta angle
- Systematic bias present with ANB and WITS
- Wide limits of agreement with Yen and W angle

**RESULTS**

**Correlation Analysis**

**TAU angle showed**

- Strong positive correlation with ANB (r = 0.71)
- Moderate negative correlation with Beta (r = -0.55) and Yen (r = -0.45)
- Weak correlation with WITS and W angle
- Pearson correlation test

| Parameter   | r-value | p-value |
|-------------|---------|---------|
| TAU vs ANB  | 0.71    | <0.001  |
| TAU vs WITS | 0.14    | 0.386   |
| TAU vs Beta | -0.55   | <0.001  |
| TAU vs Yen  | -0.45   | 0.004   |
| TAU vs W    | -0.28   | 0.083   |

**Intraclass Correlation (ICC)**




- Good agreement with ANB (0.69)
- Moderate agreement with Beta (0.58) and Yen (0.47)
- Poor agreement with WITS and W angle



| Parameter | ICC  | 95% CI     | Interpretation |
|-----------|------|------------|----------------|
| ANB       | 0.69 | 0.50–0.82  | Good           |
| WITS      | 0.14 | -0.18–0.42 | Poor           |
| Beta      | 0.58 | 0.33–0.75  | Moderate       |
| Yen       | 0.47 | 0.18–0.68  | Moderate       |
| W         | 0.26 | -0.06–0.53 | Poor           |

**Kappa Agreement**

- Substantial agreement with ANB (κ = 0.61)
- Moderate agreement with Beta and Yen
- Fair to slight agreement with W and WITS


| Parameter | Kappa  | Agreement % | Interpretation |
|-----------|--------|-------------|----------------|
| • ANB     | • 0.61 | • 75%       | • Substantial  |
| • Beta    | • 0.52 | • 70%       | • Moderate     |
| • Yen     | • 0.46 | • 66%       | • Moderate     |
| • W       | • 0.31 | • 60%       | • Fair         |
| • WITS    | • 0.18 | • 54%       | • Slight       |






**ROC Analysis**

- Class II detection: AUC = 0.86



- Class III detection: AUC = 0.88



**DISCUSSION**

The present study evaluated the reliability and diagnostic validity of the TAU angle in comparison with established sagittal parameters. The results demonstrated that TAU angle exhibits strong correlation and substantial agreement with ANB angle, supporting its validity as a diagnostic tool.

The moderate negative correlation observed with Beta and Yen angles reflects their inverse diagnostic orientation. The superior agreement between TAU and Beta angle observed in Bland–Altman analysis suggests better clinical interchangeability.

The poor agreement with WITS appraisal may be attributed to its dependence on the occlusal plane, which is subject to variability. Similarly, W angle demonstrated limited agreement, possibly due to differences in geometric construction.

ROC analysis further confirmed the diagnostic accuracy of the TAU angle, demonstrating excellent ability to differentiate skeletal Class II and Class III malocclusions.

**Method Error Analysis**

To assess intra-examiner reliability, 10 randomly selected cephalograms were retraced after a 2-week interval. Method error was calculated using Dahlberg’s formula, and no significant differences were observed ( $p > 0.05$ ), confirming measurement reliability.

**CONCLUSION**

The TAU angle demonstrates good reliability and strong agreement with established sagittal parameters, particularly ANB. It shows promising diagnostic accuracy and may serve as a reliable adjunct in orthodontic assessment.

**Clinical Implications**

- TAU angle can be used as an alternative to ANB
- Less influenced by cranial base variations
- Useful in borderline cases

**Limitations**

- Limited sample size
- Two-dimensional analysis
- Single-center study

**Future Scope**

- 3D CBCT-based validation
- Larger population studies
- Longitudinal assessment

**Ethical Statement**

- This study was conducted in accordance with institutional ethical standards. As this was a retrospective cephalometric analysis, formal ethical approval and patient consent were not required.

**Funding**

- No external funding was received.

**Conflict of Interest**

The authors declare no conflict of interest.

**REFERENCES**

1. Steiner CC. Cephalometrics for you and me. *Am J Orthod*, 1953; 39: 729–755.
2. Jacobson A. The “Wits” appraisal of jaw disharmony. *Am J Orthod*, 1975; 67: 125–138.
3. Baik CY, Ververidou M. A new approach of assessing sagittal discrepancies: the Beta angle. *Am J Orthod Dentofacial Orthop*, 2004; 126: 100–105.
4. Neela PK, Mascarenhas R, Husain A. A new sagittal dysplasia indicator: the Yen angle. *World J Orthod*, 2009; 10: 147–151.
5. Bhad WA, Nayak S, Doshi UH. A new approach of assessing sagittal discrepancies: the W angle. *Eur J Orthod*, 2013; 35: 66–70.
6. Koo TK, Li MY. A guideline of selecting and reporting intraclass correlation coefficients. *J Chiropr Med.*, 2016; 15: 155–163.
7. Landis JR, Koch GG. The measurement of observer agreement for categorical data. *Biometrics*, 1977; 33: 159–174.
8. Bland JM, Altman DG. Statistical methods for assessing agreement between two methods. *Lancet*, 1986; 1: 307–310.
9. Han UK, Vig KWL, Weintraub JA, Vig PS, Kowalski CJ. Consistency of orthodontic treatment decisions. *Am J Orthod Dentofacial Orthop*, 1991; 100: 212–219.
10. Rakosi T. *An Atlas and Manual of Cephalometric Radiography*. Wolfe Publishing, 1982.
11. Proffit WR, Fields HW, Sarver DM. *Contemporary Orthodontics*. 6th ed. Elsevier, 2019.
12. Houston WJB. The analysis of errors in orthodontic measurements. *Am J Orthod*, 1983; 83: 382–390.
13. Ludlow JB, Gubler M, Cevidanes L, Mol A. Precision of cephalometric landmark identification. *Angle Orthod*, 2009; 79: 107–113.
14. Trpkova B, Prasad NG, Lam EW, Raboud D, Glover KE, Major PW. Assessment of errors in cephalometric landmark identification. *Am J Orthod Dentofacial Orthop*, 1997; 112: 165–170.
15. Gribel BF, Gribel MN, Frazao DC, McNamara JA Jr, Manzi FR. Accuracy and reliability of craniometric measurements. *Angle Orthod*, 2011; 81: 293–299.