



THE CEYLON MEDICAL JOURNAL

Established 1887

The Official Publication of the
Sri Lanka Medical Association

Volume 48, No.2, June 2003

Quarterly ISSN 0009-0875

All communications
should be addressed to
The Editors, CMJ

Editor Emeritus

Chris G Uragoda MD, FRCP

Editors

Colvin Goonaratna FRCP, PhD
Janaka de Silva DPhil, FRCP

Assistant Editors

Dennis Aloysius MBBS, FCGP
D N Atukorala MD, FRCP
Damani de Silva MD, MRCPsych
Dulani Gunasekara MD, MRCP
Harshalal R Seneviratne DM, FRCOG
Kolitha Sellahewa MD, FCCP
Ranjan Dias MS, FRCS
Saman Gunatilake MD, FRCP
Sarath Gamini de Silva MD, FRCP
Tissa Vitarana MD, PhD
Vishvarani Wanigasekera MRCP, FRCA

International Advisory Board

Richard Smith FRCP
London, UK.

Raja Bandaranayake FRACS, PhD
New South Wales, Australia.

Kasuko Ito MD
Gifu, Japan.

R K Tandon MD, PhD
New Delhi, India.

Continued overleaf

Leading article

Detection of the small for gestational age fetus

A reliable SFH chart should be developed for Sri Lanka, and serial SFH measurements emphasised.

A fetus is generally considered to be small for gestational age (SGA) if its birth weight is below the 10th centile for its specific gestational age. The majority of SGA fetuses are constitutionally small and healthy, and when delivered at term do not carry any appreciable increased morbidity or mortality (1,2). Some of these fetuses however have had fetal growth restriction (FGR) and failed to achieve their full growth potential. They not only have a poor perinatal outcome (3,4) but also carry an increased risk of neonatal complications (5,6), impaired neurodevelopment (7), and possibly even type 2 diabetes mellitus and hypertension in adult life (8).

Over the years a great deal of research has been done to identify improved methods of early detection of FGR. Ultrasound biometry has become increasingly popular and serial measurement of growth velocity using fetal abdominal circumference, and estimated fetal weight have been found to be the best indices to predict FGR (9). It is important to note that only serial (not single) measurements can predict FGR and poor perinatal outcome (9,10). The reference chart used for ultrasound biometry should be based on longitudinal growth studies and not on cross sectional data (11,12).

Routine ultrasonography after 24 weeks gestation in low-risk pregnancy has not been useful in improving perinatal outcome (13). Even biophysical tests such as cardiotocography, amniotic fluid volume, biophysical scoring and ultrasound Doppler flow velocimetry have only a limited ability to detect FGR in low risk populations (14). Ultrasound and Doppler flow velocimetry is not freely available except in tertiary care centres. Hence it is important to detect the SGA fetus as early as possible clinically. This would enable further evaluation and monitoring of the fetuses to detect FGR.

Studies have suggested that up to 70% of SGA fetuses may not be detected by inspection and palpation of the abdomen alone (15,16). Although measurement of the symphysio-fundal height (SFH) was introduced more than three decades ago, the clinical value of this measurement has been questioned because of wide variation (17).

SFH measurements have specificities of up to 94% in the detection of SGA fetuses (18,19). Although these studies have reported sensitivities up to 84%, other large studies have shown sensitivities as low as 27% (20). The low sensitivity, high false positive rates and significant intra- and inter-observer

Stephen Lock MD, FRCP
London, UK.

Samiran Nundy FRCS, FRCP
New Delhi, India.

N Medappa MD
New Delhi, India.

Jane Smith BA, MSc
London, UK.

S K Sarin MD, DM
New Delhi, India.

David Warrell MD, FRCP
Oxford, UK.

**Advisory Board for
Statistics and Epidemiology**

R O Thattai MSc, PhD

Lalini Rajapakse MD, MSc

Kumudu Wijewardene MBBS, MD

Published by

The Sri Lanka Medical
Association
Wijerama House
6, Wijerama Mawatha
Colombo 7
SRI LANKA

Telephone +94 1 693324

Fax +94 1 698802

Internet home page

<http://www.slmaonline.org/cmj>

E-mail SLMA@eureka.lk

Printed by

S Devendra
Ananda Press
82/5, Sir Ratnajothi Saravanamuttu
Mawatha, Colombo 13
Sri Lanka

Telephone +94 1 435975

Fax +94 1 385039

E-mail anpress@sltnet.lk

variations make SFH *per se* an unreliable index for the detection of SGA fetuses (20,21,22), but in large low-risk populations it could be used as a screening test that should be supplemented with serial ultrasound biometry. It is often stated that the "SFH in centimetres (cm) should give an estimation of gestational age in weeks, ie ± 2 cm from 20-38 weeks" (23). Unfortunately this encourages over emphasis on a single SFH measurement, and the 'point to point' conversion of a single SFH measurement into gestational size in weeks. Even special tape measures, calibrated with the 'cut off levels' of SFH calculated for two week intervals have been used (24). As with ultrasound biometry, a single measurement of SFH is of limited value. Serial measurements showing the trend can improve sensitivity and specificity of SFH measurements in detecting SGA fetuses (25). The technique of SFH measurement is also important. The variable point of the fundus should be first identified, and the measurement taken to the fixed point (the symphysis pubis), with the cm values hidden from the examiner (26). It has been suggested that faulty technique may be the reason for the low sensitivity of SFH measurements in some studies (18,27).

Low birth weight of a baby could be the result of physiological variables such as its sex, and the mother's body mass index, parity and ethnicity. So reference charts of SFH measurements should be 'customised' and adjusted for such variables. Customised SFH charts should be based on longitudinal growth studies and not on cross sectional data. Use of such charts can improve the sensitivity of SFH measurements in detecting SGA fetuses (28). Multiple serial assessments using different methods such as abdominal palpation, SFH measurements, ultrasound biometry and biophysical tests may help the clinician to detect the at-risk FGR fetus early.

In this issue of the *CMJ* Senanayake and colleagues report (p 43-45) that approximately three out of four women attending peripheral as well as tertiary care antenatal clinics (ANC) in Colombo had their SFH measured in cm and documented. It is encouraging to note that more than 70% of SGA fetuses had been detected by SFH measurement in the peripheral ANCs. However, since none of the service providers at the peripheral ANC had plotted the measurements on the SFH chart provided in the pregnancy record card, they probably had not appreciated fully the trend of the SFH measurements. This is probably why only one third of these women whose SFH measurements were found to be less than expected had been referred for further evaluation in a tertiary centre. This study also found that in the tertiary care clinic detection rate of SGA was significantly lower (54%) and later in pregnancy than in the peripheral ANC. This is probably because intern house officers or senior house officers with relatively less clinical experience attend on these women after their 'booking visit' and they have to attend on about 65 patients daily. If these doctors' clinical skills could be improved and the doctor: mother ratio reduced, the detection rate may improve. The availability of ultrasonography would have encouraged them to arrange for further evaluation of the women who had low SFH measurements.

As mentioned by Senanayake and colleagues the SFH chart in the pregnancy record card used in peripheral ANCs is unsatisfactory. A reliable customised SFH chart should be developed for Sri Lanka. This will require a properly conducted multicentre longitudinal study of SFH measurements during pregnancy. Once this is developed the importance of plotting SFH measurements on the chart and studying the trend (rather than a single measurement) should be emphasised to all health care personnel who conduct antenatal clinics in the periphery as well as in tertiary care centres. This is likely to improve the detection of SGA fetuses.

References

1. Ott WJ. The diagnosis of altered fetal growth. *Obstetrics and Gynaecology Clinics of North America* 1988; **15**: 237-63.
2. Jones RA, Robertson NR. Small for dates babies: are they really a problem? *Archives of Diseases of Children* 1986; **61**: 877-80.
3. Beattie RB, Johnson P. Practical assessment of neonatal nutritional status beyond birthweight: An imperative for the 1990s. *British Journal of Obstetrics and Gynaecology* 1994; **101**: 842-6.
4. Gardosi J, Mul T, Mongelli M, Fagan D. Analysis of birthweight and gestational age in antepartum stillbirths. *British Journal of Obstetrics and Gynaecology* 1998; **105**: 524-30.
5. McIntire D, Bloom SL, Casey BM, Leveno KJ. Birthweight in relation to morbidity and mortality among newborn infants. *New England Journal of Medicine* 1999; **340**: 1234-8.
6. Fonseka P, Wijewardena K, de Silva DGH, Goonaratna C, Wijesiri WA. Neonatal and post-neonatal mortality in Galle district. *Ceylon Medical Journal* 1994; **39**: 82-5.
7. Taylor DJ, Howie PW. Fetal growth achievement and neurodevelopment disability. *British Journal of Obstetrics and Gynaecology* 1989; **96**: 789-94.
8. Barker DJ, Ghickman PD, Godfrey KM, Harrding JE, Owens JA, Robinson JS. Fetal nutrition and cardiovascular disease in adult life. *Lancet* 1993; **341**: 938-41.
9. Chang TC, Robson SC, Boys RJ, Spencer JA. Prediction of the small for gestational age infant: which ultrasonic measurement is best? *Obstetrics and Gynaecology* 1992; **80**: 1030-8.
10. De Jong CL, Francis A, Van Geijn HP, Gardosi J. Fetal growth rate and adverse prenatal events. *Ultrasound in Obstetrics and Gynaecology* 1999; **13**: 86-9.
11. Mongelli M, Gardosi J. Longitudinal study of fetal growth in subgroups of a low-risk population. *Ultrasound in Obstetrics and Gynaecology* 1995; **6**: 340-4.
12. Owen P, Donnet ML, Ogston SA, Christie AD, Howie PW, Patel NB. Standards for ultrasound fetal growth velocity. *British Journal of Obstetrics and Gynaecology* 1996; **103**: 60-9.
13. Bricker L, Neilson JP. Routine ultrasound in late pregnancy (after 24 weeks' gestation). *Cochrane Database Systematic Review* 2000; CD 001451.
14. Coomaraswamy A, Fisk NM, Gee H, Robson SC. The investigation and management of the small-for-gestational-age fetus. Green Top Guideline No. 31. London: Royal College of Obstetricians and Gynaecologists, 2002: 4-5.
15. Hall MH, Chang PK, MacGillivray I. Is routine antenatal care worthwhile? *Lancet* 1980; **2**: 78-80.
16. Rosenberg K, Grant JM, Hepburn M. Antenatal detection of growth retardation: actual practice in a large maternity hospital. *British Journal of Obstetrics and Gynaecology* 1982; **89**: 12-5.
17. Beazly JM, Underhill RA. Fallacy of the fundal height. *British Medical Journal* 1970; **4**: 404-6.
18. Mathai M, Jairaj P, Muthurathna S. Screening for light-for-gestational age infants: a comparison of three simple measurements. *British Journal of Obstetrics and Gynaecology* 1987; **94**: 217-21.
19. Calvert JP, Crean EE, Newcombe RG, Pearson JF. Antenatal screening by measurement of symphysis-fundal height. *British Medical Journal* 1982; **285**: 846-9.
20. Persson B, Stangenberg M, Lunell NO, Brodin, Holmberg NG, Vaclavincova V. Prediction of size of infants at birth by measurement of symphysis fundus height. *British Journal of Obstetrics and Gynaecology* 1986; **93**: 206-11.
21. Neilson JP. Symphysis-fundal height measurement in pregnancy. *Cochrane Database Systematic Review* 2000; CD 000944.
22. Bailey SM, Purnima S, Grant JM. A comparison of three methods of assessing inter-observer variations applied to measurement of the symphysis-fundal height. *British Journal of Obstetrics and Gynaecology* 1989; **96**: 1266-71.
23. Campbell S, Lees C. eds. *Obstetrics by ten teachers*. 17th edition. London: Arnold, 2000: 6-10.
24. Mathai M. Prediction as small-for-gestational age infant using a specially calibrated tape measure. *British Journal of Obstetrics and Gynaecology* 1988; **95**: 313-4.
25. Pearce JM, Campbell S. A comparison of symphysis-fundal height and ultrasound as screening tests for light-for-gestational age infants. *British Journal of Obstetrics and Gynaecology* 1987; **94**: 100-4.
26. Gardosi JO, Mongelli JM, Mull T. Intrauterine growth retardation. *Baillieres Clinical Obstetrics and Gynaecology* 1995; **9**: 445-63.
27. Rosenberg K, Grant JM, Tweedie P, Aitchison T, Gallagher F. Measurement of fundal height as a screening test for fetal growth retardation. *British Journal of Obstetrics and Gynaecology* 1982; **89**: 447-50.
28. Gardosi J, Francis A. Controlled trial of fundal height measurement plotted on customized antenatal growth charts. *British Journal of Obstetrics and Gynaecology* 1999; **106**: 309-17.

Malik Goonewardene, Professor and Head, Department of Obstetrics and Gynaecology, Faculty of Medicine, University of Ruhuna.