

Effect of light intensities on the stress responses and toxin production of selected Microcystin producing Cyanobacteria

HMSAT. Gunathilaka¹, WRP Wijesinghe², DN Magana-Arachchi¹

Introduction and Objectives: Low to moderate light intensity is a crucial determinant in cyanobacterial proliferation. This study aims to assess the comparative effects of two different light intensities on microcystin production and stress responses in five different cyanobacterial species.

Methods: Cyanobacterial isolates (*Microcystis* sp., *Fischerella* sp., *Nostoc* sp., *Pseudoanabaena* sp., *Leptolyngbya* sp.) were exposed to 0 $\mu\text{mol m}^{-2} \text{s}^{-1}$ and 50 $\mu\text{mol m}^{-2} \text{s}^{-1}$ light intensities. Samples in 50 $\mu\text{mol m}^{-2} \text{s}^{-1}$ were exposed to cycles of 12 h light:12 h dark. Total soluble Protein Content (TPC), Ascorbate Peroxidase activity analysis (APX), and Microcystin (MC) toxin content were analysed over 20 days. The TPC was measured using the Bradford method; absorbance was measured at 595 nm using a microplate reader (FLUOstar Omega). For the APX assay, 1 mL of extract supernatant was measured for absorbance at 290 nm for 3 minutes. For MCs, the extracted toxin was analyzed by High-Performance Liquid Chromatography (Ultimate 3000 HPLC system; VWD detector; C₁₈ column) with reference to the MC standard (SIGMA ALDRICH 33578). An Analysis of Variance was conducted to assess the statistical significance of the data.

Results: TPC levels of each condition were nearly the same as the initial (501.14+154.30 $\mu\text{g/ml}$) ($p > 0.05$). APX activity was significantly increased in high-light intensities (9.75+ 2.28 $\text{nmol min}^{-1} \text{mg}^{-1}$) compared to lower intensities (1.24+2.05 $\text{nmol min}^{-1} \text{mg}^{-1}$) ($p < 0.05$). The total MC concentration showed a significant 35.8% rise in high-light intensities than the lower intensities ($p < 0.05$). In high-light intensity, the highest concentration was recorded from *Fischerella* sp. (0.9206+0.08 mg/l), and the lowest from *Microcystis* sp. (0.2563+0.12 mg/l). In low-light intensity, only the MC-LR variant was detected from *Fischerella* sp. (0.2591+0.09 mg/l) and *Pseudoanabaena* sp. (0.3430+0.16 mg/l). Both MC-YR and MC-LR toxin variants were present in high-light-intensity conditions.

Conclusions: The results indicate APX activity and MC production increase with high-light-intensity. *Fischerella* sp. showed the highest toxin production under high light, while *Microcystis* sp. exhibited the lowest. MC-YR and MC-LR toxin variants were present in high-light conditions. Hence, cyanobacterial toxin production under light intensity can be used to predict their health risk in freshwater bodies.

Keywords: Cyanobacteria, Cyanotoxin, Light intensity, Oxidative stress

¹Molecular Microbiology and Human Diseases Programme, National Institute of Fundamental Studies, Kandy, Sri Lanka

²Department of Botany, Faculty of Science, University of Peradeniya, Sri Lanka

Address for correspondence: Prof. DN. Magana-Arachchi, Molecular Microbiology and Human Diseases Programme, National Institute of Fundamental Studies, Kandy, Sri Lanka. Telephone: +94772865367;

Email: dhammika.ma@nifs.ac.lk;  <https://orcid.org/0000-0001-5825-4626>