

Baltimore, MD. Ages ranged from 7.0 to 15.5 years. Using 2 x 2 multivariate factorial analysis of variance (MANOVA), which characterized subjects for presence or absence of TS and ADHD, and various combinations, the effects of TS status, ADHD status, and the interaction of these two factors on frontal lobe white and gray matter volumes were determined. In subjects with TS, the right frontal lobe had a larger proportion of white matter, and the right frontal white matter volume was increased. ADHD was associated with a reduction in frontal lobe volumes that was attributable to decreased gray matter, especially in the left frontal lobe. The findings support a frontostriatal pathophysiology of Tourette syndrome. (Fredericksen KA, Cutting LE, Kates WR et al. Disproportionate increases of white matter in right frontal lobe in Tourette syndrome. Neurology January (1 of 2) 2002;58:85-89). (Reprints: Dr Walter E Kaufmann, Department of Developmental Cognitive Neurology, The Kennedy Krieger Institute, 707 N Broadway, Room 522, Baltimore, MD 21205).

COMMENT. An increase in white matter in the right frontal lobe of patients with Tourette syndrome is in keeping with previous reports by the same authors of an enlarged rostrum of the corpus callosum associated with TS (Baumgardner TL et al. 1996; see Progress in Pediatric Neurology III, PNB Publ, 1997;pp322-323). In contrast, boys with ADHD were found to have a significant decrease in size of the corpus callosum rostral body. The effects of ADHD and TS on corpus callosum size were statistically independent.

ATTENTION DEFICIT DISORDER

DOPAMINE RECEPTOR AVAILABILITY IN ADHD

Striatal dopamine (D2) receptor availability was determined by iodobenzamide brain SPECT, before and 3 months after methylphenidate (MPH) therapy, in 9 children (mean age, 9.8 years) with attention deficit hyperactivity disorder (ADHD) examined at Gazi University, Ankara, Turkey. Specific D2 receptor binding ratios in ADHD subjects at baseline were higher than those reported in healthy young adults. Severity of ADHD behavioral symptoms was correlated with D2 receptor levels; the higher the Conners Teacher Rating Scale scores, the higher the baseline D2 availability in the putamen and caudate. D2 receptor availability was significantly reduced in all four regions of the striatum following MPH therapy. The higher the baseline D2 levels, the greater the behavioral response to MPH (as determined by improved hyperactivity and Conners Teacher Rating scores), whereas the response in attention deficit scores was not related to baseline D2 levels. D2 receptor levels in patients with ADHD may predict the behavioral response to MPH. (Ilgin N, Senol S, Gucuyener K et al. Is increased D2 receptor availability associated with response to stimulant medication in ADHD? Dev Med Child Neurol Nov 2001;43:755-760). (Respond: Dr N Ilgin, PK 61 Bahcelievler, 06500 Ankara, Turkey).

COMMENT. D2 receptor levels are higher than normal in untreated children with ADHD, and are reduced to near-normal values following treatment with methylphenidate. The higher the pre-treatment D2 receptor availability, the greater the severity and number of ADHD behavioral symptoms and the better the behavioral response to therapy. These findings are in accord with previous reports of clinical predictors of response to MPH therapy in ADHD (Millichap JG, Ann NY Acad Sci 1973; in Attention Deficit Hyperactivity and Learning Disorders, PNB Publ, 2001). Children who are most active are most likely to respond to MPH.