

POSTASPHYXIAL HYPOXIC-ISCHEMIC ENCEPHALOPATHY

Models of association between risk factors and severe adverse outcome within 4 hours of birth were determined in 178 infants with postasphyxial hypoxic-ischemic encephalopathy (PA-HIE) admitted consecutively between 1985 and 1992 to the regional referral neonatal ICU at the Hospital for Sick Children, Toronto, Canada. Of 165 infants with known outcomes, 88 (53%) had severe adverse outcome, 48 died (42 in the first month), and 40 had severe impairment and microcephaly at 1 year. Rates of PA-HIE and severe adverse outcome were 0.4 and 0.2/1000 births; the rate of PA-cerebral palsy was 0.1/1000 births. Delayed onset of breathing, administration of chest compressions, and seizures at less than 4 hours of age were most predictive of adverse outcome. (Ekert P, Perlman M, Steinlin M, Hao Y. Predicting the outcome of postasphyxial hypoxic-ischemic encephalopathy within 4 hours of birth. J Pediatr Oct 1997;131:613-617). (Reprints: Max Perlman MB, FRCP, Division of Neonatology, Department of Pediatrics, Hospital for Sick Children, 555 University Ave, Toronto, Ontario, M5G 1X8, Canada).

COMMENT. Three independent predictors in the first 4 hours after birth may be used as clinical markers of severe adverse outcome of asphyxia and HIE: apnea duration, chest compressions, age at onset of seizures. Measurements of CBF with PET during the neonatal period may also be predictive of childhood neurologic outcome.

Neonatal cerebral blood flow (CBF) and childhood IQ and neurologic outcome. A significant negative correlation between neonatal CBF and childhood IQ measured at ages 4 to 12 years was found in a study at Washington University School of Medicine, St Louis, MO. The mean neonatal CBF in 8 with abnormal childhood neurologic outcome was higher than in 8 with normal neurologic evaluations. Higher CBF correlated with lower IQ. (Rosenbaum JL, Almlı CR, Yundt KD, Altman DI, Powers WJ. Higher neonatal cerebral blood flow correlates with worse childhood neurologic outcome. Neurology Oct 1997;49:1035-1041).

CONGENITAL PORENCEPHALY AND HIPPOCAMPAL SCLEROSIS

MRI volumetric findings were correlated with seizure patterns and EEGs in 14 patients with intractable seizures, porencephaly and hippocampal sclerosis (HS) in a study at the University of Alabama, Birmingham, AL. Psychoparetic complex partial seizures (CPS) occurred in 10, simple partial seizures in 3, and generalized TC seizures in 1. EEGs showed ictal or interictal temporal localization in 9 (64%) patients with CPS. Porencephaly was distant from the temporal area and in the middle cerebral artery distribution in 8; it was related to the posterior cerebral in only 3. Hippocampal formation atrophy in 13 (93%) patients was concordant with CPS and EEG temporal localization in 70% cases. Ten had amygdala atrophy, concurrent with HS in 57%. Two with HS were seizure free after temporal lobectomy. A common ischemic pathogenesis is proposed for the dual pathology involving both porencephaly and mesial temporal sclerosis. HS is the most likely origin for CPS in patients with EEG temporal localization. (Ho SS, Kuzniecky RI, Gilliam F, Faught E, Bebin M, Morawetz R. Congenital porencephaly and hippocampal sclerosis. Clinical features and epileptic spectrum. Neurology Nov 1997;49:1382-1388). (Reprints: Dr Ruben I Kuzniecky, Department of Neurology, UAB Station, Birmingham, AL 35294).

COMMENT. Patients presenting with intractable complex partial seizures