



**RISK FACTORS OF ORAL MUCOSAL LESIONS AMONG ELDERLY PATIENTS AT
DENTAL SPECIALIST CENTER IN AL-KARAMAH HOSPITAL – BAGHDAD/IRAQ**

¹*Dr. Mina Hasan Auda, ²Dr. Ali Abid Hawiya, ³Dr. Amer Abdulkarim Abdul Rahman and ⁴Dr. Zainab Saad Al-Gherany

¹M. B. Ch. B, Higher Diploma in Dentistry, College of Dentistry - University Baghdad.

²M. B. Ch. B, Higher Diploma in Dentistry, College of Dentistry - University Baghdad.

³M. B. Ch. B, Higher Diploma in Dentistry, College of Dentistry - University Baghdad.

⁴M. B. Ch. B, M.Sc. College of Dentistry - University Baghdad.

^{1,2,3}Al Karama Specialized Dental Center-Baghdad /Iraq.

⁴Medical City of Baghdad – Iraq.

*Corresponding Author: Dr. Mina Hasan Auda

M. B. Ch. B, Higher Diploma in Dentistry, College of Dentistry- University Baghdad.

Article Received on 22/01/2020

Article Revised on 12/02/2021

Article Accepted on 02/03/2021

ABSTRACT

This study aimed to determine the prevalence of oral lesions in terms of sex, age, educational status, medication use, systemic diseases, the duration of denture use and tobacco or alcohol use.

A total of 140 voluntary patients (74 males and 66 females), aged ≥ 60 years, were interviewed by one investigator for demographic data, systemic diseases, tobacco or alcohol use, denture use and the duration of denture use.

The majority of the participants (87.1%) had one or more oral mucosal lesions. The prevalence of oral mucosal lesions was 45.9% in males and 40.1% in females ($P=0.97$). The most common oral mucosal lesion was a sublingual varicosity in both males and females. A statistically significant difference was observed between the three age groups (60–64, 65–69, and ≥ 70 years) in regard to the prevalence of oral mucosal lesions ($P=0.02$). There was a significant relationship between the presence of systemic diseases and oral mucosal lesions ($P=0.001$).

It can be concluded that although the majority of oral mucosal lesions detected in the present study were benign, there were some patients with premalignant and malignant lesions. Therefore, periodic oral examinations for detection of precancerous and cancerous lesions are important, especially among the elderly, smokers and denture users.

KEYWORDS: Chronic disease, dentures, elderly, oral mucosa.

INTRODUCTION

Relationship has been reported between oral mucosal lesions and aging.^[1] Due to reduced immunologic reactivity, impaired DNA repair capacity, impaired carcinogen metabolism and atrophy of oral tissues, particularly of the oral epithelium and the salivary glands, oral mucosal conditions tend to develop more frequently and more rapidly in aging populations.^[2]

Age has an important influence on the prevalence of oral mucosal lesions. The prevalence of oral mucosal lesions has been shown to be higher in older subjects than in younger individuals.^[1-6] However, age alone is not the only factor and other findings such as trauma, medications and oral and denture hygiene might affect

the development of oral mucosal lesions. Oral mucosal lesions usually occur due to systemic diseases, nutritional disorders, medication side effects or wearing ill-fitting dentures in the elderly.^[7]

In addition to age, factors such as gender, educational level, smoking and systemic diseases might pave the way for oral lesions. Since the level of education is an effective element in oral hygiene maintenance, it might be important in the formation of oral lesions. Oral health status is important for the quality of life, owing to physical, social and psychological factors. Early detection and prevention of oral lesions by dental practitioners can improve the quality of life of this population and aid in the attainment of quality aging.

The prevalence of oral mucosal conditions is an important parameter in evaluating the oral health of the elderly. Therefore, the aim of this study was to determine the prevalence of oral lesions in relation to sex, age, medication use, systemic diseases, the duration of denture usage, education level, oral hygiene status and tobacco and alcohol use.

METHODS

The study protocol was approved by the Ethics Committee of Al-Karama teaching hospital. Written informed consent was obtained from all the participants. The subjects were 140 volunteer patients aged ≥ 60 years with various dental complaints and admitted to Al Karama Specialized Dental Center at Al-Karama teaching hospital-Baghdad/Iraq during the period from March 2008 to April 2009. One oral medicine specialist interviewed all the participants for demographic data, systemic diseases, tobacco or alcohol use, use of dentures and the duration of denture use.

Intraoral examinations of the patients were performed by one investigator. As a part of the routine oral examinations, dental and medical anamnesis were taken from the patients. Tongue depressors and mirrors were used to visualize the oral mucosa with palpation being performed where indicated. Radiological examinations were also performed. In the presence of oral mucosal lesions, the diagnosis was made on the basis of the dental/medical history and clinical features according to the World Health Organization Guidelines and Oral Pathology, Clinical Pathologic Correlations.^[8,9] As for smoking, the participants were classified into three categories according to their smoking habits as non-smokers, former smokers and smokers. Oral hygiene was evaluated according to the Oral Hygiene Index (OHI) (Greene and Vermillion) and the DMFT index. The level of education of the participants was classified into six categories as illiterate, literate, primary school, middle school, high school and university. The classification of denture types was performed according to Jankittivong *et al.*^[7] Denture wearers were divided into complete denture wearer (CDW) and partial denture wearer (PDW) groups. The CDW group was further divided into sub-groups, including maxillary and mandibular complete dentures, and a maxillary or mandibular complete denture alone or in combination with crowns and/or bridges. The PDW group included patients who had either a partial denture alone or in combination with crowns and/or bridges. The duration of denture use was classified into four groups according to Jorge *et al.*^[10] as follows: < 5 , 6–10, 11–20, and ≥ 20 years of use. After preparation of a dental treatment plan, the patients were referred to other dental departments according to their requirements.^[10]

Statistical analysis

SPSS 20.0 for Windows (SPSS Inc., Chicago, USA) was used to analyze data. Chi-square test was used to test the

differences. Statistical significance was defined as $P < 0.05$.

RESULTS

The study sample of 140 elderly patients included 74 (52.8%) men and 66 (47.2%) women. The patients were divided into three age groups: 60–64 (35.0%), 65–69 (32.1%), and ≥ 70 years of age (32.9%).

The majority of participants (43.1%) were primary school graduates. The education level did not significantly affect the presence of oral mucosal lesions ($P = 0.849$).

In terms of bad habits, 47.2% of the participants never used cigarettes, 33.6% and 19.2% were former smokers and smokers, respectively. The majority of smokers (75.0%) were males and 85.0% of smokers had smoked for more than 10 years. Among the patients, 1.1% declared having an alcohol habit and were defined as drinkers.

In regard to dentures, 65.0% of the study population wore dentures. In terms of sex, the percentage of males using prostheses (55.0%) was higher than females (45.0%).

The percentage of participants with maxillary and mandibular complete dentures was 49 (35%) and 34 (24.3%), respectively. Maxillary and mandibular partial denture wearers comprised 35 (25.0%) and 44 (31.4%) of the study subjects, respectively.

The majority of the subjects with crowns and/or bridges (34.5%) and removable dentures (30.3%) had been using them for ≤ 5 years. The majority of the elderly with removable dentures (34.7%) were illiterate. The duration of denture use was significantly different in terms of the educational status ($P = 0.01$).

While 77.8% of the elderly population had natural teeth and 22.2% were edentulous. In terms of sex, the percentage of females who were edentulous was 25.7%, but this figure was 16.2% in males. However, 25.7% of the participants had functional dentition (≥ 20 teeth). The percentage of men with ≥ 20 teeth (31.1%) was higher than women (19.7%). A total of 117 (83.6%) patients had poor oral hygiene according to the OHI. The majority of subjects with poor oral hygiene (84.6%) had oral mucosal lesions. There was no significant relationship between oral hygiene and the presence of oral mucosal lesions ($P = 0.94$).

Approximately 87.1% of the participants had one or more oral mucosal lesions. The prevalence of oral mucosal lesions was 45.9% in males and 40.1% in females, with no significant difference ($P = 0.97$). The prevalence of oral mucosal lesions in the three age groups was as follows: 25.0% in the 60–64-year group, 35.1% in the 65–69-year group, and 39.9% in the ≥ 70 -

year group. A significant difference was observed between the three elderly age groups regarding the prevalence of oral mucosal lesions ($P=0.02$).

Red-blue lesions were observed with the highest frequency (45.0%) both in the study population and in both sexes (34.8% for women, 40.5% for men). The most common type of oral mucosal lesion was a sub-lingual varicosity in both males and females. The second lesion frequently observed was fissured tongue (27.3 % in females, 35.1% in males). While 9% of the patients with mucosal lesions were smokers, 55% did not smoke and 36% had stopped smoking ($P=0.001$). The most common lesions were sublingual varicosity (10.0%) and denture stomatitis (2.1%) in smokers.

When the determined malignant lesions included adenocarcinoma, melanoma and squamous cell carcinoma, lichen planus was determined as a premalignant lesion in both males and females.

The incidence of systemic diseases was high. Cardiovascular diseases were found in 89 (63.5%) participants and often identified as a systemic disease group. The majority of the elderly with systemic diseases had mucosal lesions. There was a significant relationship between the presence of systemic diseases and oral mucosal lesions. The frequencies of mucosal lesions in systemic disease groups in the elderly are shown in table (1).

The complete denture wearer group exhibited more denture-related lesions than the partial denture wearer group. Table (2) shows the prevalence of denture-related lesions among denture wearers.

The subjects who had used their dentures for ≥ 20 exhibited more denture hyperplasia than other denture wearers. Denture stomatitis was often observed in subjects wearing dentures for 6–10 years. Table (3) shows the distribution of lesions in terms of the age of the dentures.

Table (1): The percentages of mucosal lesions in some systemic diseases identified in the elderly.

Disease/disorders	Total N (%)	With mucosal lesion N (%)	Without mucosal lesion N(%)	χ^2/P -value
Cardiovascular diseases	89 (63.5)	78 (55.7)	11 (7.8)	0.001
Bone/joint disorders	74 (52.8)	65 (46.4)	9 (6.4)	0.001
Endocrine disorders	42 (30.0)	36 (25.7)	6 (4.3)	0.001
Respiratory diseases	20 (14.2)	16 (11.4)	4 (2.8)	0.001
Gastrointestinal disorders	48 (34.3)	42 (30.0)	6 (4.3)	0.001
Neuropsychiatric disorders	33 (23.5)	28 (20.0)	5 (3.5)	0.001
Hematologic disorders	14 (10.0)	12 (8.5)	2 (1.5)	0.001
Skin diseases	11 (7.8)	9 (6.4)	2 (1.4)	0.001
Genitourinary disorders	23 (16.4)	20 (14.2)	3 (2.2)	0.001

Table (2): The prevalence of denture-related lesions in complete and partial denture wearers.

	Complete denture wearer N (%)	Partial denture wearer N (%)	Total N (%)
Denture-related lesions	31 (51.7)	23 (35.4)	54 (43.2)
Denture stomatitis	26 (43.3)	47 (72.3)	73 (58.3)
Denture hyperplasia	28 (46.7)	6 (9.2)	34 (27.2)
Traumatic ulcer	13 (21.6)	8 (12.3)	21 (16.8)
Angular cheilitis	6 (10.0)	1 (1.5)	7 (5.6)
Frictional keratosis	4 (6.7)	2 (3.1)	5 (4.0)

Table (3): Distribution of denture-related lesions in terms of the age of denture

	Age of denture			
	5 years and over N (%)	6-10 years N (%)	11-20 years N (%)	years and over N (%)
Denture-related lesions	14 (25.9)	16 (29.6)	9 (16.7)	15 (27.8)
Denture stomatitis	22 (30.1)	43 (58.9)	5 (6.8)	3 (4.1)
Denture hyperplasia	5 (14.7)	9 (26.4)	8 (23.5)	12 (35.3)
Traumatic ulcer	3 (14.2)	5 (23.8)	4 (19.0)	9 (42.8)
Angular cheilitis	1 (14.2)	1 (14.2)	1 (14.2)	4 (57.1)
Frictional keratosis	1 (20.0)	1 (20.0)	1 (20.0)	2 (40.0)

DISCUSSION

The incidence of oral mucosal lesions ranged from 12% to 61.4% in the epidemiological studies related to the

prevalence of oral mucosal lesions in the elderly.^[11-20] In the present study, this incidence rate was 87.1%, similar to a study by Jainkittivong et al (83.6%).^[7] This

incidence was higher than those reported in a few studies (40.7– 45.9%).^[18-20] These variations between studies might be due to a lack of standard methods such as differences in age ranges, the sizes of the study populations, the criteria used to identify oral lesions and the types of oral mucosal lesions.

Males presented with more oral mucosal lesions than females, consistent with Mumcu *et al* and Dundar and Ilhan Kal.^[19,20] The prevalence of oral mucosal lesions was 45.9% in males and 40.1% in females in the present study. However, there was no significant difference in terms of sex. The incidence of oral mucosal lesions was higher in the >70 year age group than the other age groups, which is different from a study by Dundar and Ilhan Kal.^[20] In terms of age groups, there was a significant difference in the incidence of oral mucosal lesions. There was an increase in the incidence of lesions as the age increased in the lesion groups; this was especially observed with ulcers, tongue lesions, cancer and connective tissue lesions. Commonly observed lesions in the elderly in studies in our country and abroad vary from one study to another.^[11-20] Mumcu *et al*^[19] reported that the most common oral mucosal lesions were lingual varicosities, excessive melanin pigmentation and fissured tongues in the elderly. Fissured tongues, lingual varicosities and denture stomatitis were the most frequently identified lesions in elderly dental patients in the study of Dundar and Ilhan Kal.^[20] Lingual varicosity, fissured tongue and traumatic ulcers were the most common mucosal changes found in an elderly Thai population.^[7] Fordyce granules, herpes labialis, fissured tongue and denture stomatitis have been identified in the 65–74-year age group in Germany.^[2] Lingual varicosity, frictional keratosis and denture stomatitis were reported in Hong Kong.^[12] Hairy tongue, angular cheilitis, and lingual varicosity were the most frequently determined oral mucosal lesions in Finland.^[14]

Fissured tongue and lingual varicosity are the most common oral mucosal lesions in the elderly and the prevalence of these lesions increases with age.^[2,7,15,20] Age is an important risk factor for these two lesions.^[19-21] In the present study, these lesions were detected as the most common oral mucosal lesions.

Denture use was associated with a high prevalence of oral mucosal lesions. Most oral mucosal lesions in the elderly have been correlated with the use of removable dentures. Denture-associated lesions are the most common type of oral mucosal lesions in the aging population. The findings of oral mucosal lesions observed in denture wearers in this study were similar to those previously reported in elderly subjects.^[5,7,12,14,15,20,23]

The prevalence of oral mucosal lesions related to denture use was higher in women in some studies.^[21,23-25] In the present study, the prevalence of denture-related lesions was also higher in female subjects. This high frequency

of lesion incidence among females might be due to the fact that during or after menopause, there is atrophy of the oral mucosa, resulting in decreased protection against the chronic irritation of ill-fitting dentures.^[22]

The most common denture-related mucosal lesions observed in previous studies included traumatic ulcer, denture-induced stomatitis and denture hyperplasia.^[24,26,27] These findings are similar to our study as the most commonly observed denture-related lesion was denture stomatitis.

In agreement with several studies,^[24,25,28] in the present study there were more denture-related mucosal lesions among complete denture wearers than in partial denture wearers. In contrast to our findings, there were no significant differences in the prevalence rates of denture-related mucosal lesions between the two denture wearing groups in some studies.^[20,27]

The length of denture use has been shown to be positively related to increased denture-related mucosal lesions in previous studies.^[26,28-30] However, denture-related mucosal lesions might occur in relation to poor denture hygiene and the continuous use of dentures throughout day and night.^[31] In this study, the individuals who used their dentures for ≥ 20 years exhibited more denture hyperplasia than other denture wearers.

Epidemiological studies have shown that tongue lesions constitute a considerable proportion of oral mucosal lesions, and their prevalence rates vary in different regions of the world.^[7,32-34] In the present study, fissured tongue was the most frequent tongue lesion, similar to previous studies.^[7,19,21,33]

The oral cavity is one of the most appropriate locations for the development of oncological diseases. Oral cancer is an age-related disease and 98% of patients are >40 years old. There is a linear increase in the incidence of oral cancer with aging. Oral cancer is more common in men than women in many countries. Malignant lesions were determined in men only in the study by Dundar and Ilhan Kal and in women only in a study by Cebeci *et al.*^[20,35] In contrast to these studies, malignant lesions were determined in both women and men in our study.

The length of denture use has been shown to create a positive effect on the presence of oral mucosal lesions in the literature.^[25,28,30] The length of denture use was also a significant risk factor for having oral mucosal lesions in the current study. When dentures are used for long periods, xerostomia associated with diabetes mellitus increases the traumatic effect of dentures.^[36]

Although the majority of oral mucosal lesions detected in the present study were benign, there were patients with premalignant and malignant lesions. Therefore, it is important to carry out periodic oral examinations for the detection of precancerous and cancerous lesions,

especially in the elderly, smokers and denture users. The findings of this study support other studies that have reported positive relationships between age, sex and denture use and the presence of oral mucosal lesions.

REFERENCES

- Pindborg JJ. Oral cancer and precancer as diseases of the aged. *Community Dent Oral Epidemiol*, 1978; 6: 300–307. DOI: 10.1111/j.1600-0528.1978.tb01171.x
- Reichart PA. Oral mucosal lesions in a representative cross-sectional study of aging Germans. *Community Dent Oral Epidemiol*, 2000; 28: 390–398. DOI: 10.1034/j.1600-0528.2000.028005390.x
- Peacock ME, Carson RE. Frequency of self-reported medical conditions in periodontal patients. *J Periodontol*, 1995; 66(11): 1004–1007. doi.org/10.1902/jop.1995.66.11.1004
- Jainkittivong A, Sriwatana W. Assessment of the medical status in a dental school patient population. *CU Dent J*, 1997; 20: 35–44.
- Jainkittivong A, Aneksuk V, Langlais RP. Medical health and medication use in elderly dental patients. *J Contemp Dent Pract*, 2004; 5(1): 31–41.
- Patil S, Doni B, Maheshwari S. Prevalence and Distribution of Oral Mucosal Lesions in a Geriatric Indian Population. *Can Geriatr J*, 2015; 18(1): 11–14. doi: 10.5770/cgj.18.123
- Jainkittivong A, Aneksuk V, Langlais RP. Oral Mucosal Conditions in Elderly Dental Patients. *Oral Diseases*, 2002; 8: 218–223. DOI: 10.1034/j.1601-0825.2002.01789.x
- World Health Organization. 3rd ed. Geneva: WHO, 1995. Application of the International classification of diseases to dentistry and stomatology, ICD-DA.
- Regezi JA, Sciubba JJ, Jordan RCK. *Oral Pathology: Clinical Pathologic Correlations*. Saunders Elsevier, 2008.
- Jorge Junior J, de Almeida OP, Bozzo L, Scully C, Graner E. Oral mucosal health and disease in institutionalized elderly in Brazil. *Community Dent Oral Epidemiol*, 1991; 19: 173–175. DOI: 10.1111/j.1600-0528.1991.tb00136.x
- Vigild M. Oral mucosal lesions among institutionalized elderly in Denmark. *Community Dent Oral Epidemiol*, 1987; 15: 309–313. DOI: 10.1111/j.1600-0528.1987.tb01741.x
- Corbet EF, Holmgren CJ, Phillipsen HP. Oral mucosal lesions in 65–74-year-old Hong Kong Chinese. *Community Dent Oral Epidemiol*, 1994; 22: 392–395. DOI: 10.1111/j.1600-0528.1994.tb01599.x
- Thomson WM, Slade GD, Spencer AJ. Dental caries experience and use of prescription medication amongst people age 60+ in South Australia. *Gerodontology*, 1995; 12: 104–110. DOI: 10.1111/j.1741-2358.1995.tb00139.x
- Nevalainen MJ, Narhi TO, Ainamo A. Oral mucosal lesions and oral hygiene habits in the home-living elderly. *J Oral Rehabil*, 1997; 24: 332–337. DOI: 10.1111/j.1365-2842.1997.tb00336.x
- Lin HC, Corbet EF, Lo EC. Oral mucosal lesions in adult Chinese. *J Dent Res*, 2001; 80: 1486–1490. doi/pdf/10.1177/00220345010800052001
- García-Pola Vallejo MJ, Martínez Díaz-Canel AI, García Martín JM, González García M. Risk factors for oral soft tissue lesions in an adult Spanish population. *Community Dent Oral Epidemiol*, 2002; 30(4): 277–85. DOI: 10.1034/j.1600-0528.2002.00048.x
- Harris CK, Warnakulasuriya KA, Cooper DJ, Peters TJ, Gelbier S. Prevalence of oral mucosal lesions in alcohol misusers in south London. *J Oral Pathol Med*, 2004; 33(5): 253–9. DOI: 10.1111/j.0904-2512.2004.00142.x
- Avcu N, Ozbek M, Kurtoglu D, Kurtoglu E, Kansu O, Kansu H. Oral findings and health status among hospitalized patients with physical disabilities, aged 60 or above. *Arch Gerontol Geriatr*, 2005; 41(1): 69–79. doi.org/10.1016/j.archger.2004.11.005
- Mumcu G, Cimilli H, Sur H, Hayran O, Atalay T. Prevalence and distribution of oral lesions: a cross-sectional study in Turkey. *Oral Dis*, 2005; 11(2): 81–7. DOI: 10.1111/j.1601-0825.2004.01062.x
- Dundar N, Ilhan Kal B. Oral Mucosal Conditions and Risk Factors among Elderly in a Turkish School of Dentistry. *Gerodontology*, 2007; 53: 165–172. doi.org/10.1159/000098415
- Kovac-Kovacic M, Skaleric U. The prevalence of oral mucosal lesions in a population in Ljubljana, Slovenia. *J Oral Pathol Med*, 2000; 29(7): 331–5. DOI: 10.1034/j.1600-0714.2000.290707.x
- Mozafari PM, Dalirsani Z, Delavarian Z, Amirchaghmaghi M, Shakeri MT, Esfandyari A, Falaki F. Prevalence of oral mucosal lesions in institutionalized elderly people in Mashhad, Northeast Iran. *Gerodontology*, 2012; 29(2): 930–4. DOI: 10.1111/j.1741-2358.2011.00588.x
- Salonen L, Helldén L, Carlsson GE. Prevalence of signs and symptoms of dysfunction in the masticatory system: an epidemiologic study in an adult Swedish population. *J Cranio-mandib Disord*, 1990; 4: 241–250. DOI: 10.1111/j.1741-2358.2011.00588.x
- Jainkittivong A, Aneksuk V, Langlais RP. Oral mucosal lesions in denture wearers. *Gerodontology*, 2010; 27(1): 26–32. DOI: 10.1111/j.1741-2358.2009.00289.x
- Ercalik-Yalcinkaya S, Özcan M. Association between Oral Mucosal Lesions and Hygiene Habits in a Population of Removable Prosthesis Wearers. *J Prosthodont*, 2015; 24(4): 271–8. DOI: 10.1111/jopr.12208
- Mandali G, Sener ID, Turker SB, Ulgen H. Factors affecting the distribution and prevalence of oral mucosal lesions in complete denture wearers. *Gerodontology*, 2011; 28(2): 97–103. DOI: 10.1111/j.1741-2358.2009.00351.x

27. da Silva HF, Martins-Filho PR, Piva MR. Denture-related oral mucosal lesions among farmers in a semi-arid Northeast- ern Region of Brazil. *Med Oral Patol Oral Cir Bucal*, 2011; 16(6): 740-4. Doi:10.4317/medoral.17081
28. Coelho CM, Sousa YT, Daré AM. Denture-related oral mu- cosal lesions in a Brazilian school of dentistry. *J Oral Re- habil*, 2004; 31: 135-9. DOI: 10.1111/j.1365- 2842.2004.01115.x
29. Ferreira RC1, Magalhães CS, Moreira AN. Oral mucosal al- terations among the institutionalized elderly in Brazil. *Braz Oral Res*, 2010; 24(3): 296-302. doi.org/10.1590/S1806- 83242010000300007
30. Moskona D, Kaplan I. Oral lesions in elderly denture wear- ers. *Clin Prev Dent*, 1992; 14(5): 11-14.
31. Gendreau L, Loewy ZG. Epidemiology and etiology of den- ture stomatitis. *J Prosthodont*, 2011; 20(4): 251-60. DOI: 10.1111/j.1532-849X.2011. 00698.x
32. Campisi G, Margiotta V. Oral mucosal lesions and risk habits among men in an Italian study population. *J Oral Pathol Med*, 2001; 30(1): 22-8. DOI: 10.1034/j.1600-0714.2001. 300104.x
33. Avcu N, Kanli A. The prevalence of tongue lesions in 5150 Turkish dental outpatients. *Oral Dis*, 2003; 9(4): 188-95. DOI: 10.1034/j.1601-0825.2003. 02933.x
34. García-Pola Vallejo MJ, Martínez Díaz-Canel AI, García Martín JM, González García M. Risk factors for oral soft tis- sue lesions in an adult Spanish population. *Community Dent Oral Epidemiol*, 2002; 30(4): 277-85. DOI: 10.1034/j.1600-0528.2002.00048.x
35. Cebeci AR I, Gulsahı A, Kamburoğlu K, Orhan BK, Oztas B. Prevalence and distribution of oral mucosal lesions in an adult Turkish population. *Med Oral Patol Cir Bucal*, 2009; 1(14): 272-277.
36. Guggenheimer J, Moore PA. Xerostomia: etiology, recogni- tion and treatment. *J Am Dent Assoc*, 2003; 134(1): 61-69. doi.org/10.14219/jada.archive.2003.0018.