



ACTIVATION OF COMPLEMENT SYSTEM ALTERNATE PATHWAY BY *ANDROGRAPHIS PANICULATA*, *OCIMUM SANCTUM* AND *AZADIRACHTA INDICA*

*Balaji P, Fathima N, Lakshyajit Kakati, Jaseer K.K, Uma Krithika S and Kannan K.

Department of Pharmacy, Faculty of Engineering & Technology, Annamalai University,
Annamalainagar - 608 002, Tamil Nadu, India.

Article Received on 13/03/2015

Article Revised on 05/04/2015

Article Accepted on 28/04/2015

*Correspondence for

Author

Balaji P

Department of Pharmacy,
Faculty of Engineering &
Technology, Annamalai
University, Annamalainagar
- 608 002, Tamil Nadu,
India.

ABSTRACT

The effect of aqueous extracts of *Andrographis paniculata* (AEAP), *Ocimum sanctum* (AEOS) and *Azadirachta indica* (AEAI) on *in vitro* complement system alternate pathway (AP) activity was evaluated and compared with that of the known immunostimulant levamisole. The results suggest that there is a concentration dependent increase in the AP activity for levamisole, AEOS and AEAI. The activity exhibited by *Andrographis paniculata* was less than that of other herbs studied.

KEYWORDS: Complement system, Alternate pathway, Haemolytic activity, *Andrographis paniculata*, *Ocimum sanctum* and *Azadirachta indica*.

INTRODUCTION

Microbial infection is a very serious problem and Living beings protect themselves from infectious organisms by various defense mechanisms called as immune system. Complement system (CS) is a part of innate immune system, consist of nine proteins denoted by C1-C9 which are present in human beings and animals in inactive form.^[1, 2] These proteins can be activated by three pathways: classical pathway (CP), alternate pathway (AP) and lectin pathway (LP). The alternate pathway of the complement system is mainly innate immune system and considered as first line defense against microbial infection.^[3-5] Excessive complement activation is observed in inflammatory reactions and autoimmune disease.^[6,7] Many herbal extracts have been reported to activate the complement system Classical pathway such as *Picrorhiza scrophulariiflora*, *Tamarindus indica*, Asteraceae family.^[8-10]

Whereas there are very few reports on agents that enhance Alternate pathway activity such as Levamisole, Trichosanthin, *Aloe vera*, *Agaricus blazei* Murill.^[11-14]

Fever is mostly due to microbial infection and large number of herbal drugs is being used for fever. These herbal drugs are might be a cure for fever not only due to antipyretic action but also for the infection. Hence any plant which is used for the treatment of fever can have potential immunostimulating effect including activation of alternate pathway because it is first line defense.

Andrographis paniculata (Kalmegh) is a herbaceous plant in the family Acanthaceae and it has been reported as antibacterial, antifungal, antiviral, choleric, hypoglycemic, hypocholesterolemic, adaptogenic, anti-inflammatory, emollient, astringent, diuretic, carminative, anthelmintic, antipyretic (Dengue fever), immunosuppressant, gastric and liver tonic.^[15, 16]

Ocimum sanctum L. (Tulsi) in the family Lamiaceae and all parts of the plant are known to possess therapeutic potentials and have been used, by traditional medical practitioners, as expectorant, analgesic, anticancer, antiasthmatic, antiemetic, diaphoretic, antidiabetic, antifertility, hepatoprotective, hypotensive, hypolipidmic immunomodulatory and antistress agents. It is also been used in treatment of fever, bronchitis, arthritis, convulsions etc. Aqueous decoction of Tulsi leaves is given to patients suffering from gastric and hepatic disorders. The juice of fresh leaves is also given to patients to treat chronic fever, dysentery, hemorrhage, popular remedy for cold and dyspepsia.^[17, 18]

Azadirachta indica (Family: Meliaceae) commonly known as neem and all parts of the tree have been used medicinally for centuries. It has been used in Ayurvedic medicine for more than 4000 years due to its medicinal properties. Hot water extract of the entire plant is used for the treatment of fever, diabetes, insecticide and purgative. It also posses immunostimulant activity, hypoglycaemic activity, antiulcer effect Antifertility effect, Antimalarial activity, Antifungal activity, Antibacterial activity, Antiviral activity and Anticancer activity.^[19,20]

Under this background the present work was carried out to evaluate the effect of AP, OS and AI extracts on complement system Alternative pathway activity.

MATERIALS

Leaves of AP, OS and AI were collected from Cuddalore district and the above plants were authenticated by Dr. V. Chelladurai, Research Scientist, Botany (Scientist – C), Centre for research of Ayurveda and Siddha, Palayamkottai, Tirunelveli District, India. Voucher specimens of the plants have been deposited in the Department of Pharmacy, Annamalai University for future reference.

Drugs and Chemicals

Levamisole the standard drug was a gift sample from MMC healthcare Limited, Chennai. Veronal buffer with Mg⁺⁺ and EGTA containing 10mM barbital, 145 mM NaCl, 0.5 mM MgCl₂ and 5 mM EGTA, pH 7.4 ± 0.2] (VBS-AP buffer) was from Boston Bioproducts (Ashland, US). Anticoagulant, sodium citrate 3.8% w/v solution was from Himedia Laboratories Pvt. Ltd. Phosphate Buffer saline (PBS, pH 7.2) was from Himedia Laboratories Pvt. Ltd. India. Normal saline was from Baxter (India) Pvt. Ltd. All other chemicals used were of GR/AR grade. Human serum (HS) samples was collected from healthy volunteers.

EXPERIMENTS AND RESULTS

Preparation of Extract

Fresh plant material was, washed under running tap water to remove adhering material, dried under shade, pulverized in a mechanical grinder and passed through sieve # 40. The 25g powder of each dried leaves of *Andrographis Paniculata*, *Ocimum sanctum*, *Azadirachta Indica* was extracted separately by boiling with distilled water (1:20, w/v) for 6 hrs and then filtered. Other portions of the distilled water were added to the marc and the extraction was repeated until the last extract was colorless. The combined extract was concentrated in a rotary evaporator at a temperature not exceeding 50°C. The resulting concentrate was Lyophilised. The yield was 25.2% w/w for aqueous extract of *Andrographis Paniculata* (AEAP), 18.3% w/w for aqueous extract of *Ocimum sanctum* (AEOS), 17.45% w/w for aqueous extract of *Azadirachta Indica* (AEAI).

Preparation of 1% Rabbit erythrocytes (RbE)

Fresh Rabbit blood was collected in a sterile bottle with anticoagulant (blood: anticoagulant, 9:1 v/v), mixed and centrifuged at 2000 rpm for 10 minutes. Supernatant was removed by decantation, cells were washed twice with PBS and twice with VBS – AP buffer

(VBS/Mg₂₊/EGTA). 1% v/v suspension of RbE in VBS – AP buffer was prepared by suspending washed erythrocytes in VBS – AP buffer.

***In vitro* alternate pathway haemolytic activity** ^[21-25]

The assay was performed in flat-bottom 96-well microtitre plates (Tarsons - 941196). 5mg / ml solutions of Levamisole (Standard drug), AEAP, AEOS and AEAI, in triplicate, were prepared separately in VBS – AP buffer. Further dilutions were made in the micro-centrifuge tubes (1:4, 1:8, 1:16, 1:32, 1:64, 1:128, 1:256) with the VBS – AP buffer resulting in a final volume of 100 µL in each tube. Subsequently, 25 µL of HS was added to each tube. After incubating for 30 min at 37 °C, 25 µL RbE suspension was added to each tube and the tubes were incubated at 37 °C for 60 min. Subsequently, the tubes were centrifuged at 1000 g for 6 min. 50 µL of the supernatant was transferred to flat-bottom microtiter plate (Tarsons - 941196), mixed with 200 µL water and the absorbance was measured at 412 nm in an ELISA automatic plate reader (Multiscan EX). Controls in this assay consisted of RBC incubated in distilled water (Total lysis), RBC incubated in buffer (Blank) and the colour of HS-dilution (complement blank). The absorbance of complement blank was subtracted from absorbance values of test serum to get the corrected absorbance of test serum. Percentage haemolysis for each dilution was calculated by using the following formula:

Percentage haemolysis (y)

$$= \frac{(\text{Corrected absorbance of test serum} - \text{Absorbance of blank})}{(\text{Absorbance of Total lysis} - \text{Absorbance of blank})} \times 100$$

The concentration of drug / extract required for producing 50% haemolysis was calculated from the graph. Percentage lysis was plotted against the drug concentration and the results are given in Table 1 and Figure 1.

Table 1. *In vitro* AP activity expressed as a percentage haemolysis using VBS – AP buffer

| NHS | Percentage lysis | | | |
|-----------------------|------------------|--------------|--------------|--------------|
| | 7.93 ± 0.39 | | | |
| Concentration (µg/ml) | LEV | AEAP | AEOS | AEAI |
| 1000 | 60.58 ± 0.29 | 21.82 ± 0.11 | 53.81 ± 0.33 | 51.60 ± 0.25 |
| 500 | 53.55 ± 0.29 | 18.90 ± 0.17 | 42.40 ± 0.23 | 40.94 ± 0.23 |
| 250 | 47.37 ± 0.33 | 16.81 ± 0.12 | 37.22 ± 0.23 | 34.64 ± 0.37 |
| 125 | 40 ± 0.33 | 14.35 ± 0.18 | 32.72 ± 0.19 | 30.09 ± 0.19 |

| | | | | |
|----------------------|----------------------|--------------|-----------------------|------------------------|
| 62.5 | 31.82 ± 0.29 | 11.80 ± 0.13 | 24.33 ± 0.35 | 23.26 ± 0.37 |
| 31.25 | 23.77 ± 0.29 | 9.73 ± 0.15 | 20.72 ± 0.21 | 17.42 ± 0.29 |
| 15.625 | 16.25 ± 0.30 | 7.98 ± 0.24 | 14.33 ± 0.35 | 12.77 ± 0.35 |
| 50% hemolysis | 355.11 ± 9.72 | ● | 834.5 ± 10.30* | 873.31 ± 11.32* |

- AEAP increases the activity of Alternate pathway to lesser extent and concentration required for 50% Hemolysis couldn't been found

Values are expressed as mean ± S.E.M, n=6 in each group,

* p<0.01 AEEA and AETA when compared with Levamisole.




Figure 2. *In vitro* AP activity expressed as a percentage haemolysis using VBS – AP buffer

Statistical analysis

The data was analysed using one-way analysis of variance (ANOVA), followed by Dunnett's t – test. P<0.05 was considered significant.

DISCUSSION

Ocimum sanctum (¹⁸) and *Azadirachta indica* (¹⁹) is traditionally used for the treatment of fever. In addition to this they are reported to have immunostimulant effect. Based on these reports, AEOS and AEAI were further evaluated for their effect of complement system alternate pathway activity and the results were compared with that of levamisole. Traditionally aqueous extracts of herbs in the form of decoction is used in folklore medicines. Hence in the present study the lyophilized aqueous extracts were used and the use of organic solvent was avoided. The *in vitro* studies revealed that there is a concentration dependent increase in the AP activity for the standard immunostimulant drug levamisole (LEV), AEOS

and AEAI. The concentration required for 50% haemolysis was found to be 355.11 ± 9.72 $\mu\text{g/ml}$, 834.5 ± 10.30 $\mu\text{g/ml}$ and 873.31 ± 11.32 $\mu\text{g/ml}$ for levamisole, AEOS and AEAI respectively with VBS – AP buffer.

In the case of *Andrographis paniculata* it is reported to immunosuppressant activity (¹⁶) at the same time there are reports that *Andrographis paniculata* is used as antibacterial, antifungal (¹⁵) and in the treatment of viral infections like dengue fever.

However, AEAP slightly increased the Alternate pathway activity and increase in the activity is dose dependent. It may be due to the fact that AEAP specifically suppress certain adaptive immune system which results in autoimmune disorder stimulated excessively and stimulate innate immune system like complement system alternate pathway activity.

CONCLUSION

In the present study attempt has been made to evaluate the effect of aqueous extracts of *Andrographis paniculata*, *Ocimum sanctum* and *Azadirachta indica* on *in vitro* complement system alternate pathway activity. There was significant rise in the AP activity for AEOS, AEAI as well as levamisole even at the least dose level 15.625 $\mu\text{g/ml}$ when compared to NHS. However AEOS, AEAI exhibited less activity when compared with levamisole. AEAP exhibited dose dependent increase in the complement system alternate pathway activity though significantly less than that of the other plants studied as well as that of levamisole. There is possibility that AEAP specifically suppress certain adaptive immune system while stimulating innate immune system like complement system alternate pathway activity and the better choice for the treatment of infectious disease without causing autoimmune disorder.

Conflict of interest statement

The authors declare that there is no conflict of interest of any kind.

REFERENCES

1. Walport MJ. Complement. First of two parts. *N. Engl. J. Med*, 2001; 344: 1058–1066.
2. Anne K, Mario H, Eliska S and Mihaly J. Factor H: A Complement Regulator in Health and Disease, and a Mediator of Cellular Interactions. *Biomolecules*, (2012); 2: 46-75.
3. Morten Harboe, Ebbe Billmann Thorgersen and Tom Eirik Mollnes. Advances in assay of complement function and activation. *Advanced Drug Delivery Reviews*, 2011; 63:976–987.

4. Wen-Chao Song, Maria Rosa Sarrias, John DL. Complement and innate immunity. *Immunopharmacology*, 2000; 49:187–198.
5. Jason R Dunkelberger and Wen-Chao Song. Complement and its role in innate and adaptive immune responses. *Cell Research*, 2010; 20:34-50.
6. Dorota Tichaczek-Goska. Deficiencies and excessive human complement system activation in disorders of multifarious etiology. *Adv Clin Exp Med*, 2012; 21(1): 105–114.
7. Maria VC and Robert BS. Complement in health and disease. *Advanced Drug Delivery Reviews*, 2011; 63: 965–975.
8. Smit HF, Kroes BH, Van Den Berg AJJ, Van Der Wal D, Van Den Worm E, Beukelman CJ, Van Dijk H, Labadie RP. Immunomodulatory and anti-inflammatory activity of *Picrorhiza scrophulariiflora*. *Journal of Ethnopharmacology*, 2000; 73: 101–109.
9. Ana Paula Landi Librandi, Tais Nader Chryso stomo, Ana Elisa CSA, Carem Gledes Vargas Recchia, Sergio Akira Uyemura and Ana Isabel de Assis-Pandochi. Effect of the extract of the tamarind (*Tamarindus indica*) fruit on the complement system: Studies in vitro and in hamsters submitted a cholesterol- enriched diet. *Food and Chemical Toxicology*, 2007; 45: 1487–1495.
10. Batbayar N, Banzragch D, Inngjerdingen KT, Naran R, Michaelsen TE and Paulsen BS. Polysaccharides from Mongolian plants and their effect on the complement system: I. Polysaccharides from plants of the *Asteraceae* family. *Asian Journal of Traditional Medicines*, 2008; 3(1).
11. Jaya Kumari and Pramoda K Sahoo. Dietary levamisole modulates the immune response and disease resistance of Asian catfish *Clarias batrachus* (Linnaeus). *Aquaculture Research*, 2006; 37: 500-509.
12. Chen X, and Ma B-L. Trichosanthin, an initiator of the alternative complement activation pathway. *Clin Exp Immunol*, 1993; 93:248 252.
13. Alishahi M, Ranjbar MM, Ghorbanpour M, Peyghan R, Mesbah M and Razi jalali M. Effects of dietary *Aloe vera* on specific and nonspecific immunity of Common carp (*Cyprinus carpio*). *Journal of Veterinary Researches*, 2010; 4(3): 85 - 91
14. Shimizu S, Kitada H, Yokota H, Yamakawa J, Murayama T, Sugiyama K, Izumi H and Yamaguchi N. Activation of the alternative complement pathway by *Agaricus blazei* Murill. *Phytomedicine*, 2002; 9: 536–545.

15. Anil Kumar, Jyotsna Dora, Anup Singh and Rishikant Tripathi. A review on king of bitter (Kalmegh). *International journal of research in pharmacy and chemistry*, 2012; 2(1): 116-124.
16. Dayana J and Parameswari CS. Emerging Novel Phytochemicals as Immunosuppressants – Review. *Journal of Advanced Pharmaceutical Research*, 2013; 4(1): 33-45
17. Prakash P and Neelu Gupta. Therapeutic uses of *ocimum sanctum* linn (tulsi) with a note on eugenol and its pharmacological actions: a short review. *Indian J Physiol Pharmacol*, 2005; 49 (2) : 125–131.
18. Caroline Jeba R, Rama Vaidyanathan and Rameshkumar G. Immunomodulatory activity of aqueous extract of *Ocimum sanctum* in rat. *International Journal on Pharmaceutical and Biomedical Research*, 2011; 2(1): 2011: 33-38.
19. Imam Hashmat, Hussain Azad and Ajij Ahmed. Neem (*Azadirachta indica* A. Juss) - A Nature's Drugstore: An overview. *International Research Journal of Biological Sciences*, 2012; 1(6): 76-79.
20. Devmurari VP and Jivani N.P. Hepatoprotective Activity of Methanolic and Aqueous Extracts of *Azadirachta Indica* leaves. *International Journal of PharmTech Research*, 2010; 2(2):1037-1040.
21. Cos P, Hermans N, Van Poel B, De Bruyne T, Apers S, Sindambiwe JB, Vanden Berghe D, Pieters L and Vlietinck AJ. Complement modulating activity of Rwandan medicinal plants. *Phytomedicine*, 2002; 9: 56–61.
22. Jaqueline Dalbello Biller-Takahashi, Leonardo Susumu Takahashi, Cleni Mara Marzocchi Machado, Fabio Sabbadin Zanuzzo, Rafael Estevan Sabioni and Elisabeth Criscuolo Urbinati. Hemolytic activity of alternative complement pathway as an indicator of innate immunity in pacu (*Piaractus mesopotamicus*). *R. Bras. Zootec*, 2012; 41(2): 237-241.
23. Klerx JPAM, Beukelman, CJ, Van Dijk H and Willers JMN. Microassay for colorimetric estimation of complement activity in guinea pig, human and mouse serum. *J Immunol Methods*, 1983; 63: 215-220.
24. Mqadmi A, Zheng X, Song J, Abramowitz S, Giclas P and Yazdanbakhsh K. Prevention of complement-mediated immune hemolysis by a small molecule compound. *Biochem Biophysical Res Commun*, 2004; 325: 1465-1471.
25. Deharo E, Baelmans R, Gimenez A, Quenevo C and Bourdy G. *In vitro* immunomodulatory activity of plants used by the Tacana ethnic group in Bolivia. *Phytomed*, 2004; 11: 516-22.