

**ANTIBACTERIAL AND ANTIFUNGAL ACTIVITY OF  
MACROCYCLIC SCHIFF'S BASES AND THEIR METAL CHELATES  
WITH COBALT (II)****Mayank Pandya, N.S.Chundawat\* and K.P. Sharma**Department of Science, F.A.S.C., Mody University of Science and Technology,  
Lakshmanagarh- 332311, Rajasthan, India.

Article Received on 18/05/2015

Article Revised on 13/06/2015

Article Accepted on 05/07/2015

**\*Correspondence for  
Author****Dr. N.S.Chundawat**Department of Science,  
F.A.S.C., Mody University  
of Science and Technology,  
Lakshmanagarh- 332311,  
Rajasthan, India.**ABSTRACT**

The synthesized macrocyclic Schiff's bases and their metal chelates with cobalt (II) were screened for antibacterial and antifungal activity. The biological activities of metal chelates revealed enhanced activity as compared to corresponding ligands.


**KEYWORDS:** Macrocyclic Schiff's bases, metal chelates, antibacterial, antifungal activity.

**INTRODUCTION**


The design and synthesis of well-arranged metal-containing macrocycles is an interesting field of macrocyclic synthetic chemistry.<sup>[1,2]</sup> In addition to their particular structural features, supramolecular species formed by self-assembly of transition metals introduce many special functional and biological properties. The macrocyclic Schiff base ligands have received vital attention not only because of their pharmacological properties like antifungal, anticancer, antiviral, antibacterial character<sup>[3,4]</sup> and their mixed soft-hard donor character, versatile coordination behavior<sup>[5,6]</sup>, but also for their capacity for chemical recognition of anions and metals of biochemical, medical and environmental importance.<sup>[7,8]</sup> In case of N and C based functionalized macrocyclic ligands, the mode of metal incorporation is very much similar to that of metalloproteins in which the requisite metal is bound in a macrocyclic cavity or cleft produced by the conformational arrangement of the protein. The attachment of metal ions to proteins such as monoclonal antibodies can create new tools for use in biology and medicine.<sup>[9]</sup>

In light of above facts the present paper has been devoted to the studies of biological activities of metal chelates of cobalt (II) with macrocyclic Schiff's bases and compare them with the biological activities of pure ligands. Macrocyclic Schiff's bases chosen for this purpose are as follows.

- a. Macrocyclic Schiff's base derived from benzil and o-phenylenediamine



- b. Macrocyclic Schiff's base derived from benzil and ethylenediamine



## MATERIAL AND METHODS

### Culture and Maintenance of microorganisms

Pure cultures of all experimental bacteria and fungi were obtained from the Microbial Type Culture Collection and Gene Bank (MTCC), Institute of Microbial Technology (IMTECH), Chandigarh. The pure bacterial cultures were maintained on nutrient agar medium and fungal culture on potato dextrose agar (PDA) medium. Each bacterial and fungal culture was further maintained by subculturing regularly on the same medium and stored at 4°C before use in experiments.

For the present study following pure bacterial and fungal cultures were taken.

### Bacterial culture

S.No.	Name	Type	MTCC No.
1	<i>Pseudomonas aeruginosa</i>	Gram negative	424
2	<i>Klebsiella pneumonia</i>	Gram negative	39
3	<i>Escherichia coli</i>	Gram negative	40

### Fungal culture

S.No.	Name	MTCC No.
1	<i>Aspergillus flavus</i>	3306
2	<i>Aspergillus tubingensis</i>	2208

### Microbiological screening

Antimicrobial activities of synthesized Schiff's bases and their metal complexes were evaluated by the agar well diffusion method (Murray *et al.*)<sup>[10]</sup> modified by (Olurinola).<sup>[11]</sup>

**Agar-well diffusion method:** 5 ml aliquot of nutrient broth was inoculated with the test organism and incubated at 37<sup>0</sup>C for 24 hours. Sterile nutrient agar plates were also prepared and holes of 5 mm diameter were cut using a sterile cork borer ensuring proper distribution. The test organisms after 24 hours of incubation were spread onto separate agar plates. The chemical compounds were dissolved in DMSO at a particular concentration and poured into appropriately labeled holes using a pipette in aseptic conditions. A hole containing DMSO served as a control. The plates were left at room temperature for two hours to allow the diffusion of the sample followed by incubation at 37<sup>0</sup>C for 24 hours in inverted position. The antimicrobial activity was determined by measuring the diameter of the zone (mm) showing complete inhibition with respect to control (DMSO).

### RESULTS

In the present investigation, the inhibitory effect of synthesized Schiff's bases and their metal chelates with cobalt (II) were evaluated against both fungicidal and bacterial strains. The antimicrobial activity was determined using agar well diffusion method and summarized in Table I-IV. The activity was quantitatively assessed on the basis of zone of inhibition.

Table-I: Effect of macrocyclic Schiff's bases on bacterial growth.

Compd.	Zone of inhibition (mm)								
	100 µg/ml			500 µg/ml			1000 µg/ml		
	P.aeruginosa (424)	K.pneumoniae (39)	E.coli (40)	P.aeruginosa (424)	K.pneumoniae (39)	E.coli (40)	P.aeruginosa (424)	K.pneumoniae (39)	E.coli (40)
a	20	35	11	17	22	14	-	18	12
b	19	33	12	23	24	-	21	21	13
DMSO	-	-	-	-	-	-	-	-	-

Table-II: Effect of metal chelates on bacterial growth.

SB + Cobalt (II)	Zone of inhibition (mm)								
	100 µg/ml			500 µg/ml			1000 µg/ml		
	P.aeruginosa (424)	K.pneumoniae (39)	E.coli (40)	P.aeruginosa (424)	K.pneumoniae (39)	E.coli (40)	P.aeruginosa (424)	K.pneumoniae (39)	E.coli (40)
1	23	39	14	18	25	16	12	20	14
2	21	36	16	28	26	15	23	27	19
DMSO	-	-	-	-	-	-	-	-	-

**Antifungal activity:** The antifungal activity was performed by cup plate method at concentration 100 µg / ml, 500 µg / ml and 1000 µg / ml.

Table-III: Antifungal activity of macrocyclic Schiff's bases.

Compd.	Zone of inhibition (mm)					
	100 µg / ml.		500 µg / ml.		1000 µg / ml.	
	A.tubingensis (2208)	A.flavus (3306)	A.tubingensis (2208)	A.flavus (3306)	A.tubingensis (2208)	A.flavus (3306)
a	-	-	52	50	81	78
b	-	-	50	48	84	85
DMSO	-	-	-	-	-	--

Table-IV: Antifungal activity of metal chelates.

SB + Copper (II)	Zone of inhibition (mm)					
	100 µg / ml.		500 µg / ml.		1000 µg / ml.	
	A.tubingensis (2208)	A.flavus (3306)	A.tubingensis (2208)	A.flavus (3306)	A.tubingensis (2208)	A.flavus (3306)
1	17	13	57	57	83	80
2	13	11	52	52	88	89
DMSO	-	-	-	-	-	--

(-) Indicates No activity

## CONCLUSION

On comparing the results of biological activity, it may be concluded that metal chelates have greater inhibiting power than the free ligands against all the microbes. Although it is difficult to make out an exact structure activity relationship between the biological activity and the structure of these complexes. These results also indicate a combination of metal and ligand environment plays a vital role in increasing a degree of inhibition.

**ACKNOWLEDGEMENT:** Authors are thankful to the Dean FASC, MUST, for providing necessary research facilities in the department.

## REFERENCES

1. Fekner, T., Gallucci, J., & Chan, M. Ruffling-induced chirality: synthesis, metalation, and optical resolution of highly nonplanar, cyclic, benzimidazole-based ligands. *J. Am. Chem. Soc.*, 2004; 126(1): 223-36; BIBLIOGRAPHY \l 1033 Sheikh, R. A., Shreaz, S., Khan, L. A., & Hashmi, A. A. Synthesis, characterization and antifungal activity of transition metal. *J. Chem. Pharm. Res.*, 2010; 2(3): 274-286.
2. BIBLIOGRAPHY \l 1033 Sessler, J., Katayev, E., Pantos, G., & Ustynyuk, Y. Synthesis and study of a new diamidodipyrromethane macrocycle. An anion receptor with a high sulfate-to-nitrate binding selectivity. *Chem. Commun.*, 2004; 11: 1276-1277; BIBLIOGRAPHY \l 1033 Chandra, S., Gupta, N., & Gupta, L. Synthesis and EPR Spectral Studies of Mono- and Binuclear Cobalt(II) and Nickel(II) Complexes with New 20-Membered Dithiatetraazamacrocyclic [N<sub>4</sub>S<sub>2</sub>] Ligand. *Synth. React. Inorg. Met.-Org. Chem.*, 2004; 34(5): 919-927.
3. BIBLIOGRAPHY \l 1033 Wang, M., Wang, L., Li, Y., & Xu, Z. Antitumour activity of transition metal complexes with the thiosemicarbazone derived from 3-acetylbulliferone. *Tran. Met. Chem.*, 2001; 26(3): 307-310; BIBLIOGRAPHY \l 1033 Asiri, A., & Khan, S. Palladium(II) Complexes of NS Donor Ligands Derived from

- Steroidal Thiosemicarbazones as Antibacterial Agents. *Molecules*, 2010; 15(7): 4784-4791.
- BIBLIOGRAPHY \l 1033 Shiekh, R. A., Rahman, I. A., Malik, M. A., Masudi, S. M., & Luddin, N. Synthesis, Spectral, Electrochemical and Biological Studies of Nitrogen Donor Macrocyclic Ligand and its Transition Metal complexes. *Int. J. Electrochem. Sci.*, 2012; 7: 12829-12845; BIBLIOGRAPHY \l 1033 Sheikh, R., Shreaz, S., Khan, L., & Hashmi, A. Development and characterization of bioactive macrocyclic metal complexes, use as a potential drug . *J. Chem. Pharm. Res.*, 2010; 2(2): 172-185.
  - BIBLIOGRAPHY \l 1033 Maji, M., Chatterjee, M., Ghosh, S., Chattopadhyay, S., Wu, B., & Mak, T. Chemistry of ruthenium(II) complexes of the tridentate NNS donor methyl 2-pyridyl ketone 4-(4-tolyl)thiosemicarbazone. Isolation and structural characterisation of a novel ruthenium(II) complex containing a co-ordinated imine of an  $\alpha$ -N heterocyclic ketone. *J. Chem. Soc. Dalton Trans.*, 1999; 2: 135-140.
  - BIBLIOGRAPHY \l 1033 Sengupta, P., Dinda, R., Ghosh, S., & Sheldrick, W. Synthesis and characterization of some biologically active ruthenium(II) complexes of thiosemicarbazones of pyridine 2-aldehyde and thiophene 2-aldehyde. *Polyhedron*, 2003; 22(3): 447-453.
  - BIBLIOGRAPHY \l 1033 Labisbal, E., Sousa, A., Castineiras, A., Gracia-Vazquez, J., Romero, J., & West, D. Spectral and structural studies of metal complexes of isatin 3-hexamethyleneiminylthiosemicarbazone prepared electrochemically. *Polyhedron*, 2000; 19: 1255-1262; BIBLIOGRAPHY \l 1033 Srivastava, M., Mishra, B., & Nizamuddin. *Ind. J. Chem.*, 2001; 40B(4): 342-344.
  - BIBLIOGRAPHY \l 1033 Canadas, M., Lopez-Torres, E., Arias, A., Mendiola, M., & Sevilla, M. Spectroscopic and electrochemical properties of nickel(II), iron(III) and cobalt(II) complexes with benzilbisthiosemicarbazone. *Polyhedron*, 2000; 19(18-19): 2059-2068.
  - BIBLIOGRAPHY \l 1033 Gao, E., Sun, H., Liao, D., Jiang, Z., & Yan, S. Synthesis of and magnetic interactions in binuclear M(II) (M=Cu, Ni and Mn) complexes of macrocyclic oxamido ligands . *Polyhedron*, 2002; 21(4): 359-364.
  - BIBLIOGRAPHY \l 1033 Murray, P., Baron, E., Pfaller, M., Tenover, F., & Tenover, H. *Manual of Clinical Microbiology* (6th Ed. ed.). Washington DC; ASM Press., 1995.
  - BIBLIOGRAPHY \l 1033 Olurinola, P. A laboratory manual of pharmaceutical microbiology. Nigeria; National Institute for Pharmaceutical Research., 1996.